



A Prospective Study on the Degree of Hyponatremia in Transurethral Resection of Prostate

Maryam Fatima^{1*}, Zainab Ali¹ and Zubia Zafar¹

¹Islam Medical College, Sialkot, Pakistan.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i47A33054

Editor(s):

(1) Dr. Giuseppe Murdaca, University of Genoa, Italy.

Reviewers:

(1) Aurobinda Samal Ispat, India.

(2) Naser sabah Hussein, Al karamh teaching hospital Baghdad, Iraq.

(3) Manjeet Kumar, IGMC Shimla, India.

Complete Peer review History: <https://www.sdiarticle4.com/review-history/74973>

Original Research Article

Received 07 August 2021

Accepted 14 October 2021

Published 29 October 2021

ABSTRACT

Background: Systemic uptake of hypotonic fluid during irrigation performed at the time of Transurethral Resection of the Prostate (TURP) may cause electrolyte changes including sodium, potassium and calcium. The present study was aimed to assess the degree of sodium changes during Transurethral Resection of Prostate (TURP).

Methods: 50 males aged 54 to 86 years, presented for TURP, were enrolled in the study with ASA classification of I to IV grades. Glycine 1.5% was used with height of irrigation kept at 60 cm. One day preoperatively and one hour postoperative, levels of serum sodium were measured for all the patients.

Results: Spinal anaesthesia was given in 31 patients, whereas 19 patients were managed under general anaesthesia, with mean duration time being 72.42 ± 24.77 min. The mean size of the resected prostate was 54.82 ± 25.04 g. 58% patients developed mild hyponatremia, 4% had asymptomatic moderate hyponatremia and no patient developed severe hyponatremia or TURP Syndrome. The systemic diseases like hypertension, DM, or IHD found to be insignificant.

Conclusion: TURP was associated with a high incidence of asymptomatic mild hyponatremia. Duration of operation was one of the most important factors.

Keywords: Serum sodium; benign prostatic hyperplasia; transurethral resection of prostate; hyponatremia; TURP.

1. INTRODUCTION

From past 60 years, the benign prostatic hyperplasia (BPH) remains a common illness in men. This ailment is featured by presence of urinary incontinence, altered urinary frequency, urgency of urination, a weak and nocturnal urine flow [1,2]. BPH is characterized by increased prostate size (more than 20 g) with irritative and/or obstructive symptoms [3]. Surgical treatment of BPH involves the Transurethral resection of the prostate (TURP), which is an urological intervention, done under spinal or general anaesthesia using endoscopy. It is being advocated as the treatment of choice in 95% cases with simple prostatectomy, being more secure and effective than open prostatectomy [4-7].

After the surgical intervention, flushing and irrigation of the bladder is done using large triple lumen catheter [8]. Different irrigation solutions used are glycine, sorbitol, distilled water, and mannitol [9]. Besides irrigation, TURP procedure also involves the dilation of surface of mucosa, removing blood; and cleaning the resected pieces of prostatic tissue for visualizing the area properly.

A serious complication observed with TURP procedure is the uptake of hypotonic irrigation fluid systemically either directly in the vascular system via the prostatic venous plexus or indirectly into retroperitoneal or perivesical space [10]. This could be because of high irrigation fluid pressure that surpasses the venous pressure (1.5 kPa) [11].

In more than 20% of patients treated with TURP, TURP syndrome was observed with a complex pathophysiology requiring a multidisciplinary management [12-13]. TURP syndrome is a water intoxication that occurs iatrogenically because of augmented intake of fluid, causing hyponatremia within 15 min to 24 hours of resection [14]. Different types of signs and symptoms are reported in patients with TURP syndrome, involving fear, fatigue, lethargy, dizziness, confusion, headache, dyspnea, restlessness, nausea and vomiting, cardiovascular manifestations, arrhythmia, bradycardia and hypertension. If the condition is not detected and managed in time, TURP syndrome can manifest with exacerbated clinical symptoms like

commencement of hypotension, cyanosis, cardiac arrest and death if not early detected and promptly treated [11]. In every TURP procedure, varying amount of irrigation fluids is absorbed through venous sinuses of prostate (ranging from 20 ml/min to several litres). With diffusion of 1L of the irrigating fluid in circulation, in an hour, acute decline in the concentration of serum sodium levels (5–8 mmol/L) has been observed [15]. Acute hyponatremia is a serious condition that occurs with serum sodium concentration decreased to a level of 115–120 mEq/L [16]. It is often complicated by intravascular hemolysis and increase in serum potassium levels, increasing the risk of hyperkalemic cardiotoxicity [17]. In TURP syndrome, a combination of hyponatremia and hyperkalemia can give rise to cardiovascular complications [18].

2. MATERIALS AND METHODS

This prospective study was conducted in Islam Medical College Sialkot, Pakistan from May 2019 to April 2020. Ethical approval was obtained from the ACH Ethics and Internal Review Board (IBR) committee. 50 male patients aged 54 – 86 years presented for TURP were enrolled in the study with American Society of Anesthesiologists (ASA) classification of grade I and IV. Informed and written consent was obtained from all study subjects.

Following the fasting regimen protocol for each patient, preoperative assessment and evaluation was done after routine investigations. Patients with the history of electrolytes imbalance, impaired renal or hepatic function, heart failure, and patients undergoing treatment with diuretics were excluded from the study. Monitoring of hemodynamic parameters was carried out as per ASA standard monitoring. Systolic blood pressure (SBP), diastolic blood pressure (DBP), Heart rate (HR), oxygen saturation and respiratory rate (RR) was recorded during and after procedure. 1.5% glycine was used as irrigation fluid at 60cm height. Preoperative serum sodium was measured one day preoperatively and one hour from the end of the procedure.

Data was entered and analyzed using the Statistical Package for Social Sciences (SPSS version 21.0). The statistical significance was significant at p-value < 0.05.

3. RESULTS

The mean of all study subjects was reported to be 71.58 ± 7.76 years. The majority of the patients had grade I or III. Spinal anaesthesia was given to 31 patients (62%) and 19 (38%) patients were subjected to general anaesthesia. The mean duration of procedure was 72.42 ± 24.77 min. The mean size of the resected prostate was 54.82 ± 25.04 g with the minimum and maximum size being 21.4 g and 131 g. Blood transfusion was done in only 5 patients (10%) (Table 1).

19 patients (38%) were reported with normal range of serum sodium, 29 (58%) had mild hyponatremia and 2 patients (4%) had an asymptomatic moderate hyponatremia, with an insignificant difference ($p\text{-value} > 0.05$) statistically. No patient developed severe hyponatremia or TURP Syndrome (Table 2).

The intergroup comparison was done between normal with mild; and normal with moderate hyponatremia for various parameters like hypertension, DM, IHD, size of resected segment, type of anaesthesia and duration of surgery. It was observed that both intergroup comparisons

showed an insignificant difference statistically except for duration of surgery (Table 3).

4. DISCUSSION

TURP is a common surgical intervention being done in males aged above 60 years. At this age, males are commonly reported with various comorbidities. It has been observed that diuretics are commonly used in this age group that can result in dehydration and deficiency of essential electrolytes such as potassium, sodium, and calcium. During Endourology surgery, absorption of intravesical irrigation fluid systemically is one of the serious intraoperative complications.

In 1990, Han et al. demonstrated that the primary reason of pathogenesis of TURP syndrome is water intoxication and hyponatremia [19]. Changes in intravascular and plasma osmolar volume are characteristics of TURP syndrome. The dilutional hyponatremia has also been reported as a result from the absorption of small amounts of fluid. But, in our study, we observed that patients develop mild to moderate hyponatremia, but no patient had developed severe hyponatremia or TURP Syndrome.

Table 1. General characteristics

Variables	Mean	SD	Minimum	Maximum
Age (Yrs)	71.58	7.76	54	86
ASA				
I	11 (22%)			
II	24 (48%)			
III	13 (26%)			
IV	02 (4%)			
Types of Anesthesia				
General Anesthesia	19 (38%)			
Spinal Anesthesia	31 (62%)			
Size of Prostate	54.82	25.04	21.4	131
≤ 60 g	39 (78%)			
≥ 60 g	11 (22%)			
Blood transfusion				
Yes	5 (10%)			
No	45			
Duration of Surgery	72.42		30	120
≤ 60 minutes	22 (44%)			
≥ 60 minutes	28 (56%)			

Table 2. Grades of hyponatremia

Variables	N (%)
Normal (135 – 145)	19 (38%)
Mild (134 – 130)	29 (58%)
Moderate (129 – 125)	2 (4%)
Severe (≤ 124)	0

Table 3. HTN, DM, IHD, Size of Prostate, Type of Anesthesia and Duration of the Procedure

Variables	Normal hyponatremia (n = 19)	Mild hyponatremia (n = 29)	P-Value	Normal hyponatremia (n = 19)	Moderate hyponatremia (n = 31)
HTN					
No	11	23	0.110	11	24
Yes	8	6		8	7
DM					
No	13	24	0.245	13	25
Yes	6	5		6	6
IHD					
No	16	24	0.895	16	26
Yes	3	5		3	5
Size of resected Prostate					
≤ 60 g (38)	16	20	0.233	16	21
≥ 60 g (12)	3	9		3	10
Type of Anesthesia					
Spinal	11	18	0.772	11	20
General	8	11		8	11
Duration of Surgery					
≤ 60 minutes (46%)	13	10	0.021*	13	10
≥ 60 minutes (54%)	6	19		6	21

*p-value<0.05 is significant

Symptoms noticed with hyponatremia are dizziness, headache, nausea, dyspnea, weakness, muscle spasms and seizures. More severe symptoms are observed when level of absorption increases more than 3 litres. In 1973, Madsen et al. conducted a study and revealed that different important factors that determine the rate of absorption of irrigants: hydrostatic pressure in the prostate area, dependence on the quantity of irrigation cylindrical vessel and the pressure developed inside the bladder during the operation [20]. Similar to our study, Petrusheva AP et al. in 2015 observed significant changes in serum sodium levels by changing the sterile water volume, the procedure duration and the resected prostate volume [21].

We observed a transient increase in serum potassium that could be due to intracellular absorption. A study done by Han et al. observed a significant increase in levels of serum potassium during the absorption of sterile water intraoperatively [16,22].

In our study, we observed no significant difference in level of hyponatremia with anaesthetic techniques. It has been found in various studies that to detect early signs of TURP syndrome, regional anaesthesia is preferred over general anaesthesia. Regional anaesthesia not only decreases the chances of venous thrombosis postoperatively, but also no significant difference is observed with general anaesthesia, in relation to cognitive function, mortality and blood loss postoperatively [23].

Although water gives better visualization for hemolysis, but acute water intoxication is a major difficulty of significant assimilation. Thus, in our study, we used 1.5% of glycine for irrigation. It was preferred being less costly and has less chances to cause renal failure and hemolysis. Glycine in concentration of 2.2% is also being used in many studies. But it has more side effects than 1.5% glycine as irrigant. Transient blindness is observed with high concentration of glycine, because it is an inhibitory neurotransmitter in the retina [24].

The other commonly used irrigants is the combination of 2.7% sorbitol and 0.54% mannitol. The solutions that are less frequently used are 2.5–4% dextrose, 3% mannitol, 3.3% sorbitol, and 1% urea. These solutions are hypotonic, thus a sufficient amount of water absorption can take place, especially when irrigation is done under pressure. These solutions were reported to cause hyperglycemia and increase the intravascular volume as compared to 1.5% glycine. [10].

Our study showed a significant change in serum sodium (hyponatremia) related to the duration of the operation. Procedure more than 1 hour is associated with higher level of hyponatremia. With more than 90 minutes duration of TURP, higher morbidity and mortality was observed [25,26].

5. CONCLUSION

Our study revealed that hyponatremia is a common finding in patients undergoing TURP. We observed that duration of operation is one of the deciding factors to determine the level of hyponatremia. We advocate that sodium levels should be assessed as a mandatory investigation during long TURP procedure for early detection and management of hyponatremia.

CONSENT

Informed consent was obtained from all patients involved in the study.

ETHICAL APPROVAL

This prospective study was conducted in Islam Medical College Sialkot, Pakistan from May 2019 to April 2020. Ethical approval was obtained from the ACH Ethics and Internal Review Board (IBR) committee.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Thorpe A, Neal D. Benign prostatic hyperplasia. *The Lancet*. 2003;361(9366): 1359-1367.
2. Nickel JC. Inflammation and benign prostatic hyperplasia. *UrolClin North Am*. 2008;35(1):109-vii.
3. Garraway W, Lee R, Collins G. High prevalence of benign prostatic hypertrophy in the community. *The Lancet*. 1991;338(8765):469-471.
4. Rassweiler J, Teber D, Kuntz R, Hofman R. Complications of Transurethral Resection of the Prostate (TURP) – Incidence, Management, and Prevention. *EurUrol* 2006; 50(5): 969–80.
5. Anson K, Nawrocki J, Buckley J, Fowler C, Kirby R, Lawrence W, et al. A multicenter, randomized prospective study of endoscopic laser ablation versus transurethral resection of the prostate. *Urology* 1995; 46(3): 305–10.
6. Ali MN. The outcome of Transurethral Resection of the Prostate *J Coll Physicians Surg Pak*. 2001; 11(12):743–6.
7. Lin YH, Hou CP, Chen TH, et al. Transurethral resection of the prostate provides more favorable clinical outcomes compared with conservative medical treatment in patients with urinary retention caused by benign prostatic obstruction. *BMC Geriatr*. 2018;18(1):15.
8. Cury J, Coelho RF, Bruschini H, Srougi M. Is the ability to perform transurethral resection of the prostate influenced by the surgeon's previous experience?. *Clinics (Sao Paulo)* 2008;63(3):315-320.
9. Oestring J. E. Benign prostatic hyperplasia: medical and minimally invasive treatment options. *N Engl J Med*. 1995; 332: 99–110.
10. Hahn RG. Fluid absorption in endoscopic surgery. *Br J Anaesth* 2006;96:8–20.
11. Gravenstein D. Transurethral resection of the prostate (TURP) syndrome: A review of the pathophysiology and management. *AnesthAnalg*. 1997; 84: 438–46.
12. Okamura K, Terai A, Nojiri Y, Okumura K, Saito S, Ozawa H, et al. Evolution of common clinical path for transurethral resection of prostate (TURP). *NipponHinyokikaGakkaiZasshi*. 2007; 98(1):3-8.
13. Hawary A, Mukhtar K, Sinclair A, Pearce I. Transurethral resection of the prostate syndrome: almost gone but not forgotten. *J Endourol*. 2009 Dec; 23(12): 2013–20.
14. Hurlbert BJ, Wingard DW. Water intoxication after 15 minutes TURP. *Anesthesiology*. 1979; 50: 355–356.
15. Hahn RG. Intravesical pressure during fluid absorption in TURP. *ScandUrolNephrol*. 2000; 34: 102–8.

16. Moorthy H. K, Philip S. Serum electrolytes in turp syndrome – is the role of potassium under estimated? *Indian J Anaesth.* 2002;46(6):441–444.
17. Chen SS, Lin AT, Chen KK, Chang LS. Hemolysis in transurethral resection of the prostate using distilled water as the irrigant. *J Chin Med Assoc.* 2006 Jun;69(6):270-5.
18. Schreingraber S, Heitmann L, Weder W, Finsterer U. Are there acid-base changes during TURP? *AnesthAnalg.* 2000; 90: 946–50.
19. Hahn RG. Fluid and electrolyte dynamics during development of the TURP syndrome. *Br J Urol.* 1990; 66: 79–84.
20. Madsen PO, Naber KG. The importance of the pressure in the prostatic fossa and absorption of irrigating fluid during transurethral resection of the prostate. *J Urol.* 1973; 109: 446–52.
21. PanovskaPetrusheva A, Kuzmanovska B, Mojsova M, Kartalov A, Spirovskaa T, Shosholcheva M, TemelkovskaStevanoska M, Zdravkovska M, Dohchev S, Stankov O. Evaluation of changes in serum concentration of sodium in a transurethral resection of the prostate. *Pril (MakedonAkadNaukUmet Odd Med Nauki).* 2015;36(1):117-27.
22. Ayus JC, Arieff AI. Glycine-induced hyponatremia. *Arch Intern Med.* 1997; 157: 223–6.
23. Dobson PM, Caldicott LD, Gerrish S P, Cole JR, Channer KS. Changes in hemodynamic variables during transurethral resection of the prostate: Comparison of general and spinal anesthesia. *Br J Anaesth.* 1994; 72: 267–71.
24. Basnal S, Afzal L, Mammen K, Osahan NK, Abraham J. Transient blindness-An unusual symptom of Transurethral prostatic resection reaction-A Case report. *Indian J Anaesth.* 2002; 46: 221–3.
25. Demirel I, Ozer AB, Bayar MK, Erhan OL. TURP syndrome and severe hyponatremia under general anaesthesia. *BMJ Case Rep.* 2012 Nov 19;2012:bcr-2012-006899.
26. Boukatta B, Sbai H, Messaoudi F, Lafrayiji Z, El Bouazzaoui A, Kanjaa N. Transurethral resection of prostate syndrome: report of a case. *Pan Afr Med J.* 2013;14:14.

© 2021 Fatima et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle4.com/review-history/74973>