



Estimating Genetic Parameters for DSSAT CROPGRO-Cotton Model Calibration and Validation

G. Srinivasan ^{a*≡}, S. Pazhanivelan ^{b^o}, S. Murali Krishnasamy ^{a[#]},
N. S. Sudarmanian ^{b[†]}, S. Rajeswari ^{c[‡]} and Balaji Kannan ^{d[¥]}

^a Department of Agronomy, Tamil Nadu Agricultural University, Coimbatore -03, India.

^b Department of Remote Sensing and GIS, Tamil Nadu Agricultural University, Coimbatore -03, India.

^c Department of Cotton, Tamil Nadu Agricultural University, Coimbatore -03, India.

^d Agricultural Engineering College and Research Institute, Coimbatore -03, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2021/v33i2430764

Editor(s):

(1) Dr. Hon H. Ho, State University of New York, USA.

Reviewers:

(1) Engr. Imran Arshad, UAE.

(2) Umar shahbaz, University of Trento, Italy.

(3) Shaik Sameera, Acharya NG Ranga Agricultural University, India.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:

<https://www.sdiarticle5.com/review-history/78969>

Received 06 October 2021

Accepted 13 December 2021

Published 14 December 2021

Original Research Article

ABSTRACT

DSSAT CROPGRO-Cotton Model (version 4.7.5) was generally used to forecast the effect of climate change on productivity. The objective of this study was to calibrate and validate this model in Tamil Nadu, India for simulation of development, growth and seed cotton yield of Suraj cotton cultivars under varied planting dates viz., 28th July, 11th August, 18th August, 25th August, 8th September and 15th September. The model was calibrated with data (phenology, biomass and yield components) collected during 2019. Calibration of CROPGRO-Cotton model with genetic coefficients of cultivar Suraj for seed cotton yield (kg ha^{-1}). Simulation of days to flowering, days from planting to first pod and physiological maturity, LAI and seed cotton yield with normalized RMSE (NRMSE) values of less than 10% across all the various planting dates densities were

[≡] Ph.D. Scholar

^o Professor and Head

[#] Professor

[†] Research Associate

[‡] Professor and Head

[¥] Professor (SWCE)

*Corresponding author: E-mail: srinivasan993.sv@gmail.com;

considered excellent. Finally, we discovered that planting at the right time can mitigate many of the negative effects of fluctuating weather on cotton productivity. As a result conclude that DSSAT model will be used to make decision on cotton planting in changing climates.

Keywords: *Gossypium hirsutum*; DSSAT CROPGRO-cotton; cotton cultivars; crop growth and yield simulation.

1. INTRODUCTION

In India Cotton is the most important cash/fiber crops commonly referred to as "King of Natural Fiber and White Gold". *Gossypium spp.* plays an important part in the agricultural and textile industrial economies around the world. Cotton supplies 65 % of the textile industry's demand in India and > 70 countries widely growing cotton in tropical/subtropical climates. There are approximately 1500 mills, four million hand-looms, 1.7 million power looms and thousands of garments, hosiery, and processing units, providing employment directly or indirectly to about 45 million people [1]. Cotton was grown in areas with rainfall ranging from 600 to 2500 mm. To grow cotton crop at least 500 mm (20 inch) of water (rainfall/irrigation) is necessary in a frequent and timely pattern throughout crop growing season [2].

India is the only country in the world to grow all the four cultivated species of cotton in addition to hybrids and has the distinction of having the largest area under cotton cultivation, accounting for approximately 42 % of the world's total area under cotton cultivation, ranging from 12.5 mha to 13.5 mha. Cotton may be grown in three different agro - ecological zones viz., northern, central and southern zone. Nearly 70 % of the crop is cultivated under rainfed conditions in the central and southern regions of the country. In India during 2019-2020 (provisional), production of Cotton was 354.91 lakh bales cultivated under an area of 133.7 lakh hectares [3] (Directorate of Economics & Statistics, 2020-21).

Crop simulation models are one of the most important instruments for integrating agronomic and information sciences [4]. Through the mathematical and conceptual relationship that governs a living plant's growth in the Soil, Water, Plant, and Atmosphere continuum, these crop models made it possible to replicate a living plant. Crop simulation models can simulate the crop and interact with the environment. Agricultural growth models are useful for assessing the influence of climate change on

crop production stability under various management strategies [5]. Crop growth simulation models allows researchers to measure the impact of climate change on soil, crop growth, productivity and the long-term viability of agriculture. Cotton is extremely sensitive to adverse environmental conditions and field management [6]. Hence these technologies are used to examine yield gaps in a variety of crops and can decrease the need for costly and need more time to evaluate field trial. Crop simulation model is very beneficial because it connects crop process analysis and performance evaluation [7,8].

The Decision Support System for Agricultural Technology Transfer (DSSAT) is a crucial tool for agricultural technology transfer and the prospect for DSSAT is really valuable [9]. Using long-term weather and soil data information, it is possible to scale out short-duration field experimental results [10,11]. DSSAT model can simulate cotton development, growth, and yield under a wide range of soil, diverse meteorological parameters and agronomic management conditions [12]. This study's purpose is determine how well perform DSSAT CROPGRO-Cotton model in simulating growth and yield of Suraj cotton.

2. MATERIALS AND METHODS

DSSAT CROPGRO-Cotton model calibration was carried in the selected monitoring site of farmer's field during July, 2019 to January, 2020. The selected monitoring site is located in the Cuddalore district, Tamilnadu at 11° 32' N, 79° 8' E and at an altitude of 68 m AMSL.

2.1 Selection of Cotton variety

This experiment involved six dates of sowing (28th July, 11th August, 18th August, 25th August, 8th September and 15th September) with Suraj cotton variety which is the most commonly grown cotton variety in the study area is preferred for simulating growth and yield.

2.2 Data Collection

The data collection was guided by technical reports from the DSSAT software. Sample analysis, observations and the usage of existing data were used to create data sets. From the planting date to harvest date, daily maximum and minimum air temperature ($^{\circ}\text{C}$), precipitation (mm), relative humidity (%) and solar radiation ($\text{MJ m}^{-2} \text{day}^{-1}$) were needed to create weather file. Soil information such as soil class, texture, bulk density, organic carbon percent, sand percent, silt percent, clay percent, pH, and cation exchange capacity in the surface layer and subsurface layer were needed to create soil file in SBuild -DSSAT. Crop management data (XBuild) such as planting method, planting date, plant density, row spacing, fertilizer application, irrigation data, harvesting date, harvesting method, grain yield/ m^2 , and leaf area index were gathered. The Weather data prevailed during crop growing season in the study area was presented in the Fig. 1.

2.3 Crop Management

Cotton (*Gossypium hirsutum* L.) was sown in the *kharif* season in the study area. The Suraj cotton variety was sown evenly at one meter apart, by ridges and furrow method. Thinning was done after crop emergence with uniform plant to plant spacing of 60 cm. When the field capacity reaches to 50% soil moisture depletion, irrigation

is provided to maintain the soil moisture preferable to crop growth. The crop was fertilized with entire fertilizer dose as recommendation of TNAU crop production guide.

2.4 DSSAT CROPGRO-Cotton Model Calibration

Genotype-specific parameters (GSPs) that are specific to each cultivar in DSSAT agricultural models and it enable the model to replicate the performance of several cultivars under a variety of soil, weather, and management situations. [13]. There are three input files were created in DSSAT to run model namely,

- Weather file: Weatherman program and collected weather data
- Soil file: SBuild program and soil data
- Experimental data file: XBuild program and crop management data

The model was calibrated using collected data from selected monitoring site of farmer's field during 2019-2020 through determination of genetic coefficient for Suraj variety in DSSAT 4.7.5. Then model was validated using the data from selected monitoring site of farmer's field during 2019-2020 by relating the observed results and simulated results. The methodology of DSSAT CROPGRO-Cotton model was presented in the Fig. 2.




Fig. 1. Weather data prevailed during crop growing season in the study area




Fig. 2. Methodology of DSSAT CROPGRO-Cotton

2.5 Crop Model Validation and Test Criteria

Validation is the process of comparing model simulation results versus observations from crops that were not involved in the calibration. In addition to its ability to predict phenology and yield, a model should be carefully evaluated under a wide range of environmental variables to evaluate the performance of vital processes. Before any model can be implemented with confidence, it must first be validated. It must undergo proper validation or an assessment of the severity of the mistakes that may happen as a consequence of its use. Model validation, in its simplest form is a comparison between simulated and observed values. Several criteria were used to quantify the differences between observed and simulated data.

2.5 Statistical Approach of Model Evaluation

The root mean square error (RMSE) values indicate how much the model over or under estimate compared to observed measurements. Lower the RMSE values higher the performance of model. RMSE tests the accuracy of the model and set of RMSE values were calculated using the formulae given below (Wallach and Goffinet, 1989). A smaller RMSE means less deviation of the simulated values from the observed values and indicates better performance. If Normalized RMSE is less than 10%, the simulation is considered as excellent, good if larger than 10% but less than 20%, medium if greater than

20% and poor if it is greater than 30% [14] and Agreement percent (Jemison et al., 1994).

$$RMSE = \left[\frac{\sum_{i=1}^n (P_i - O_i)^2}{n} \right]^{1/2}$$

$$NRMSE = 100 \times \frac{RMSE}{O_i}$$

$$Agreement (\%) = 100 \times \frac{(O_i - RMSE)}{O_i}$$

Where, P - Predicted data, O - Observed data and n - the number of observations

3. RESULTS AND DISCUSSION

3.1 Calibration and Validation Results of DSSAT Model

The calibration of the DSSAT CROPGRO-Cotton model was based on phenology and yield components were recorded at the time of harvesting. The "trial-and-error" method of DSSAT v 4.7.5 was used to evaluate how the cotton cultivar's genetic coefficients will influence the phenological stages in the CROPGRO models. Improvements were made to match observed crop phenology and yield to simulated values, as well as to keep the calibrated genetic coefficient within the cultivar's predefined error ranges. The genetic coefficients obtained from the GLUE run as part of the Calibration exercise are shown in table below (Table 1). The cultivar-specific parameters generated were within the DSSAT cultivar database's range. As a result, we can utilize the genetic coefficient generated in the model application for the research area.

Table.1. Calibrated genotypic coefficients of Suraj Cotton - DSSAT CROPGRO-Cotton model

No.	Code	Description	Suraj variety
1	CSDL	Critical Short Day Length below which reproductive development progresses with no daylength effect (for shortday plants) (hour)	23.00
2	PPSEN	Slope of the relative response of development to photoperiod with time(positive for short day plants) (1/hour)	0.01
3	EM-FL	Time between plant emergence and flower appearance (R1) (photothermal days)	50.0
4	FL-SH	Time between first flower and first pod (R3) (photothermal days)	11.0
5	FL-SD	Time between first flower and first seed (R5) (photothermal days)	14.0
6	SD-PM	Time between first seed (R5) and physiological maturity (R7) (photothermal days)	49.00
7	FL-LF	Time between first flower (R1) and end of leaf expansion (photothermal days)	75.00
8	LFMAX	Maximum leaf photosynthesis rate at 30 C, 350 vpm CO ₂ , and high light (mg CO ₂ /m ² -s)	1.16
9	SLAVR	Specific leaf area of cultivar under standard growth conditions (cm ² /g)	174.0
10	SIZLF	Maximum size of full leaf (three leaflets) (cm ²)	293.0
11	XFRT	Maximum fraction of daily growth that is partitioned to seed + shell	0.70
12	WTPSD	Maximum weight per seed (g)	0.170
13	SFDUR	Seed filling duration for pod cohort at standard growth conditions (photothermal days)	35.0
14	SDPDV	Average seed per pod under standard growing conditions	30.00
15	PODUR	Time required for cultivar to reach final pod load under optimalconditions (photothermal days)	11.0
16	THRSH	Threshing percentage. The maximum ratio of (seed/(seed+shell)) at maturity. Causes seeds to stop growing as their dry weight increases until the shells are filled in a cohort.	68.0
17	SDPRO	Fraction protein in seeds (g(protein)/g(seed))	0.153
18	SDLIP	Fraction oil in seeds (g(oil)/g(seed))	0.120

Table.2. Observed and predicted Anthesis, first pod day, Physiological maturity day and yield at harvest maturity under different planting window

Day After Planting	Observed Value	Simulated Value	RMSE	NRMSE	Agreement (%)
28th July 28, 2019					
Anthesis	60	59	1	1.67	98.33
First pod day	74	73	1	1.35	98.65
Physiological maturity day	143	145	2	1.40	98.60
Yield at harvest maturity	2589	2782	193	7.45	92.55
11th August, 2019					
Anthesis	59	61	2	3.39	96.61
First pod day	74	75	1	1.35	98.65
Physiological maturity day	145	149	4	2.76	97.24
Yield at harvest maturity	2634	2735	101	3.83	96.17
18th August, 2019					
Anthesis	59	60	1	1.69	98.31
First pod day	75	74	1	1.33	98.67
Physiological maturity day	145	149	4	2.76	97.24
Yield at harvest maturity	2681	2842	161	6.01	93.99
25th August, 2019					
Anthesis	60	61	1	1.67	98.33
First pod day	74	75	1	1.35	98.65

Day After Planting	Observed Value	Simulated Value	RMSE	NRMSE	Agreement (%)
Physiological maturity day	149	150	1	0.67	99.33
Yield at harvest maturity	2650	2804	154	5.81	94.19
8th September, 2019					
Anthesis	60	62	2	3.33	96.67
First pod day	74	75	1	1.35	98.65
Physiological maturity day	144	148	4	2.78	97.22
Yield at harvest maturity	2639	2745	106	4.02	95.98
15th September, 2019					
Anthesis	58	62	4	6.90	93.10
First pod day	73	76	3	4.11	95.89
Physiological maturity day	146	147	1	0.68	99.32
Yield at harvest maturity	2688	2759	71	2.64	97.36




Fig. 3. LAI of cotton under different dates of sowing

For calibration, information for key phenological events (anthesis day, LAI, first pod day, days to physiological maturity and yield at harvest maturity) and yield-related data are needed. The model simulation was started with the initial values for similar soils in other regions have been available in the model. Banterng et al. describe the experiment, data gathering, and model calibration in detail [15].

3.2 Days to Anthesis

The CROPGRO-Cotton model reasonably simulated days taken to flowering under different dates of sowing. The RMSE for calibrated treatment for observed and simulated days to flowering of Suraj cotton variety under different sowing dates are represented in Table 2. The crop simulation model showed similar to the observed flowering days. Results from the crop simulation model evaluation showed that the crop reached flowering stage between 59 and 62

days after sowing for all the various planting dates (similar results are found by [16]).

3.3 Leaf Area Index (LAI)

The observed LAI of the Suraj cultivar between 1.85 and 2.75 at harvest whereas DSSAT CROPGRO-Cotton simulated LAI between 2.1 to 2.5 at harvest. LAI was given in Fig.3 at different dates of sowing.

3.4 Days to First Pod and Physiological Maturity Day

The crop establishment to first pod under different planting dates are compared with observed and simulated values, the agreement was range from 95.89 to 98.67 (Table 2). The model performance was good as compared in first pod day after planting. Crop physiological maturity was 143 to 149 days for observed and 147 to 150 days for simulated days in the model

and found to be more than 97 per cent agreement. Similar result was found in cotton reported by Arshad and Muhammad [17].

3.5 Yield at Harvest Maturity

The cotton yield at harvest maturity was found to be 2688 to 2589 kg ha⁻¹ (observed). The highest DSSAT simulate cotton yield at harvest maturity was found to be 2842 kg ha⁻¹ when the crop was sown during 18th August, 2019 followed by 2804 and 2782 kg ha⁻¹ with sowing dates of 25th August and 25th July, 2019, respectively. The RMSE was found to be 71 to 193 for Suraj cotton cultivar. Similar findings also reported in various crop like rice, cotton, maize, groundnut, sorghum in DSSAT model by Torre et al. [18]; Kumar et al. [19]; Venkatesan and Pazhanivelan [20], Angel et al. [21]; Deiveegan et al. [22]; Sabarinathan et al. [23], respectively.

4. CONCLUSION

The experiment was carried out to calibrate and validate the DSSAT CROPGRO-Cotton model. The model was capable to simulate all the studied parameters of different dates of sowing. The implementation of calibrated DSSAT-CROPGRO-Cotton model by optimizing crop specific parameters of Suraj cotton genotypes followed by evaluation of the model using another independent set of data showed that DSSAT CROPGRO-Cotton model performed better in comparison with simulate phenology and yield. Hence it indicates that the DSSAT CROPGRO-Cotton model can be used as decision support tool for all these optimized Suraj cotton genotype with their respective coefficients for different applications viz., optimizing dates of sowing, plant population and spacing, and fertilizer application. It can be concluded from the our findings that the evaluation of DSSAT CROPGRO-Cotton model was found good enough research tools to predict the phenological occurrence, yield and harvest index of the cotton crop in advance and model will facilitate the farmers to make broad decision on the crop management operations. DSSAT simulation model is quite useful since it connects agriculture process analysis and good performance evaluation.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Anonymous. Indian Textiles Industry Report, Ministry of Textiles, Indian Textile Journal, Department of Industrial Policy and Promotion, Press Information Bureau; 2020.
2. Doorenbos, J., Pruitt, W., 1984. Guidelines for Predicting Crop Water Requirements. (Food and Agriculture Organisation of the United Nations).
3. Annual Report 2020-21, Department of Agriculture, Cooperation & Farmers' Welfare pg. 06 Available:<http://www.agricoop.nic.in/>
4. Jung, Jinha, Murilo Maeda, Anjin Chang, Mahendra Bhandari, Akash Ashapure, Juan Landivar-Bowles. The potential of remote sensing and artificial intelligence as tools to improve the resilience of agriculture production systems. *Current Opinion in Biotechnology*. 2021;70:15-22.
5. Hoogenboom G, Tsuji GY, Jones JW, Singh U, Godwin DC, Pickering NB, Curry RB. Decision support system to study climate change impacts on crop production. *ASA Spec. Publ. No. 59. Am. Soc. Agron, Madison, WI*. 1995;51–75.
6. Sowmiya R, Sakhivel N. Performance of cotton genotype TCH 1819 to high density planting system under winter irrigated condition at the Western agroclimatic zone of Tamil Nadu. *Journal of Applied and Natural Science*. 2021;13(SI):130-134.
7. Araya A, Prasad PVV, Ciampitti IA, Jha PK. Using crop simulation model to evaluate influence of water management practices and multiple cropping systems on crop yields: A case study for Ethiopian highlands. *Field Crops Research*. 2021;260:108004.
8. Kostkova M, Hlavinka P, Pohankova E, Kersebaum KC, Nendel C, Gobin A, Olesen JE, et al. Performance of 13 crop simulation models and their ensemble for simulating four field crops in Central Europe. *The Journal of Agricultural Science*. 2021;1-21.
9. Wajid, Aftab, Khalid Hussain, Ayesha Ilyas, Muhammad Habib-ur-Rahman, Qamar Shakil, Gerrit Hoogenboom. Crop models: Important tools in decision support system to manage wheat production under vulnerable environments. *Agriculture*. 2021;11(11):1166.
10. Jones JW, Hoogenboom G, Porter CH, Boote KJ, Batchelor WD, Hunt LA, Wilkens

- PW, Singh U, Gijsman AJ, Ritchie JT. DSSAT cropping system model. *Eur. J. Agron.* 2003;18:235–265.
11. Liu L, Basso B. Linking field survey with crop modeling to forecast maize yield in smallholder farmers' fields in Tanzania. *Food Security.* 2020;12(3):537-548.
 12. Amin A, Nasim W, Mubeen M, Ahmad A, et al. Simulated CSM-CROPGRO-Cotton yield under projected future climate by SimCLIM for southern Punjab, Pakistan. *Agric, Syst.* 2018;167: 213–222.
 13. Hunt LA, Pararajasingham S, Jones JW, Hoogenboom G, Imamura DT, Ogoshi RM. GENCALC-software to facilitate the use of crop models for analysing field experiments. *Agron. J.* 1993;85: 1090–1094.
 14. Loague, Keith, Richard E. Green. Statistical and graphical methods for evaluating solute transport models: overview and application. *Journal of Contaminant Hydrology.* 1991;7(1-2): 51-73.
 15. Banterng P, Patanothai A, Pannagetch K, Jogly S, Hoogenboom G. Seasonal variation in the dynamic growth and development traits of peanut lines. *J. Agric. Sci. (Camb.).* 2003;141:51–62.
 16. Dakhore KK, Rathod AS, Kadam DR, Shinde GU, Kadam YE, Ghosh K. Prediction of Kharif cotton yield over Parbhani, Maharashtra: Combination of extended range forecast and DSSAT-CROPGRO-Cotton model. *MAUSAM.* 2021;72(3):635-644.
 17. Arshad, Muhammad Naveed. Adapting DSSAT model for simulation of cotton yield for nitrogen levels and planting dates. *Agronomy.* 2017;109(6).
 18. Torre. D, Daniel Marc G, Jay Gao, Cate Macinnis-Ng. Remote sensing-based estimation of rice yields using various models: A critical review. *Geo-spatial Information Science.* 2021;1-24.
 19. Kumar, Sagar, Ram Niwas, Khichar ML, Yogesh Kumar, Amit Singh Premdeep. Sensitivity analysis of DSSAT CROPGRO-Cotton model for cotton under different growing environments. *Indian J Ecol.* 2017;44(4):237-241.
 20. Venkatesan M, Pazhanivelan S. Estimation of Maize yield at spatial level using DSSAT crop simulation model. *Madras Agricultural Journal.* 2018;1:105.
 21. Angel M, Astrid B, Francisca S, Jaime L. Sowing date affects maize development and yield. *Agriculture.* 2019;9:67.
 22. Deiveegan M, Pazhanivelan S, Panneerselvam S, Setiyono TD. Modelling the spatial level yield variability of groundnut (*Arachis hypogea*) in rain fed districts of Tamil Nadu using crop simulation model. *Journal of Agrometeorology.* 2019;21(Special Issue-1):217-223.
 23. Sabarinathan B, Pazhanivelan S, Ragunath KP, Sivamurugan AP, Sudarmanian NS. Calibration and validation of CERES-sorghum module of DSSAT model for rabi sorghum under rainfed condition. *The Pharma Innovation Journal.* 2021;10(1):715-718.

© 2021 Srinivasan et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/78969>*