



Effect of Different Inorganic Fertilizers and Bio-Fertilizers on Growth, Yield and Quality of Sponge Gourd (*Luffa cylindrica* L.)

Niharika^{a++*}, Deepanshu^{a#}, Devi Singh^{a†}
and Amchi C. M. Marak^{a++}

^a Department of Horticulture, Naini Agricultural Institute, SHUATS, Prayagraj, Uttar Pradesh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i92593

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/104328>

Original Research Article

Received: 02/06/2023

Accepted: 05/08/2023

Published: 10/08/2023

ABSTRACT

The study titled Effect of different inorganic fertilizers and bio-fertilizers on growth, yield, and quality of Sponge gourd (*Luffa cylindrica* L.) was conducted at the Horticulture Research Farm, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj, during the *zaid* season of 2022. The experiment was designed in a randomized block design with fifteen treatment combinations replicated three times. The treatments were made out of N, P, K, Azospirillum, Azotobacter, and PSB. The results showed that the application of T₈ (75% NP + 100% K + Azotobacter + Azospirillum + PSB) was the most effective for growth, yield and quality of sponge gourd giving maximum average vine length of (278.34 cm) at harvest, maximum number of nodes per plant (15.98) at 40 DAS, minimum number

⁺⁺M.Sc. Scholar;

[#]Assistant Professor;

[†]Associate Professor;

*Corresponding author: E-mail: kumariniharika987@gmail.com;

of days to appearance of first male and female flower at (34.83) and (40.77) DAS respectively, minimum number of days to first appearance of male and female flowers in 50% plants at (38.78) and (44.72) DAS respectively, minimum number of days to first fruit picking at (46.72) DAS, highest average number of (16.17) fruits per plant, maximum fruit length and diameter of (24.33cm) and (5.05 cm) respectively, highest average fruit weight of (137.17g), maximum average fruit yield per plant of (2.22 kg), maximum yield per hectare of (30.8 tonnes), maximum total soluble solids of 5.81 (^oBrix), and maximum ascorbic acid content of (20.29 mg/100 g).

Keywords: *Sponge gourd; bio-fertilizers; growth; yield; PSB; azospirillum; azotobacter.*

1. INTRODUCTION

Cucurbits are undoubtedly among the most intriguing crops cultivated for seed. They provide a broad variety of fruit in terms of its sizes, colours, and shapes, and there doesn't seem to be any end to the new innovations that periodically surface. The crop *Luffa* is a cross-pollinated species with 26 chromosomes ($2n = 26$). The *Luffa*, also known as sponge gourd, loofah, vegetable sponge, bath sponge, or dish cloth gourd (*Luffa cylindrica* (L.) Roem syn *L. aegyptiaca* Mill), is a member of the cucurbitaceous family. India is where the *Luffa* species first appeared. *L. acutangula* still exists in wild forms in northwest India. Sometimes it is claimed that *L. cylindrica* descended from *L. acutangula*, although this does not seem possible given how drastically different the two species are from one another in terms of appearance and flowering period. Additionally, *L. cylindrica* is more common than *L. acutangula* and has a tendency to overgrow in many areas of the Tropics. The best types for fibre (sponges), which have undergone several changes, may currently be found in Japan.

Global land shortages make agricultural production limited, necessitating higher yields and better plant nutrition using fertilisers to boost output and improve soil fertility. Excessive application of inorganic fertilizers without integration of biofertilizers deteriorates the physical, chemical, and biological aspects of soil. It has a very diminishing effect on the health of plants and living organisms including the soil microbes, which has caused a severe decrease in humus content and increase in soil, air, and water pollution. Whereas application of biofertilizers in combination with inorganic fertilizers enhances the soil nutrient content more sustainably than alone use of inorganic fertilizers.

The biofertilizers or bioinoculants comprises of microorganisms which tends to multiply swiftly when applied in the soil or as seed treatment or

as seedling root dip, creating a dense population in the rhizosphere. *Azotobacter* are aerobic, nitrogen-fixing bacteria in soil, synthesizing cellular proteins using air nitrogen, providing essential nitrogen for crop plants. *Azotobacter* can successfully replace at least some of the mineral nitrogen fertilisers used in agricultural production by fixing roughly 20 kg of nitrogen per hectare annually [1]. According to one investigation, applying a mixed culture of *Azotobacter* strains could cut the demand for nitrogen fertilisers by up to 50% [2]. A bacterial genus known as *Azospirillum* is gram-negative, microaerophilic, non-fermentative, and nitrogen-fixing. It has the capacity to create various phytohormones, such as auxins, cytokinins, and gibberellins, as well as fix atmospheric nitrogen [3]. It also produce several phytohormones, including auxins, cytokinins, and gibberellins [4]. PS bacteria and plants interact to preserve soil fertility, increase crop output, and make previously unavailable P sources accessible to metabolically healthy plants. They boost plant growth by promoting biological nitrogen fixation efficiency, increasing trace element availability, and synthesizing phytohormones [5-7].

Nitrogen and Phosphorus serve as the most important nutrient in fertilization of *Sponge Gourd* and are also required in large quantities compared to other nutrients. Inorganic fertilizers along with Biofertilizers increases the production by direct nutrient supplement, nitrogen fixation, and phosphate solubilization which increases the amount of readily available nitrogen, phosphorus, and potassium for plant use without causing much harm to the environment. Thus, the aim of this investigation is to study and screen out the best effect of different treatment combinations of inorganic and biofertilizers on the growth, yield and quality of *sponge gourd*.

2. MATERIALS AND METHODS

The present investigation was carried out to study the Effect of different inorganic fertilizers

and bio-fertilizers on growth, yield and quality of sponge gourd (*Luffa cylindrica L.*). The experiment was carried out during *Zaid* season of 2022 at Horticulture Research farm, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P). Prayagraj is situated at an elevation of 78 meters above sea level at 25.87° North latitude and 81.15°E longitude. This region has a sub-tropical climate prevailing in the South-East part of U.P. with both the extremes in temperature, i.e., the winter and the summer. In cold winters, the temperature sometimes is as low as 32°F in December – January and very hot summer with temperature reaching up to 115°F in the months of May and June. During winter, frosts and during summer, hot scorching winds are also not uncommon. The experiment materials consist of hybrid TMSG-1609 which were planted with the spacing of 120x60 cm after seed inoculation with the bio-fertilizers according to the treatments. A total of fifteen treatments were tried in RBD and replicated thrice which consisted of different combinations of inorganic fertilizers and bio-fertilizers, viz., T₁: RDN (60:40:30) +FYM (20 tonnes), T₂: 75% N + 100% PK + Azotobacter, T₃: 75% N + 100% PK + Azospirillum, T₄: 75% P + 100% NK + PSB, T₅: 75% N + 100% PK + Azotobacter + Azospirillum, T₆: 75 % NP + 100% K + Azotobacter + PSB, T₇: 75% NP + 100% K + Azospirillum + PSB, T₈: 75% NP + 100% K + Azotobacter + Azospirillum +PSB, T₉: 50% N + 100% PK + Azotobacter, T₁₀: 50% N + 100% PK + Azospirillum, T₁₁: 50% P + 100% NK + PSB, T₁₂: 50% N + 100% PK + Azotobacter + Azospirillum, T₁₃: 50% NP + 100% K + Azotobacter + PSB, T₁₄: 50% NP + 100% K + Azospirillum + PSB, T₁₅: 50% NP + 100% K + Azotobacter + Azospirillum +PSB. Five plants were randomly selected for recording observations on growth, yield and quality attributing parameters viz., vine length at harvest (cm), number of nodes per plant at 40 DAS, days to appearance of 1st male flower, days to appearance of 1st female flower, days to first appearance of male flowers in 50% plants of sponge gourd, days to first appearance of male flowers in 50% plants of sponge gourd, days to first fruit picking, number of fruits per plant, fruit weight (g) using physical balance, length of fruit (cm) using measuring scale, Diameter of Fruit (cm) using vernier calipers, average fruit yield per plant (kg), yield per hectare (tonnes/hectare), TSS (°Brix) using hand refractometer, ascorbic acid (mg/100 g). The data collected

during course of investigation were subjected to statistical analysis by adopting appropriate method of analysis of variance as described by Fisher (1950).

3. RESULTS AND DISCUSSION

3.1 Growth Parameters

3.1.1 Vine length

It is revealed from Table 1 that growth characters of sponge gourd were influenced significantly due to application of different combinations of inorganic fertilizers and bio-fertilizers. As shown in Table 1 among the different treatments T₈, which consisted of 75% NP + 100% K + Azotobacter + Azospirillum + PSB recorded significantly the longest vine length of 278.34cm, whereas T₁ {RDN (60:40:30) + FYM (20 tonnes)}recorded the shortest vine length of 254.22 cm at harvest due to the combined use of inorganic fertilizers and bio-fertilizers. The highest vine length in the best treatment might be due to ready availability of nutrients, their improved absorption and translocation by plants more quickly, which resulted in higher photosynthetic activity than other treatments. Azospirillum and Azotobacter improve plant N availability and content, affecting growth characteristics. N is crucial for protein formation, protoplasm formation, cell division, and enlargement, ultimately enhancing plant growth [8]. Phosphobacteria convert unavailable phosphorus to available form during early crop growth, aiding in plant absorption of essential nutrients. They also enhance growth through biosynthesis of growth-promoting substances like vitamin B12 and Auxin [9].

The results of present investigation are in agreement with the findings of Prabhu et al., [9] in cucumber, Prasad et al., [10] in bitter gourd, Anmol and Singh (2018) in cucumber and Sonkamble et al., [11] in watermelon.

3.1.2 Number of nodes

Number of nodes varies significantly with the treatments. The maximum number of nodes 15.98 at 40 days after sowing was recorded in T₈ (75% NP + 100% K + Azotobacter + Azospirillum +PSB), whereas minimum number of nodes 10.89 was recorded in T₁ { RDN (60:40:30) + FYM (20tonnes)} shown in Table 1. The vine length is directly proportional to the number of nodes present in the plant and combined use of

inorganic fertilizers and biofertilizers show a beneficial effect on plant growth by increasing the available N, P and their uptake.

The results of present investigation are in agreement with the findings of Prasad et al., [10] in Bitter gourd.

3.2 Earliness Parameters

3.2.1 Days to appearance of first male flower

Days to appearance of 1st male flower varies significantly within the treatments. Sponge gourd required maximum number of days to appearance of first male flower i.e., 41.53 days in Table 1 when it was treated with T₁ {RDN (60:40:30) + FYM (20 tonnes)}. On the other hand plants of the plots with treatment T₈ (75 % NP + 100 % K + Azotobacter + Azospirillum +PSB) took least number of days i.e., 34.83 days. Significant and early male flower initiation was due to combination of bio-fertilizers especially PSB which increased the availability of P which is crucial for promoting earliness [9]. The results of present investigation are in agreement with the findings of Prabhu et al., [9] in cucumber and Prasad et al., [10] in bitter gourd.

3.2.2 Days to appearance of first female flower

With a significant variation plants treated with T₈ (75% NP + 100% K + Azotobacter + Azospirillum +PSB) took least number of days (40.77 days) to appearance of 1st female flower whereas T₁{RDN (60:40:30) + FYM (20 tonnes)} took the maximum of 50.02 days for the same. The earliness in appearance of female flower in Table 1 might be due to the enhanced production of growth substances like gibberellic acid, indole acetic acid, dihydrozeatin from bio-fertilizers which had positive influence on the physiological activity of plants which could assist the plants to induce female flowers [10]. The results of present investigation are in agreement with the findings of Prabhu et al., [9] in cucumber, Prasad et al., [10] in bitter gourd, Sonkamble et al., [11] in watermelon and Sahu et al., (2022) in cucumber.

3.2.3 Days to first appearance of male flowers in 50% plants of sponge gourd

Significant variation was found in the days to first appearance of male flowers in 50% plants of Sponge Gourd in Table 1. Maximum number of 45.62 days were taken by T₁ {RDN (60:40:30) +

FYM (20 tonnes)} to first appearance of male flowers in 50% plants while it took only 38.78 days by T₈ (75% NP + 100% K + Azotobacter + Azospirillum +PSB) for the same. This might be due early appearance of male flowers and better availability of nutrients by the combined use of inorganic and bio-fertilizers especially P which plays a crucial role in early induction of flowers on the plants [9].

3.2.4 Days to first appearance of female flowers in 50% plants of sponge gourd

Minimum number of 44.72 days were taken by T₈ (75% NP + 100% K + Azotobacter + Azospirillum +PSB) to first appearance of female flowers in 50% plants while it took maximum number of 54.11 days by T₁ {RDN (60:40:30) + FYM (20 tonnes)} for the same. This significant variation and earliness observed in Table 1 was favoured by the early appearance of female flowers and enhanced production of growth substances like gibberellic acid, indole acetic acid, dihydrozeatin from bio-fertilizers which had positive influence on the physiological activity of plants which could assist the plants to induce female flowers [10].

3.2.5 Days to first fruit picking in Sponge gourd

Days to first fruit picking varies significantly in Table 1 due to application of different combinations of inorganic fertilizers and bio-fertilizers. Plants treated with T₈ (75% NP + 100% K + Azotobacter + Azospirillum +PSB) took least number of 46.72 days to first harvest whereas maximum number of 57.85 days were taken by T₁{RDN (60:40:30) + FYM (20 tonnes)} for the same. Combined use of biofertilizers might have created balanced nutrition which accelerated the synthesis of chlorophyll and amino acids, there by photosynthates were translocated from leaves to fruits [12,13]. The results of present investigation are in agreement with the findings of Prasad et al., [10] in bitter gourd, Sonkamble et al., [11] in watermelon and Sahu et al.,(2022) in cucumber.

3.3 Yield Parameters

3.3.1 Number of fruits per plant

Number of fruits per plant varies significantly among the treatments. The maximum number of fruits per plant 16.17 was recorded in T₈ (75% NP + 100% K + Azotobacter + Azospirillum +PSB) whereas minimum number of fruits per

plant 9.79 was found in T₁ {RDN (60:40:30) + FYM (20 tonnes)} in Table 1. This might be due to the increased nutrient availability from the organic phosphorus through phosphobacteria and IAA from Azospirillum could have increased the various endogenous hormonal levels in the plant tissue, thereby enhancing pollen germination and tube growth, which ultimately increased the number of fruits per plant. It may also be due to higher percentage of productive flowers in these treatments [14]. The results of present investigation are in agreement with the findings of Prabhu et al., [9] in cucumber, Prasad et al., [10] in bitter gourd, Anmol and Singh (2018), Sonkamble et al., [11] in watermelon and Sahu et al.,(2022) in cucumber.

3.3.2 Fruit weight

Fruit weight varies significantly among the treatments. Heavier fruits of 137.17 g were harvested from plants treated with T₈ (75% NP + 100% K + Azotobacter + Azospirillum +PSB) whereas lighter fruits of 121.20 g were harvested from T₁ {RDN (60:40:30) + FYM (20 tonnes)} in Table 1. This significant increase in weight might be attributed to the balanced nutrition due to integration of bio-fertilizers which accelerated the synthesis of chlorophyll and amino acids, there by translocation of photosynthates from leaves to fruits causing increase in fruit weight [12,13]. The results of present investigation are in agreement with the findings of Prabhu et al., [9] in cucumber, Prasad et al., [10] in bitter gourd, Anmol and Singh (2018), Sonkamble et al., [11] in watermelon and Sahu et al., (2022) in cucumber.

3.3.3 Fruit length

Fruit length varies significantly among the treatments. Maximum fruit length of 24.33 cm were harvested from plants treated with T₈ (75% NP + 100% K + Azotobacter + Azospirillum +PSB) whereas fruits with minimum fruit length of 17.74 cm were harvested from T₁ {RDN (60:40:30) + FYM (20 tonnes)} in Table 1. This might be due to the application of inorganic sources of nitrogen in combination with bio-fertilizers (Azotobacter and PSB) which led the plant growth favourably with the production of more carbohydrates which perhaps accelerated flow of assimilates to sink and might be the reason of higher fruit length [10]. The results of present investigation are in agreement with the findings of Prabhu et al., [9] in cucumber, Prasad et al., [10] in bitter gourd, Anmol and

Singh (2018) and Sahu et al., (2022) in cucumber.

3.3.4 Fruit diameter

Fruit diameter varies significantly among the treatments. Fruits with maximum diameter of 5.05 cm were harvested from plants treated with T₈ (75% NP + 100% K + Azotobacter + Azospirillum +PSB) whereas fruits with minimum fruit diameter of 3.01 cm were harvested from T₁ {RDN (60:40:30) + FYM (20 tonnes)} in Table 1. The application of inorganic sources of nitrogen in combination with bio-fertilizers (Azotobacter and PSB) lead the plant growth favourably with the production of more carbohydrates which perhaps accelerated flow of assimilates to sink and might be the reason of higher fruit diameter [10]. The results of present investigation are in agreement with the findings of Prabhu et al., [9] in cucumber, Prasad et al., [10] in bitter gourd, Anmol and Singh (2018) and Sahu et al., (2022) in cucumber.

3.3.5 Average fruit yield per plant

Average fruit yield per plant varies significantly among the treatments. Maximum fruit yield per plant of 2.22 kg was obtained from T₈ (75 % NP + 100 % K + Azotobacter + Azospirillum +PSB) whereas a minimum of 1.19 kg was recorded in T₁ {RDN (60:40:30) + FYM (20 tonnes)} for the same in Table 1. The quick availability of plant nutrient from inorganic source, balanced C/N ratio, synthesis of auxin, growth substances, anti fungal due to inoculation of Azotobacter and conversion of insoluble phosphate to soluble form by PSB perhaps helped to increase fruit yield [10]. The results of present investigation are in agreement with the findings of Prabhu et al., [9] in cucumber, Prasad et al., [10] in bitter gourd, Anmol and Singh (2018) in cucumber and Sonkamble et al., [11] in watermelon.

3.3.6 Fruit yield per hectare

Fruit yield per hectare varies significantly among the treatments. Maximum fruit yield per hectare of 30.8 tonnes was obtained from T₈ (75% NP + 100% K + Azotobacter + Azospirillum +PSB) whereas a minimum of 16.49 tonnes was recorded in T₁ {RDN (60:40:30) + FYM (20 tonnes)} for the same in Table 1. This might be due to soil inoculation of biofertilizers which adds nitrogen from the atmosphere by fixing and solubilizing insoluble phosphates, promoting plant growth and potentially enhancing fruit yield.

Table 1. Effect of different inorganic fertilizers and bio-fertilizers on growth, yield and quality of sponge gourd

Notation	Vine length at harvest	Number of nodes at 40 DAS	Days to appearance of first male flower	Days to appearance of first female flower	Number of days to first appearance of male flowers in 50% plants	Number of days to first appearance of female flowers in 50% plants	Days to first fruit picking	Number of fruits per plant	Fruit Weight	Fruit Length	Fruit Diameter	Fruit Yield per plant	Yield per hectare	TSS	Ascorbic acid content
T ₁	254.22	10.89	41.53	50.02	45.62	54.11	57.85	9.79	121.20	17.74	3.01	1.19	16.49	3.74	15.35
T ₂	260.89	12.11	39.00	45.3	43.04	49.34	52.28	12.07	122.87	20.01	3.64	1.48	20.6	4.36	18.27
T ₃	262.22	12.35	38.63	44.97	42.66	49	51.79	12.32	123.83	20.17	3.72	1.53	21.19	4.51	18.43
T ₄	258.98	11.09	40.07	47.03	44.15	51.11	55.28	11.10	122.17	18.40	3.14	1.36	18.83	3.98	17.67
T ₅	264.89	13.11	38.46	44.63	42.47	48.64	51.36	13.21	125.73	20.47	4.11	1.66	23.07	4.74	18.89
T ₆	267.11	14.46	37.99	43.2	41.96	47.17	49.89	14.64	129.43	21.07	4.61	1.90	26.32	5.41	19.66
T ₇	266.89	14.24	38.20	43.63	42.18	47.61	50.49	14.32	128.77	20.97	4.56	1.84	25.61	5.22	19.47
T ₈	278.34	15.98	34.83	40.77	38.78	44.72	46.72	16.17	137.17	24.33	5.05	2.22	30.8	5.81	20.29
T ₉	260.33	11.96	39.40	45.4	43.45	49.45	52.35	11.78	122.67	19.98	3.53	1.45	20.07	4.22	18.07
T ₁₀	259.89	11.71	39.73	45.83	43.79	49.89	53.16	11.54	122.60	19.97	3.36	1.41	19.65	4.13	17.94
T ₁₁	259.78	11.35	39.77	46.54	43.84	50.61	54.09	11.32	122.50	18.89	3.23	1.39	19.26	4.06	17.83
T ₁₂	263.22	12.78	38.55	44.86	42.57	48.88	51.59	12.88	125.33	20.41	3.84	1.61	22.42	4.65	18.64
T ₁₃	266.44	13.90	38.26	43.97	42.25	47.96	50.95	13.97	127.13	20.70	4.34	1.78	24.67	5.01	19.21
T ₁₄	265.02	13.54	38.38	44.37	42.38	48.37	51.20	13.76	126.77	20.65	4.20	1.74	24.23	4.96	19.00
T ₁₅	268.42	15.14	37.76	42.86	41.72	46.82	49.49	14.87	133.87	22.54	4.78	1.99	27.65	5.52	19.71
F- test	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S
SE.d (±)	4.33	0.24	0.75	0.65	0.75	0.65	0.5	0.32	1.45	0.71	0.08	0.04	0.61	0.06	0.09
CD	8.88	0.49	1.54	1.33	1.54	1.33	1.03	0.66	2.97	1.46	0.17	0.09	1.25	0.13	0.19
C.V ^{0.05}	2.01	2.25	2.38	1.77	2.16	1.62	1.19	3.05	1.41	4.26	2.53	3.29	3.29	1.61	0.6

The results of present investigation are in agreement with the findings of Prabhu et al., [9] in cucumber, Prasad et al., [10] in bitter gourd, Anmol and Singh (2018), Sonkamble et al., [11] in watermelon and Sahu et al., (2022) in cucumber.

3.4 Quality Parameters

3.4.1 TSS

TSS varies significantly among the treatments. Maximum TSS value of 5.81 °Brix was obtained from T₈ (75% NP + 100% K + Azotobacter + Azospirillum +PSB) whereas a minimum TSS value of 3.74 °Brix was recorded in T₁ {RDN (60:40:30) + FYM (20 tonnes)} in Table 1. This may be brought on by the speedy metabolic conversion of pectin and starch into soluble molecules, as well as the swift movement of sugars from leaves to growing fruits due to integration of inorganic fertilizers with bio-fertilizers. The results of present investigation are in agreement with the findings of Sayed et al., (2015) in cucumber, Das et al.,(2009)in bottle gourd, Anmol and Singh (2018) in cucumber.

3.4.2 Ascorbic acid

Ascorbic acid varies significantly among the treatments. Maximum vitamin –C content 20.29 mg per 100 g was obtained from T₈ (75% NP + 100% K + Azotobacter + Azospirillum +PSB) whereas a minimum 15.35mg per 100g was recorded in T₁ {RDN (60:40:30) + FYM (20 tonnes)} for the same in Table 1. This might be due the integration of inorganic fertilizers with bio-fertilizers which led to the speedy metabolic conversion of starch into ascorbic acid. The results of present investigation are in agreement with the findings of Sayed et al., (2015) in cucumber, Das et al., (2009)in bottle gourd, Anmol and Singh (2018) in cucumber.

4. CONCLUSION

From the present investigation, it is concluded that the influence of different inorganic fertilizers and bio-fertilizers with different treatment combinations rendered their significant effect on, growth, yield and quality of sponge gourd. Treatment T₈ (75% NP + 100% K + Azotobacter + Azospirillum +PSB) recorded best in vine length at harvest (cm), number of nodes per plant at 40 DAS, days to appearance of 1st male flower, days to appearance of 1st female flower, days to first appearance of male flowers in 50%

plants of sponge gourd, days to first appearance of male flowers in 50% plants of sponge gourd, Days to first fruit picking, number of fruits per plant, fruit weight (g), length of fruit (cm), diameter of fruit (cm), average fruit yield per plant (kg), yield per hectare (tonnes/ hectare), TSS (°Brix), ascorbic acid (mg/100 g fruit wt.).

ACKNOWLEDGEMENT

The author is thankful to the Advisor and Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj(U.P.) India for providing necessary facilities for undertaking this research. This manuscript has not received any grant from any sources.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kizilkaya R. Nitrogen fixation capacity of Azotobacter spp. strains isolated from soils in different ecosystems and relationship between them and the microbiological properties of soils. J. Environ. Biol. 2009; 30(1):73-82.
2. Romero-Perdomo F, Abril J, Camelo M, Moreno-Galván A, Pastrana I, Rojas-Tapias D, Bonilla R. Azotobacter chroococcum as a potentially useful bacterial biofertilizer for cotton (*Gossypium hirsutum*): Effect in reducing N fertilization. Revista Argentina de microbiologia. 2017;49(4):377-383.
3. Dobereiner J, Marriel IE, Nery M. Ecological distribution of Spirillum lipoferum Beijerinck. Canadian Journal of Microbiology. 1976;22(10):1464-1473.
4. Reynders L. Conversion of tryptophan to indoleacetic acid by Azospirillum Brasilense; 1979.
5. Ponmurugan P, Gopi C. *In vitro* production of growth regulators and phosphatase activity by phosphate solubilizing bacteria. African Journal of Biotechnology. 2006; 5(4):348-350.
6. Wani P, Khan M, Zaidi A. Co-inoculation of nitrogen-fixing and phosphate-solubilizing bacteria to promote growth, yield and nutrient uptake in chickpea. Acta

- Agronomica Hungarica. 2007;55(3):315-323.
7. Zaidi A, Khan MS, Ahemad M, Oves M. Plant growth promotion by phosphate solubilizing bacteria. Acta Microbiologica et Immunologica Hungarica. 2009;56(3):263–284.
 8. Bakly SA. Effect of fertilization treatments on the yield of chryslar imperial rose plants. Agricultural Research Review; 1974.
 9. Prabhu M, Natarajan S, Srinivasan K, Pugalendhi L. Integrated nutrient management in cucumber. Indian J. Agric. Res. 2006;40(2):123-126.
 10. Prasad PH, Mandal AR, Sarkar A, Thapa U, Maity TK. Effect of biofertilizers and nitrogen on growth and yield attributes of bitter melon (*Momordica charantia* L). In Proceedings, International Conference on Horticulture. 2009;738-739.
 11. Sonkamble AM, Mapari A, Patil SR, Tayade. Effect of organic manures and biofertilizers on growth and yield of watermelon (*Citrullus lanatus* Thunb.) International Journal of Agricultural and Applied Sciences. 2022; 3(2):41-45.
 12. Eifediyi EK, Remison SU. Growth and yield of cucumber (*Cucumis sativus* L.) as influenced by farmyard manure and inorganic fertilizer. J Plant Breed Crop Sci. 2010;2(7):216-20.
 13. Moharana DP, Mohan L, Singh BK, Singh AK, Kumar H, Mahapatra AS. Effect of Integrated Nutrient Management on growth and yield attributes of Cucumber (*Cucumis sativus* L.) cv. Swarna Ageti under polyhouse conditions. The Bioscane. 2017; 12(1):305-8.
 14. Rajagopal V, Rao IM. Changes in the endogenous level of auxins and gibberellin-like substances in the shoot apices of nitrogen-deficient tomato plants (*Lycopersicon esculentum* Mill). Australian Journal of Botany. 1974;22(3):429-435.

© 2023 Niharika et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/104328>