

Asian Journal of Research and Reports in Gastroenterology

Volume 6, Issue 1, Page 127-131, 2023; Article no.AJRRGA.107129

Distal Esophageal Squamous Cell Carcinoma: A Rare, but Possible Localization

Livia Marieta Negoita a* and Catalin Stefan Ghenea a

^a Clinical Department of Gastroenterology, Bucharest Emergency Clinical Hospital, Romania.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

https://www.sdiarticle5.com/review-history/107129

Received: 29/07/2023 Accepted: 04/10/2023 Published: 10/10/2023

Case Report

ABSTRACT

Aim: To report a rare but possible localization of a distal esophageal squamous cell carcinoma. **Case Presentation:** A 71-year-old man with severe progressive dysphagia and epigastric pain for 2 months. Endoscopy revealed normal esophagus up to 37 cm from the dental arch where a tight stenosis that prevents the endoscope from advancing is visualized. The histopathological result revealed squamous cell carcinoma. Tumor Board decided to initiate neoadjuvant chemoradiotherapy followed by surgery.

Conclusion: Management of patients with esophagus tumors is challenging. Localization of squamous cell carcinoma in the distal esophagus, although rarer, is possible.

Keywords: Esophageal squamous cell carcinoma; dysphagia; biopsy; esophageal stent; esophagectomy.

*Corresponding author: Email: negoita.livia@yahoo.com;

ABBREVIATIONS

AJCC: American Joint Committee on Cancer

CDDP: Cisplatin

EADC: Esophageal adenocarcinoma

ESCC: Esophageal squamous cell carcinoma

EGJ: Esophagogastric junction EUS: Endoscopic ultrasound CT: Computer tomography

5-FU: Fluorouracil

1. INTRODUCTION

"Esophagus cancer is the ninth most common malignancy and ranks as the sixth most frequent cause of death cancer in the world, constituting 7% all gastrointestinal cancers"[1]. of "Esophagus cancer exists in two main forms with different ethiological and pathological characteristics: esophageal squamous (ESCC) and carcinoma esophageal adenocarcinoma (EADC)"[2]. "ESCC is the most common form of esophageal cancer worldwide and the most prevalent type of oesophageal cancer in the East (Asian and Eastern European countries)"[3]. "Primary risk factors oesophageal cancer development (squamous cell carcinoma) are smoking and alcohol consumption"[4]. "Other risk factors include: previous treatment due to squamous cell carcinoma of the head and neck or lung, oesophageal post caustics burning, oesophageal achalasia. Plummer-Vinson syndrome genetically-dependent hand-foot hyperkeratosis"[5]. "ESCC is located mainly in the upper and middle parts of the oesophagus, while EADC is usually diagnosed in the lower part. ESCC occurs in 30% of the distal esophagus"[2]. "In the AJCC 8th edition, tumors involving the EGJ that have midpoint within the proximal 2 cm of the cardia/proximal stomach are staged as esophageal cancers"[6].

2. CASE PRESENTATION

2.1 Clinical History

A 71-year-old man presented to our clinic with severe progressive dysphagia for solid food felt substernal, epigastric pain and weight loss (5 kg) for 2 months. He's a smoker (50 cigarettes/years) and mild alcohol intake (60 grams alcohol per week). Diagnostic evaluation included a complete medical history, physical examination, laboratory blood analyzes and pulmonary and cardiac function tests." with

"Diagnostic evaluation included a complete medical history, physical examination, laboratory blood tests and pulmonary and cardiac function tests. All laboratory exams were normal, except the lower level of albumin 29 g/L. The patient underwent esophagogastroscopy with biopsies, chest-abdomen and pelvis CT with contrast enhancement and EUS. Endoscopy revealed normal esophagus up to 37 cm from the dental arch where a tight stenosis (Fig.1) that prevents the endoscope from advancing is visualized. A Savary guide is inserted and dilated with Savary disposable bougie dilator of 10, 12 and 14 Fr. After that, multiple biopsies are taken from the lower esophagus (Fig. 2) and in retroversion, from the level of cardia and finally, a metallic esophageal stent is inserted (Fig. 3) to allow the patient to feed.




Fig. 1. Cardia stenosis




Fig. 2. Biopsies from the lower esophagus

The histopathological result revealed esophageal mucosa with infiltrative carcinoma tumor proliferation (Fig. 4) and immunohistochemistry established the diagnosis of squamous cell

carcinoma (CK7 negative: CK56 cytoplasmic positive in tumor tissue: p63 positive nuclear in tumor cells). A chest CT revealed a stenotic circumferential parietal thickening affecting the distal esophagus for a length of ~35-40 mm. Also, two lymph node images at the lower periesophageal level and the abdominal-pelvic CT examination did not show suggestive images of a secondary determination. EUS revealed a tumor infiltration of muscularis propria. A search screening for other diseases did not show any. The tumor stage was determined according to the TNM staging system (pT2-N1-M0). Being an advanced local esophageal cancer, in a patient with good performance status (ECOG 0), the Tumor Board decided to initiate neoadjuvant chemo-radiotherapy (two cycles of CDDP and 5-FU) followed by surgery.




Fig. 3. Metalic esophageal stent




Fig. 4. Invasive squamous cell carcinoma (100x)

After surgery, the patient will be monitored clinically and imaging once every 6 months in the first three years and endoscopically every 4-6 months in the first year. Also, being a squamous cell carcinoma of the esophagus, the patient will perform an ENT examination and an annual bronchoscopy in order to detect a second cancer.

3. DISCUSSION

"The esophagus cancer is a relatively uncommon but extremely lethal malignancy"[7]. "The most common presenting symptoms are dysphagia, odynophagia and weight loss"[8]. "The disease most commonly spreads to adjacent and supraclavicular lymph nodes, liver, lungs, pleura, and bone"[9]. The diagnosis of esophageal cancer is based on the following methods: the patient's history and clinical examination, radiological examination (classic and tomography computerized), digestive endoscopy biopsy/cytological examination and endoscopy[10]. "Squamous cell carcinomas and adenocarcinomas cannot be distinguished radiographically or endoscopically"[11].

diagnosis of certainty esophageal carcinoma is establishes by digestive endoscopy histopathological examination. appears to have a more favorable prognosis with non-surgical available and surgical interventions compared to EADC"[12]. Surgery is the standard treatment for a candidate with a localized, non-superficial tumor[13]. "In patients with resectable. locally unresectable borderline unresectable cancer without contraindications to surgery, it is important to consider preoperative radiochemotherapy"[14].

In this case with advanced esophageal cancer, the endoscopy easily diagnosed the esophagus cancer. For a smoker and long-term alcohol consumption, like the patient of this case, the risk of esophageal SCC is increased and this tumor should be always considered during the initial medical assessment. Multidisciplinary approach was used for treating this patient. "The use of preoperative chemotherapy and radiation therapy followed by esophageal resection appears to prolong survival as compared with controls in small, randomized trials and some reports suggest that no additional benefit accrues when surgery is added if significant shrinkage of tumor has been achieved by the chemoradiation combination"[15].

4. CONCLUSION

The prognosis for patients with esophageal carcinoma is poor. Localization of squamous cell carcinoma in the distal esophagus, although rarer, is possible. Surgical treatment consists of resecting the tumor by esophagectomy with esogastric/jejunal or eso-colic anastomosis. accompanied bγ lymph node dissection (mediastinal and coronary gastric). Intervention is difficult, requiring an approach double (both thoracic and abdominal), and should be performed in experienced services. Patients with locally advanced disease should also receive preoperative chemoradiation or perioperative chemotherapy.

5. LIMITATIONS OF THE STUDY

The limits of our study are represented by the lack of follow-up after chemotherapy and surgical intervention of the patient.

CONSENT

The informed consent was obtained from the patient for publication and any accompanying images.

ETHICAL APPROVAL

The treatment strategy/study protocol was approved by a local tumor board/ethics committee.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- 1. CQ. Liu et al, Epidemiology of esophageal cancer in 2020 and projections to 2030 and 2040. Thorac. Cancer. 2023;14(1):3–11.
 - DOI:10.1111/1759-7714.14745
- Jain S. Dhingra S, Pathology of esophageal cancer and Barrett's esophagus, Ann. Cardiothorac. Surg. 2017;6(2). DOI:10.21037/acs.2017.03.06
- 3. Zhang HZ, Jin GF, Shen HB, Epidemiologic differences in esophageal cancer between Asian and Western populations. Chinese Journal of Cancer. 2012;31(6).

- DOI:10.5732/cic.011.10390
- 4. Yang X. et al, Smoking and alcohol drinking in relation to the risk of esophageal squamous cell carcinoma: A population-based case-control study in China. Sci. Rep. vol. 2017;7(1):17249. DOI: 10.1038/s41598-017-17617-2
- Tarazi M, Chidambaram S, Markar SR, Risk factors of esophageal squamous cell carcinoma beyond alcohol and smoking, Cancers. 2021;13(5).
 DOI: 10.3390/cancers13051009
- Zhang S, Orita H, Fukunaga T, Current surgical treatment of esophagogastric junction adenocarcinoma. World Journal of Gastrointestinal Oncology. 2019;11(8). DOI:10.4251/wigo.v11.i8.567
- 7. Marom G, Esophageal cancer staging, Thoracic Surgery Clinics. 2022;32(4). DOI: 10.1016/i.thorsurg.2022.06.006
- 8. Ueda H. et al. Clinical evaluation of palliative chemoradiotherapy for metastatic esophageal cancer, Oncotarget. 2017;8(46).

DOI:10.18632/oncotarget.17925

- 9. M. Watanabe et al, Recent progress in multidisciplinary treatment for patients with esophageal cancer, Surgery Today. 2020;50(1).
 - DOI: 10.1007/s00595-019-01878-7
- Vasile MI. et al, Update on Endoscopy-Based Imaging Techniques in the Diagnosis of Esophageal Cancer. Curr. Heal. Sci. J. 2017;43(4):295–300. DOI:10.12865/CHSJ.43.04.01
- 11. Abbas H, Ghazanfar H, UI Hussain AN, Baiomi Α, Ihimoyan Α, atypical presentation of esophageal squamous cell Carcinoma masquerading as diffuse Case Esophagitis. severe Rep. Gastroenterol. 2021;15(2). DOI:10.1159/000517129
- Schlottmann F, Casas MA, Molena D, Evidence-based approach to the treatment of esophagogastric junction tumors, World J. Clin. Oncol. 2022;13(3): 159–167.
 - DOI:10.5306/wjco.v13.i3.159
- Kalff MC, Van Berge Henegouwen MI, 13. Gisbertz SS. Textbook outcome for esophageal cancer surgery: An international consensus-based update of a quality measure. Dis. Esophagus. 2021:34(7).
 - DOI:10.1093/dote/doab011
- Jin Z, Zhu K, Sun JJ. Zhang, Zhang B, Minimally invasive versus open

esophagectomy after neoadjuvant therapy for esophageal cancer: a metaanalysis. J. Cardiothorac. Surg. 2023; 18(1):90.

DOI:10.1186/s13019-023-02180-x

 Lordick F, Hölscher AH, Haustermans K, Wittekind C, Multimodal treatment of esophageal cancer. Langenbeck's Archives of Surgery. 2013;2:398. DOI: 10.1007/s00423-012-1001-1

© 2023 Negoita and Ghenea; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
https://www.sdiarticle5.com/review-history/107129