



# Assess the Magnitude of Genetic Diversity in Advance Breeding Line of Mungbean with Respect to Seed Yield and Component Traits

Pallavi Sonaniya <sup>a\*</sup>, Lekha Ram <sup>a</sup>, Gundlapalli Soumith <sup>a</sup>  
Ravindra Solanki <sup>a</sup>, Ghanshyam Bamaniya <sup>a</sup>  
and Rahul Sonaniya <sup>b</sup>

<sup>a</sup> Department of Genetics and Plant Breeding, Rajmata Vijayaraje Scindia Krishi Vishwa Vidyalaya, Gwalior Madhya Pradesh-474001, India.

<sup>b</sup> Department of Genetics and Plant Breeding, Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur, Madhya Pradesh-482004, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/JSRR/2024/v30i52013

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/113653>

Original Research Article

Received: 07/02/2024  
Accepted: 11/04/2024  
Published: 19/04/2024

## ABSTRACT

The study was conducted across four distinct environments in Madhya Pradesh during the 2021 kharif season, utilizing a Randomized Complete Block Design involving fourteen Mungbean genotypes with three replications. Examination of genetic parameters unveiled a notable pattern: the phenotypic coefficient of variation (PCV) consistently exceeded the genotypic coefficient of variation (GCV) across all observed traits of particular interest were the traits demonstrating the highest PCV and GCV values, notably seed yield per plant in E2 followed by biological yield per

\*Corresponding author: E-mail: [pallavisonaniya.gpb@gmail.com](mailto:pallavisonaniya.gpb@gmail.com);

plant in E1. These findings strongly suggest the prevalence of additive gene action influencing these traits, as indicated by their high heritability estimates. The traits with the highest heritability values were seed yield per plant in E3 and E4, biological yield per plant in E2 and E4, and number of pods per plant in E1. These results underscore the genetic basis underlying these traits and their potential for targeted breeding efforts. Cluster I was the largest among all the clusters comprising 8 genotypes, whereas cluster II had 5 genotypes. While the clusters III were solitary cluster consisting one genotype only. Cluster II showed maximum intra cluster D2 value, whereas clusters III showed zero value for Intra cluster distance. The highest inter cluster divergence was observed between genotypes of cluster II and III. The percent contribution of individual characters toward the total divergence was found high for Harvest index, whereas Number of primary branches showed low percentage of contribution

**Keywords:** PCV; GCV; heritability; genetic advance; D2; clusters; environments.

## 1. INTRODUCTION

Mungbean, scientifically known as *Vigna radiata* (L.) R. Wilczek var *radiata*, is a legume from the Fabaceae family. With a chromosome count of  $2n = 22$  and a compact 579 Mb genome, it goes by various names like green gram, moong, green soy, green bean, mash bean, and golden gram [1]. This crop, thriving in tropical and subtropical regions, stands as a significant food and cash crop. Its seeds offer easily digestible dietary protein, with an ideal daily intake of about 40 grams per person or 14.6 kg annually [2].

The protein content in green gram surpasses that of cereals by two to three times, comprising 51 percent carbohydrates, 26 percent protein, and 4 percent each of minerals and essential vitamins like A, B1, B2, C, niacin, folate, iron, calcium, and zinc. This nutrient profile complements and diversifies cereal-based diets effectively.

Mungbean has gained importance in double and intercropping systems due to its short growing cycle and nitrogen-fixing capabilities ( $58-109 \text{ kg-1 ha-1}$ ), which significantly enhance soil fertility [3]. Its agronomic, nutritional, and economic advantages have led to a substantial surge in both production and consumer demand worldwide over the last two decades.

Globally, Mungbean cultivation spans various latitudes and seasons, occupying over 6 million hectares. In the context of pulses in India, which covers 28.79 million hectares with a production of 25.46 million tonnes and a productivity rate of 885 Kg per hectare, Mungbean covers 5.55 million hectares, yielding 3.17 million tonnes at a productivity rate of 570 Kg per hectare [4]. Notably, its production has escalated from 1.60 million tonnes in 2015-16 to 3.17 million tonnes in 2021-22. Rajasthan, Maharashtra, Karnataka, Andhra Pradesh, and Madhya Pradesh stand out

as key Mungbean cultivating states. Among these, Madhya Pradesh contributes 938.10 hectares, 1134.52 tonnes, and 1209 Kg per hectare in terms of area, production, and productivity, respectively.

Enhancing the genetic traits of this crop primarily relies on understanding genetic variability and heritability factors. Analyzing parameters like phenotypic and genotypic coefficients of variability, genetic advance, and heritability [2] becomes imperative. This knowledge aids in pinpointing the most favorable yield attributes for selection or hybridization, laying the foundation for effective crop improvement.

Genetic diversity is an important factor and also a prerequisite in any hybridization programme. Inclusion of diverse parents in hybridization programme serves the purpose of combining desirable recombinations. Multivariate analysis by means of Mahalanobis  $D^2$  statistic is a powerful tool in quantifying the degree of divergence at genotypic level.

## 2. MATERIALS AND METHODS

The study took place in the experimental areas of the All India Coordinated Research Project on MULLaRP across four diverse environments in Madhya Pradesh (R.A.K. College of Agriculture, Sehore; K.V.K. Barwani; K.V.K. Jhabhua; and College of Agriculture, Gwalior) during the 2021 Kharif season. Fourteen different mungbean genotypes were cultivated using a Randomized Complete Block Design with three replications. The crop rows spanned 4 meters in length, with a spacing of 30 cm between rows and 10 cm between plants. The fields exhibited uniformity, gentle slopes, proper drainage, and normal fertility levels, where all recommended agronomic practices were implemented to foster a robust crop.

Data collection involved observing five randomly selected plants within each plot. These observations encompassed various characteristics such as days to 50% flowering, days to maturity, plant height, number of branches per plant, number of pods per plant, pod length, number of seeds per pod, 100-seed weight, biological yield per plant, seed yield per plant, and harvest index.

Analysis of variance followed the methodology outlined by Burton [5], while the estimation of range was conducted based on Johnson et al.'s method [6]. The Phenotypic Coefficient of Variation (PCV) and Genotypic Coefficient of Variation (GCV) were calculated using Burton's formula [5]. Heritability was determined using Allard's formula [7], and genetic advance was calculated as a percentage using Johnson et al.'s formula [6]. The data were subjected to Mahalanobis  $D^2$  statistics as per Mahalanobis [8] method and genotypes were grouped into different clusters following Toucher's method as suggested by Rao (1952).

### 3. RESULTS AND DISCUSSION

Results of the present study on fourteen genotypes were done to understand the genetic diversity. The experimental results of the present investigation have been mentioned under following:

#### 3.1 Analysis of Variance

The analysis of variance highlighted significant differences among genotypes across most traits, with exceptions noted in specific environments for traits like days to maturity in E3, plant height in E3 and E4, and the number of primary branches and seeds per pod across all environments. When pooling data across environments, significant differences among genotypes were observed for most traits, except for the number of primary branches and seeds per pod, where highly significant differences in mean sum of squares were evident, likely due to minimal genotype  $\times$  environment interaction for these specific traits within the studied material (Table 1). These findings align with previous research by Sopan et al. [9] and Mwangi et al [10].

#### 3.2 Parameters of Genetic Variability

The genetic variability parameters namely phenotypic coefficient of variation (PCV), genetic coefficient of variation (GCV), heritability in broad sense (%), genetic advance and expected

genetic advance (as per cent of mean) for all ten traits were Estimated and have been presented in Table 2.

#### 3.3 Phenotypic Coefficient of Variation (PCV), Genetic Coefficient of Variation (GCV)

In this study, the phenotypic coefficient of variation (PCV) consistently surpassed the genotypic coefficient of variation (GCV) across all traits analysed. The traits with the highest PCV and GCV were seed yield per plant in E2 (31.36, 28.58) followed by biological yield per plant in E1 (27.41, 24.80). Moderate PCV and GCV values were observed for biological yield per plant in E4 (19.35 for PCV) followed by number of primary branches in E4 (16.15) and harvest index in E3 (14.61). For GCV, notable values were recorded for seed yield per plant in E4 (18.29), biological yield per plant in E3 (17.06), and number of pods per plant in E1 (16.42).

The traits with the lowest PCV were number of pods per plant in E4 (9.64), followed by 100 seed weight (8.39), and days to maturity in E1 (3.41). As for GCV, the lowest values were observed for plant height in E4 (1.31), number of primary branches in E1 (3.93), and plant height in E2 (8.02).

These results suggest that selecting traits with higher PCV and GCV for further breeding efforts could be more effective. These findings align with previous research by Nitesh et al. (2017) for seed yield per plant, harvest index, and number of pods per plant, Tushar kumar et al. (2019) and Mariyammal et al. [11] for seed yield per plant, Ramakrishnan et al. [12], Zida et al. [13], and Sineka et al. [14] for number of pods per plant.

#### 3.4 Heritability (Broad Sense) and Genetic Advance

Heritability, indicating the inheritance of traits from parents to offspring, aids breeders in selecting superior genotypes. Higher heritability suggests traits less influenced by the environment and primarily controlled by additive genetic effects. Robinson et al. (1949) categorized heritability into high (>60%), moderate (30-60%), and low (<30%). In this study, seed yield per plant displayed the highest heritability in E3 (97.32) and E4 (92.95), followed by biological yield per plant in E2 (88.66) and E4 (87.91), and number of pods per plant in E1 (63.66). Moderate heritability was noted for days to 50% flowering in E1 (59.59) and plant height in E3 (47.21) (Table 2).

**Table 1. Analysis of variance for ten various characters in mungbean**

Source Variations	of	df	Mean Squares								
			Days to 50% flowering					Days to maturity			
			E1	E2	E3	E4	POE	E1	E2	E3	E4
Replicate	2	1.88	1.45	3.50	4.57	8.73	6.02	13.02	0.73	7.73	7.06
Genotypes	13	9.83*	6.85*	8.11*	6.79*	20.67*	8.60*	13.42*	10.30	12.08*	16.11*
Error	26	3.31	2.01	1.55	1.57	0.88	2.94	4.15	6.02	2.68	2.23
Source Variations	of	df	Mean Squares								
			No. of pods/plant					No. of primary branches.			
			E1	E2	E3	E4	POE	E1	E2	E3	E4
Replicate	2	0.16	8.00	18.50	5.42	8.32	0.73	0.28	0.30	0.30	0.18
Genotypes	13	36.19*	25.57*	41.15*	7.87*	199.21*	0.71	1.51	0.72	1.86	17.45
Error	26	5.80	4.25	6.98	1.76	7.92	0.50	0.90	0.54	1.15	0.48
Source Variations	of	df	Mean Squares								
			Plant height					No. of seeds per pod			
			E1	E2	E3	E4	POE	E1	E2	E3	E4
Replicate	2	1.59	0.28	1.73	8.00	7.52	0.73	0.21	0.28	0.16	0.25
Genotypes	13	58.45*	83.03*	14.79	7.42	140.84*	1.12	0.58	1.06	1.06	15.94
Error	26	22.26	18.33	7.96	5.53	1.36	0.66	0.39	0.61	0.83	0.38
Source Variations	of	df	Mean Squares								
			Biological yield per plant					Harvest index			
			E1	E2	E3	E4	POE	E1	E2	E3	E4
Replicate	2	1.28	3.50	5.07	3.42	1.30	26.84	7.94	18.93	28.48	18.72
Genotypes	13	38.85*	43.40*	28.61*	29.47*	82.00*	30.42*	31.65*	141.5	32.14*	161.57*
Error	26	2.67	2.65	2.49	3.24	3.99	11.23	14.54	12.22	10.08	25.16
Source Variations	of	df	Mean Squares								
			100 seed weight					Seed yield per plant			
			E1	E2	E3	E4	POE	E1	E2	E3	E4
Replicate	2	0.02	0.38	0.10	0.04	0.06	0.39	0.009	0.08	0.17	0.11
Genotypes	13	0.18*	0.16*	0.20*	0.24*	5.99*	2.02*	3.90*	2.64*	1.02*	15.06*
Error	26	0.02	0.04	0.05	0.02	0.05	0.12	0.24	0.12	0.06	0.17

**Table 2. Estimates of genetic parameters for 10 different characters of mungbean genotypes**

Sr. No.	Characters	Environment	Mean	Range		PCV (%)	GCV (%)	h <sup>2</sup> (broad sense)	Genetic advance	Gen. Adv as % of Mean
				Mini	Maxi					
1	Daysto50%flowering	E1	41.95	39.00	45.00	5.58	3.52	59.59	3.93	9.37
		E2	40.45	38.66	43.00	4.70	3.14	67.43	3.59	8.88
		E3	41.92	39.00	44.00	4.61	3.53	78.51	4.77	11.38
		E4	41.42	39.00	43.66	4.39	3.18	72.55	4.05	9.78
		POE	41.44	39.75	43.75	4.85	2.75	65.16	2.77	6.68
2	Daystomaturity	E1	64.28	62.00	69.00	3.41	2.14	59.07	3.65	5.68
		E2	65.47	62.00	68.33	4.11	2.69	62.67	4.83	7.38
		E3	65.61	63.00	69.33	4.15	1.82	59.16	2.25	3.43
		E4	65.45	61.00	67.66	3.68	2.70	73.82	5.45	8.33
		POE	65.20	62.00	68.25	3.86	1.95	60.64	2.75	4.22
3	No.ofpodspersplant	E1	19.38	13.00	25.00	20.59	16.42	63.56	10.55	54.44
		E2	21.14	17.33	26.00	15.94	12.61	62.54	8.79	41.58
		E3	24.35	20.00	31.66	17.59	13.86	61.98	11.05	45.38
		E4	20.21	16.00	22.66	9.64	7.06	53.63	4.41	21.82
		POE	21.27	18.25	23.75	16.53	7.25	55.64	2.89	13.59
4	No.ofprimarybranchesper plant	E1	6.62	6.00	7.00	11.45	3.93	56.83	0.47	7.10
		E2	6.64	6.00	8.00	15.82	6.79	63.39	0.91	13.70
		E3	5.74	4.66	6.66	13.52	4.35	50.28	0.45	7.84
		E4	7.31	6.00	8.00	16.15	6.68	62.07	0.95	13.00
		POE	6.58	5.91	7.25	14.57	3.55	66.98	0.35	5.32
5	Plantheight(cm)	E1	57.05	49.33	63.33	10.27	6.09	60.15	8.59	15.06
		E2	57.93	50.33	64.66	10.90	8.02	79.05	14.17	24.46
		E3	60.69	55.33	65.00	5.27	2.49	47.21	3.03	4.99
		E4	60.50	56.00	62.66	4.10	1.31	40.18	1.15	1.90
		POE	59.04	53.91	62.08	8.06	3.80	57.24	4.47	7.57
6	No.ofseedsperspod	E1	10.33	9.00	11.33	8.74	3.82	49.09	0.83	8.03
		E2	9.57	9.00	10.33	7.07	2.65	43.97	0.51	5.33
		E3	9.71	8.66	10.66	9.02	3.99	49.51	0.81	8.34
		E4	8.83	8.00	10.00	10.80	3.14	43.46	0.45	5.10


Sr.	Characters	Environment	Mean	Range	PCV	GCV	h <sup>2</sup> (broad	Genetic	Gen.Adv	
		POE	9.61	9.00	10.33	8.94	3.83	48.27	0.75	7.80
7	Biologicalyieldperplant(gm)	E1	14.00	10.73	20.60	27.41	24.80	86.83	9.82	70.11
		E2	16.42	12.00	23.66	24.52	22.44	88.66	10.52	64.07
		E3	17.29	12.66	21.83	19.35	17.06	82.73	8.14	47.05
		E4	15.71	12.00	23.00	22.03	18.82	87.91	7.91	50.35
		POE	15.85	12.40	22.23	23.20	18.81	84.62	10.07	63.53
8	Harvest index (%)	E1	21.48	17.40	28.50	19.54	11.77	71.28	6.37	29.66
		E2	23.31	17.83	29.70	19.29	10.24	63.18	5.33	22.87
		E3	24.54	18.73	26.96	14.61	3.27	40.05	0.83	3.38
		E4	20.10	15.96	26.73	20.76	13.49	77.16	7.35	36.57
		POE	22.36	18.96	26.31	18.46	8.06	60.07	3.35	14.98
9	100seedweight(gm)	E1	2.96	2.56	3.36	9.66	7.74	74.63	0.85	28.72
		E2	3.51	3.26	3.93	8.39	5.64	55.83	0.65	18.52
		E3	2.95	2.40	3.53	11.68	8.86	67.98	0.93	31.53
		E4	2.60	2.16	3.00	12.20	10.43	83.27	1.05	40.38
		POE	3.01	2.71	3.25	10.36	5.30	67.22	0.43	14.29
10	Seedyieldperplant(gm)	E1	2.97	1.93	4.40	29.19	26.77	94.13	2.36	79.46
		E2	3.86	2.20	5.40	31.36	28.58	93.05	3.22	83.29
		E3	4.21	2.96	5.76	23.26	21.74	97.32	2.75	65.32
		E4	3.09	2.26	4.00	20.08	18.29	92.95	1.70	55.02
		POE	3.53	2.47	4.82	26.71	21.37	81.05	1.97	55.81

Genetic advance, influenced by selection intensity, heritability, and phenotypic standard deviation, ranged from 0.12 to 4.98 across environments. The highest genetic advance was seen in biological yield per plant in E2 (10.52) and number of pods per plant in E3 (11.55), while the lowest was observed in 100 seed weight in E2 (0.65). Other traits had genetic advance values around 7.19 (biological yield per plant in E4), 9.59 (plant height in E1), 7.35 (harvest index in E4), 4.77 (days to 50% flowering in E3), 1.70 (seed yield per plant in E4), and 0.95 (number of primary branches in E4) (Table 2). These findings align with previous research by Aparna et al. [15] highlighting high heritability and low genetic advance for days to 50% flowering and traits like biological yield per plant and number of pods per plant, as observed by Malli et al. [16] and Sineka et al. (2021).

### 3.5 Genetic Divergence Analysis

To calculate  $D^2$  values, the correlated average values of characteristics were converted into standard uncorrelated averages through the application of Tocher's method. The statistical distance (Mahalanobis  $D^2$ ) between pairs of genotypes was determined by summing up the squared differences between the pairs of corresponding uncorrelated values for any two genotypes analysed simultaneously.

The analysis of variance showed significant differences between Mungbean genotypes for all the characters studied. All the fourteen genotypes were grouped into three clusters (Table 3 and Fig. 1). Cluster I was the largest among all the clusters comprising 8 genotypes, whereas cluster II had 5 genotypes. While the clusters III were solitary cluster consisting one genotype only.



Cluster No.	No. of genotypes	Name of the Genotypes
I	8	RVSTM 22-1, RVSTM 22-2, IPM 410-3 (Shikha), IPM 205 -7 (Virat), RVSM 18-1, MI 98-64, MI 181-1, MI 750-1
II	5	RVSM 22-3, RVSM 22-4, RVSM 22-5, RVSM 22-6, RVSM 22-7
III	1	RVSM 22-8

Fig. 1. Dendrogram based on genetic distance, summarizing the data on differentiation between 14 Mungbean genotypes according to Mahalanobis'  $D^2$  method

Table 3. Clustering pattern of 14 genotypes of Mungbean based on Mahalanobis'  $D^2$ -values and the member present in each respective cluster

1= RVSTM 22-1	8= RVSM 22-8
2= RVSTM 22-2	9= RVSM 18-1
3= RVSM 22-3	10= MI 98-64
4= RVSM 22-4	11= MI 181-1
5= RVSM 22-5	12= MI 750-1
6= RVSM 22-6	13= IPM 410-3(Shikha)
7= RVSM 22-7	14= IPM 205-7 (Virat)

The pattern of group constellations indicated that significant variability existed among the genotypes as observed from the clusters. Cluster I originating from different places indicated that there was no parallelism between clustering pattern and geographic distribution of genotypes. Similar findings were reported by Henry and Mathur (2017) and Rahim [17]. This kind of genetic diversity was recorded among the genotypes belonging to the same geographic origin might be due to difference in adoption, selection criteria, selection pressure and environmental condition.

The intra and inter cluster D<sup>2</sup> mean values are presented in table 4. On the basis of D<sup>2</sup> values, 14 genotypes were grouped into three clusters. Intra cluster distance ranged from 0.00 to 4.98. Cluster II showed maximum intra cluster D<sup>2</sup> value (D<sup>2</sup> = 4.98), cluster I (D<sup>2</sup> = 4.71), whereas cluster III showed zero value for intra cluster distance. The highest inter cluster divergence was observed between genotypes of cluster II and III (8.46), followed by cluster I and cluster II (8.15). Cluster distance was lowest between cluster I and cluster III (7.20).

Cluster II showed highest cluster mean for six characters viz., Days to 50% flowering, Days to

maturity, No. of pods/plant, Plant height, Biological yield per plant and Seed yield per plant. Cluster III recorded highest mean value for No. of primary branches, No. of seeds per pod and Harvest index, while cluster I recorded highest mean value for 100 seed weight only (Table 5).

The selection and choice of parents mainly depend upon contribution of character towards divergence Loganathan *et al.*, [18] and the contribution towards genetic divergence is represented in Table 6. It was observed that among all the traits, contribution of seed yield per plant was maximum. The percent contribution of individual characters toward the total divergence was found high for Harvest index (%) (34.07%) followed by 100 seed weight (g) (20.88%), Biological yield per plant (g) (17.58%), Seed yield per plant (g) (10.99%), Plant height (cm) (4.40%), Days to 50% flowering and Number of pods per plant (3.30%), Days to maturity and Number of seeds per pod (2.20%) and Number of primary branches (1.10%) showed low percentage of contribution and it also contributed towards total divergence. Similar results were reported by Appalaswamy and Reddy [16] and Henry and Mathur [20].

**Table 4. Average intra (Bold) and inter cluster D<sup>2</sup> values in Mungbean genotypes**

<b>Clusters</b>	<b>Cluster I</b>	<b>Cluster II</b>	<b>Cluster III</b>
Cluster I	4.71	8.15	7.2
Cluster II		4.98	8.46
Cluster III			0.00

**Table 5. Cluster means for yield attributed characters in Mungbean genotypes**

<b>Clusters</b>	<b>Days to 50% flowering</b>	<b>Days to maturity</b>	<b>No. of pods/plant</b>	<b>No. of primary branches</b>	<b>Plant Height (cm)</b>
Cluster I	41.54	63.75	18.17	6.46	53.88
Cluster II	43.07	65.07	22.00	6.80	61.6
Cluster III	39.67	64.67	16.00	7.00	59.67
<b>Clusters</b>	<b>No. of seeds per pod</b>	<b>Biological yield per plant (gm)</b>	<b>Harvest index</b>	<b>100 seed weight (gm)</b>	<b>Seed yield per plant (gm)</b>
Cluster I	10.38	11.91	20.41	3.07	2.41
Cluster II	10.20	17.99	21.81	2.79	3.87
Cluster III	10.67	10.73	28.50	2.97	3.03



**Table 6. Relative contribution of different characters for genetic divergence in Mungbean genotypes**

S. No.	Character	Contribution %
1	Days to 50% flowering	3.30
2	Days to maturity	2.20
3	No. of pods/plant	3.30
4	No. of primary branches	1.10
5	Plant height	4.40
6	No. of seeds per pod	2.20
7	Biological yield per plant	17.58
8	Harvest index	34.07
9	100 seed weight	20.88
10	Seed yield per plant	10.99
	<b>Total</b>	<b>100%</b>

It is well known that crosses between divergent parents usually produce greater heterotic effect than closely related ones [21-24]. Considering the importance of character towards total divergence, the present study indicated that parental lines selected from cluster I (RVSTM 22-1, RVSTM 22-2, IPM 410-3 (Shikha), IPM 205 -7 (Virat), RVSM 18-1, MI 98-64, MI 181-1, MI 750-1) and from cluster II (RVSM 22-3, RVSM 22-4, RVSM 22-5, RVSM 22-6, RVSM 22-7) could be used in crossing programme to achieve desired segregants [25-27].

#### 4. CONCLUSIONS

The pooled analysis of variance shows significance across all genotypes, except for the number of primary branches, signifying ample genetic variability within the population. Traits such as seed yield per plant and biological yield per plant exhibited the highest phenotypic and genetic coefficient of variation, suggesting a strong influence of genetic variability on their expression. The combination of high heritability and substantial genetic advance as a percentage of the mean was observed prominently in biological yield per plant, followed by seed yield per plant and the number of pods per plant. This suggests that these traits are predominantly governed by additive gene action, making them suitable candidates for direct selection in breeding programs. On the basis of these traits superior genotypes are selected and used in hybridization programme as a donor parent. crossing programme could be made among the genotypes belonging in cluster I and cluster II for getting maximum heterotic combinations, especially for yield of Mungbean.

#### ACKNOWLEDGEMENT

I am immensely grateful to Dr. Lekharam, our esteemed guide and chairman, a Scientist in Plant Breeding and Genetics at R.A.K. College of Agriculture, Sehore (M.P.), for initiating this research problem and providing invaluable guidance, unwavering support, and insightful suggestions throughout this investigation. His inspiration was instrumental in the successful completion of this dissertation.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Rahangdale S, Lakhani JP, Singh SK, Sharma S, Barela A, Kumar P, Prajapati SS. Study of Selection Criteria and Genetic Variability in Mungbean (*Vigna radiata* L. Wilczek) Genotypes for Seed Yield and its Contributing Traits. *Environment and Ecology*. 2023;41(1):145—151.
- Afroz M, Thirumala Rao V, Sridhar V, Hari Y. Assessment of genetic variability in mung bean [*Vigna radiata* (L.) Wilczek.] *The Pharma Innovation Journal*. 2022;11(11):1371-1373
- Haften SV, Dudley C, Kang Y, Smith D, Nair RM, Douglas CA, Potgieter A, Robinson H, Hickey LT, MR. Building a better Mungbean: Breeding for reproductive resilience in a changing climate. *Food and Energy Security*; 2023. Available: <https://doi.org/10.1002/fes3.467>

4. Anonymous. Report of All India Coordinated Research Project on KHARIF PULSE, ICAR-Indian Institute of Pulses Research, Kanpur. 2022-23;42-45.
5. Burton GW. Quantitative Inheritance in grasses. Proceeding of 6th International Grassland Congress. 1952;(1):277-283.
6. Johnson HW, Robinson HF, Comstock PE. Estimate of genetic and environmental variability in Soybeans. Agron. J. 1955;47:314-318.
7. Allard RW. Principles of Plant breeding. John Wiley and Sons, Inc., New York. 1960;485.
8. Mahalanobis PC. On the generalized distance in statistics. Sankhyā: The Indian Journal of Statistics, Series A. 2008;80:S1-S7.
9. Sopan Z, Gandhi AG, Vishwavidyalaya K, Sao IA, Nair IS, Nanda HC, Nair SK. Combining ability analysis for seed yield, its contributing traits and protein content in Mungbean (*Vigna radiata* (L.) Wilczek). International Journal of Chemical Studies. 2018;6(5):761-764.
10. Mwangi JW, Okoth OR, Kariuki MP, Piero MN. Genetic and phenotypic diversity of selected Kenyan mung bean (*Vigna radiata* L. Wilczek) genotypes. Journal of Genetic Engineering and Biotechnology. 2021;19:142.
11. Mariyammal I, Pandiyan M, Vanniarajan C, Kennedy JS, Senthil N. Genetic variability in segregating generations of green gram (*Vigna radiata* L. Wilczek) for quantitative traits. Electronic Journal of Plant Breeding. 2019;10(1):293 – 296.
12. Ramakrishnan CK Divya, Savithramma DL, Vijayabharathi A. Studies on genetic variability, correlation and path analysis for yield and yield related traits in Green gram [*Vigna radiata* (L.) Wilczek]. Int. J. Curr. Microbiol. App. Sci. 2018;7(3):2753-2761.
13. Zida WPFMS, Batiemo TB. J, Ouedraogo TJ, Sawadogo M. Agro morphological evaluation of 44 lines of Mung Bean (*Vigna radiata* (L.) Wilczek) introduced in Burkina Faso. European Scientific Journal, ESJ. 2021;17(40), 20.
14. Sineka T, Murugan E, Sheeba A, Hemalatha G, Vanniarajan C. Genetic relatedness and variability studies in greengram (*Vigna radiata* (L.) Wilczek) Electronic Journal of Plant Breeding. 2021;12(4):1157 – 1162.
15. Aparna R, Singh SK, Sharma V, Pathak R. Genetic variability, heritability, genetic advance and path analysis in mungbean (*Vigna radiata* (L.) Wilczek). Legume Res. 2015;38(2):157-163.
16. Malli SK, Lavanya GR, Nikhil BSK. Genetic variability, heritability and genetic advance in mungbean (*Vigna radiata* L. Wilczek) genotypes. Green Farming. 2018;9(2): 235-238,
17. Rahim MA. Genetic variability, character association and genetic divergence in mungbean (*Vigna radiata* L. Wilczek). Plant Omics. 2020;3(1):1-6.
18. Loganathan P.K. Saravanan and J. Ganesan. Genetic analysis of yield and related components in green gram (*Vigna radiata* L.). Research on Crops. 2021;1:34-36.
19. Appalaswamy A, Reddy GLK. Genetic divergence and heterotic studies on mungbean (*Vigna radiata* L. Wilczek). Legume Res. 2020;27(2):115-118.
20. Henry A, Mathur, BK. Correlation studies of quality parameters, seed yield and maturity period in clusterbean. J. Arid Legumes. 2017;5(1):70-74.
21. Asari T, Patel BN, Patel R, Patil GB, Solank C. Genetic variability, correlation and path coefficient analysis of yield and yield contributing characters in mung bean [*Vigna radiata* (L.) Wilczek]. International Journal of Chemical Studies. 2019;7(4): 383-387.
22. Desai, V. K., Parmar, L. D., Chaudhary, A. R. and Chaudhary, N. B. (2020). Genetic variability, correlation, path coefficient and stability analysis for yield and its attributing traits in summer Green gram [*Vigna radiata* (L.) Wilczek] accessions. Int. J. Curr. Microbiol. App. Sci. 2020;9(6): 2942-2955.
23. Jagdhane NM, Suresh BG, Ram BJ, Yadav P. Genetic variability and character association for seed yield in Mungbean [*Vigna radiata* (L.) Wilczek]. Journal of Pharmacognosy and Phytochemistry. 2017;6(4):1388-1390.
24. Mehendi S, Singh IP, Bohra A, Singh CM. Multivariate analysis in greengram (*Vigna radiata* (L.) Wilczek). Legume Res. 2018; 38(6):758-762.
25. Ovir, Singh AK. Analysis of morphological characters interrelationship in the germplasm of mungbean (*Vigna radiata* (L.) Wilczek) in the hot arid climate.; Legume Res. 2016;39(1):14-19.
26. Paliwal S, Sharma S, Pathak N. Principal component analysis in mungbean [*Vigna radiata* L. Wilczek] genotypes under two

- seasons The Pharma Innovation Journal. 2022;11(3):1500-1505.
27. Payasi DK. Genetic variability analysis for seed yield and its components in mungbean (*Vigna radiata* (L.) Wilczek). Inter. J. Plant Breed. Genet. 2015;9:177-188

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

*<https://www.sdiarticle5.com/review-history/113653>*