



The Association between Depression, Daytime Sleepiness, Chronotype and Fatigue among the Students

**Nishi Soni ^a, Saurabh Jaiswal ^a, Sudhir Kumar ^a,
Shalie Malik ^a and Sangeeta Rani ^{a*}**

^a Department of Zoology, University of Lucknow, Lucknow-226007, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajmah/2024/v22i71061>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/119138>

Original Research Article

Received: 20/04/2024

Accepted: 23/06/2024

Published: 27/06/2024

ABSTRACT

Fatigue and excessive daytime sleepiness (EDS) are frequently conflated in everyday life, but they are separate concepts with distinct symptoms. Despite their potential to coexist or occur independently, their interplay with depression remains largely unexplored. This study investigates the complex relationship between fatigue, excessive daytime sleepiness (EDS), and depression among students, aiming to unravel their relative associations. A cross-sectional survey of Indian school students (N=450) aged 11 to 20 (15.2±1.46) was analysed. The study used self-reported measures such as the Fatigue Severity Scale (FSS), Epworth Sleepiness Scale (ESS), and the Center for Epidemiologic Studies Depression Scale (CES-D). The findings revealed significant associations between depression and various combinations of sleepiness and fatigue, with the E⁺F⁺ group showing the strongest correlation. Depression was significantly higher in excessive daytime sleepers (N= 165, 69.9%) (P < 0.001) and fatigue (N= 167, 70.8%) (p < 0.001). Depression was

*Corresponding author: E-mail: sangeetarani6219@gmail.com;

Cite as: Soni, Nishi, Saurabh Jaiswal, Sudhir Kumar, Shalie Malik, and Sangeeta Rani. 2024. "The Association Between Depression, Daytime Sleepiness, Chronotype and Fatigue Among the Students". *Asian Journal of Medicine and Health* 22 (7):204-13. <https://doi.org/10.9734/ajmah/2024/v22i71061>.

found positively correlated with fatigue ($r = 0.332$, $p < 0.001$) and daytime sleepiness ($r = 0.213$, $p < 0.001$). Chronotype was found negatively correlated with fatigue ($r = -0.124$, $p < 0.001$), daytime sleepiness ($r = -0.105$, $p < 0.05$), and depression ($r = -0.198$, $p < 0.001$). These findings underscore a significant association ($p < 0.001$) between depression and the presence of both excessive daytime sleepiness and fatigue. This revelation demands a nuanced approach to mental health interventions, recognizing the shared underlying mechanisms and advocating for comprehensive strategies that address these intertwined facets of student well-being.

Keywords: Chronotype; depression; fatigue; sleepiness.

ABBREVIATIONS

r-MEQ : Reduced Morningness Eveningness Questionnaire
CT : Chronotype
MT : Morning Type
ET : Evening Type
NT : Neither Type
EDS : Excessive Daytime Sleepiness
FSS : Fatigue Severity Scale
ESS : Epworth Sleepiness Scale
CES-D : Center for Epidemiologic Studies Depression Scale

1. INTRODUCTION

“Fatigue and excessive daytime sleepiness (EDS) frequently have similar meanings in everyday life. These two terms, however, are distinct concepts” [1,2]. “Fatigue and EDS can coexist, but they can also exist independently and with distinct symptoms” [1]. “A population-based study in the Netherlands found that 30.5% of adults have chronic fatigue for more than six months” [3]. “Fatigue is also widespread in the Korean general population, with estimates ranging from 21-31%” [4,5]. The association between EDS and depression has been established in both sleep disorders [6] and the general population [7]. “Several self-report measures have been developed to assess the severity and prevalence” [8]. “The Fatigue Severity Scale is the most commonly used for fatigue measurement” [9,10,11,12]. “Furthermore, fatigue is more common in people who have neurological diseases like multiple sclerosis, Parkinson's disease, and traumatic brain injury” [13]. “It is also linked to psychological issues such as depression, anxiety, and emotional stress [14]. Sleepiness, characterized by an overwhelming urge to sleep during waking hours, and fatigue, encompassing both physical and mental exhaustion, represent significant facets of sleep-related disturbances, in contrast to fatigue” [15].

“Excessive daytime sleepiness (EDS) is the tendency to fall asleep despite deliberate attempts to remain alert. Chronic sleep loss and poor sleep quality are the primary underlying causes of EDS” [16]. “They are present in several sleep disorders, including obstructive sleep apnea and narcolepsy, as well as psychiatric disorders, particularly depression. Approximately 80% of depressive episodes are accompanied by comorbid insomnia, whether related or not to EDS” [16]. “Sleep is regulated by the interaction of sleep homeostasis and circadian rhythm, whereas exertion plays a pivotal role in the occurrence and consequences of fatigue” [17]. “Fatigue and sleepiness are frequently used interchangeably because both conditions produce tiredness symptoms that overlap. However, people who complain of fatigue are not always tired, and vice versa” [1]. “Excessive daytime sleepiness (EDS) is commonly recognised as a symptom of sleep disorders such as obstructive sleep apnea, periodic limb movement disorder, poor sleep hygiene, and chronic sleep deprivation” [18]. “Aside from sleep, circadian rhythms are important in the relationship between depression and EDS” [19]. “Depression is strongly linked to biological rhythms on a variety of levels, including neurobiological systems that support both the pathophysiology of depression, such as the serotonergic system, and the clinical aspects of depression” [19]. “Circadian misalignment can be caused by human circadian system disorders, resulting in sleep disturbances (specifically, insomnia and/or EDS), reduced attention and impaired daytime alertness, a lack of energy, poor performance, a negative mood, and gastrointestinal disorders. Interestingly, these symptoms also occur in depression, which further extends the idea that depression and circadian rhythms are intimately connected” [19].

The effects of Excessive Daytime Sleepiness (EDS) and fatigue on depression constitute a complex interplay that has garnered inadequate attention in scientific inquiry. Despite their

potential to manifest independently or sequentially, the nuanced dynamics between these factors remain largely unexplored. The current study endeavors to fill this gap by delving into the intricate relationship between fatigue, EDS, and depression among students. While their implications for mental health have been recognized, their comparative influence on depression remains elusive. By examining the relative associations of fatigue and EDS with depressive symptoms, this study aims to unravel their differential impacts within the student population.

2. MATERIALS AND METHODS

2.1 Participants

The data for this study was obtained from a comprehensive cross-sectional epidemiological survey that investigated chronotype, depression, excessive daytime sleepiness (EDS), and the Fatigue Severity Scale (FSS). The study included a broad and representative sample of Indian school students aged 11 to 20 years (15.2 ± 1.46), with 305 males (67.8%) and 145 females (32.2%). In addition to evaluating the aforementioned characteristics, the authors conducted an investigation into the participants' demographic profiles and socioeconomic statuses in March 2022. It is important to note that all participants provided explicit written consent before they participated in the study, demonstrating the ethical integrity of the research procedure.

2.2 Evaluation of Fatigue, Sleepiness, Chronotype and Depression

2.2.1 Fatigue Severity Scale (FSS)

The self-reported Fatigue Severity Scale (FSS), one of the most commonly used fatigue assessment indexes, was used to measure fatigue [9]. This questionnaire has nine items that measure the exerting influence of fatigue on specific situations of functioning and the behavioral consequences of fatigue. Each item is rated from 1 (strongly disagree) to 7 (strongly agree). The authors calculated the FSS score as the total summation of all nine items. We defined the presence of fatigue as an FSS score ≥ 36 .

2.2.2 Epworth Sleepiness Scale (ESS)

Daytime sleepiness was assessed using the Epworth Sleepiness Scale (ESS), an eight-item

questionnaire that measures the likelihood of dozing in eight situations [20]. The total ESS score ranges from 0 to 23, and scores ≥ 11 indicate EDS in this study. The combination of EDS and fatigue was classified into four categories as follows; EDS with fatigue (E^+F^+ , $ESS \geq 11$, $FSS \geq 36$), fatigue without EDS (E^-F^+ , $ESS < 11$, $FSS \geq 36$), EDS without fatigue (E^+F^- , $ESS \geq 11$, $FSS < 36$), and no fatigue and EDS (E^-F^- , $ESS < 11$, $FSS < 36$).

2.2.3 reduced-Morningness Eveningness Questionnaire (r-MEQ)

The r-MEQ was developed by Adan et al. [21] and only includes items 1, 7, 10, 18, and 19 of the original MEQ [21]. r-MEQ is used to self-assess the chronotype (CT) of the individuals. The questionnaire has 5 questions and the score range from 4 to 26. Whereby the higher score indicates the Morningness chronotype. The same cut-off scores for determining CT groups were used in Adan et al. [21], (Eveningness chronotype: < 12 ; Neither chronotype: $12-17$; morning: > 17). The validated scoring was used where the first four questions were scored 1–5 and question 5 was scored 0–6.

2.2.4 Center for Epidemiologic Studies Depression Scale (CES-D)

Depressive symptoms were measured using the Center for Epidemiologic Studies Depression Scale (CES-D) [22]. CES-D is among the most frequently used measures for assessing the level of depressive symptoms and is a well-validated self-report measure of depressive symptoms in adults in the community [23]. CES-D is a 20-item and the score range from 0-60. CES-D measure that assesses the frequency of depression during the past week on a 0–3 Likert scale (“rarely or none of the time” to “most or all of the time”). The standard cut-off is ≥ 16 for “possible depression” [22].

2.3 Statistical Analyses

The normality of the dataset was determined using the Kolmogorov-Smirnov test. Categorical variables were shown as frequency and Percentage. Categorical variables were compared using the chi-square test ($p < 0.05$). One-way analysis of variance (ANOVA) was used to compare the means of two or more independent groups. P values < 0.05 were considered statistically significant with a 95% confidence interval (CI). Pearson / Spearman's

correlations were used to assess the strength of relationships between variables. Multiple regression analyses were used to investigate the relationship between daytime sleepiness, fatigue, depression, and chronotype. The dependent variable for linear regression was depression, while the independent variables were fatigue, daytime sleepiness, and chronotype. The data analysis and graph preparation were performed using Statistical Package for the Social Sciences (SPSS) 26.0 (IBM Corporation, NY, USA) and GraphPad Prism Software version 8.0, San Diego, USA.

3. RESULTS

3.1 Characteristics of Participants

A total of 470 participants responded to this questionnaire study. The authors excluded twenty participants due to missing data of fatigue, depression, or the sleepiness scale. After exclusions, the authors analyzed data from 450 participants. The proportion of females was N=145 (19.4%) and males was N=305 (40.8%).

3.2 Comparison of Sleepiness, Depression, and Fatigue with Chronotype

Figs. 1-3 demonstrate the composite scores of CES-D (>16 =D+), ESS (≥11 =E+), and FSS (≥36 =F+) among 450 participants. Results (Fig. 1) show that 52.4% exhibited symptoms of depression, 23.7% were daytime sleepers, and 22.6% were fatigued. Fig. 2 further breaks down these findings by chronotype, Fig. 2(a) demonstrated the depression score among the chronotype where MT (23.02±5.77), NT (24.73±6.08), and ET (28.62±8.02) which revealed that people with an evening type (ET) demonstrated higher levels of depression and ET

was also found highly significant different with MT (p=0.000) and NT (p=0.013). Fig. 2(b) shows the sleepiness score among the chronotype MT (13.28±2.22), NT (13.59±2.20), and ET (13.00±1.22) whereas Fig. 2(c) shows the fatigue score among the chronotype MT (40.50±3.56), NT (41.25±3.75), and ET (42.62±3.66), both were not found significantly different from each other.

Fig. 3 Delves into gender differences among the chronotypes. Fig. 3(a) shows the depression score with chronotype where MT male (22.23±4.87) female (25.22±7.43), NT male (23.77±5.85) female (25.96±6.19), ET male (28.29±7.14) female (29.00±9.25), showing that females exhibited higher rates of depression across all chronotypes. ET males were found to be highly significantly different from MT(p=0.001) and NT (p=0.017) males. Fig. 3(b) shows the sleepiness score among the chronotype with gender differences where MT male (13.35±2.36) female (13.08±1.88), NT male (13.16±2.39) female (14.00±1.96), ET male (13.57±1.13) female (12.33±1.03). Fig. 3(c) shows the fatigue score MT male (40.62±3.66) female (41.00±3.53), NT male (40.52±3.28) female (41.89±4.07), ET male (42.71±3.35) female (42.50±4.32).

Table 1 lists the study participants' demographic, r-MEQ, and social behavior characteristics according to the presence of depression. Participants with depression showed a significant preponderance for being female (N = 89, 37.7%) (p < 0.05). Age range from 16-20 was significantly related to depression (N=108, 45.8%) (p < 0.05). The subjects who preferred to be in the group were significantly related to depression (N=153, 64.8%) (p <0.001). Depression was significantly higher in ESS (N= 165, 69.9%) (p < 0.001) and fatigue (N= 167, 70.8%) (p < 0.001), respectively.




Fig. 1. Represent the Depression, Sleepiness and Fatigue scores among the participants
 D+, presence of depression; E+, presence of sleepiness; F+, presence of fatigue

Table 1. Characteristics of participants according to the presence of depression

Variables		Total N (%)	Depression (-) N (%)	Depression (+) N (%)	X ² -value	df	p-value
Gender	Male	305 (67.8)	158 (73.8)	147 (62.3)	6.848	1	0.009
	Female	145 (32.2)	56 (26.2)	89 (37.7)			
Age	11-15	264 (58.7)	136 (63.6)	128 (54.2)	4.015	1	0.045
	16-20	186 (41.3)	78 (36.4)	108 (45.8)			
Education level	Class 8-10	240 (53.3)	120 (56.1)	120 (50.8)	1.232	1	0.267
	Class 11-12	210 (46.7)	94 (43.9)	116 (49.2)			
Belongs to	Ruler	128 (28.4)	66 (30.8)	62 (40.3)	1.152	1	0.283
	Urban	322 (71.6)	148 (69.2)	174 (73.7)			
Family status	Joint	186 (41.3)	91 (42.5)	95 (40.3)	0.238	1	0.625
	Nuclear	264 (58.7)	123 (57.5)	141 (59.7)			
Prefer to	Live alone	129 (28.7)	46 (21.5)	83 (35.2)	10.262	1	0.001
	In group	321 (71.3)	168 (78.5)	153 (64.8)			
Chronotype	MT	179 (39.8)	92 (43.0)	87 (36.9)	5.774	2	0.056
	NT	234 (52.0)	111 (51.9)	123 (52.1)			
	ET	37 (08.2)	11 (05.1)	26 (11.0)			
Daytime Sleepiness	Normal	343 (76.2)	178 (83.2)	71 (30.1)	10.892	1	0.001
	Sleepiness	107 (23.8)	36 (16.8)	165 (69.9)			
Fatigue	Normal	348 (77.3)	181 (84.6)	69 (29.2)	12.223	1	0.000
	Fatigue	102 (22.7)	33 (15.4)	167 (70.8)			
ESS<11, FSS<36	E-F ⁻	286 (63.6)	155 (34.4)	131 (29.1)	18.608	3	0.000
ESS≥11, FSS<36	E+F ⁻	62 (13.8)	26 (05.8)	36 (08.0)			
ESS≥11, FSS≥36	E+F ⁺	45 (10.0)	10 (02.2)	35 (07.8)			
ESS < 11, FSS ≥ 36	E-F ⁺	57 (12.7)	23 (05.1)	34 (07.6)			




Fig. 2. Comparison of Depression, Sleepiness and Fatigue with Chronotype among the participants. Fig. 2a shows the depression, 2b shows the Sleepiness and 2c shows the Fatigue score among the chronotype

Fig. 3a, shows the depression, 3b shows sleepiness and 3c shows the fatigue score in between the gender and chronotype

Table 2. Characteristics of participants according to excessive daytime sleepiness according to two categories

Variables		Total N (%)	E ⁺ F ⁻ N (%)	E ⁻ F ⁺ N (%)	X ² -value	df	p-value
Gender	Male	72 (60.5)	41 (66.1)	31 (54.4)	1.714	1	0.191
	Female	47 (39.5)	21 (33.9)	26 (45.6)			
Age	11-15	59 (49.6)	34 (54.8)	25 (43.9)	1.432	1	0.231
	16-20	60 (50.4)	28 (45.2)	32 (56.1)			
Education level	Class 8-10	50 (42.0)	25 (40.3)	25 (43.9)	0.153	1	0.696
	Class 11-12	69 (58.0)	37 (59.7)	32 (56.1)			
Belongs to	Rural	30 (25.2)	15 (24.2)	15 (26.3)	0.071	1	0.790
	Urban	89 (74.8)	47 (75.8)	42 (73.7)			
Family status	Joint	55 (46.2)	30(48.4)	25 (43.9)	0.245	1	0.621
	Nuclear	64 (53.8)	32 (51.6)	32 (56.1)			
Prefer to	Live alone	33 (27.7)	22 (35.5)	11 (19.3)	3.882	1	0.049
	In group	86 (72.3)	40 (64.5)	46 (80.7)			
Chronotype	MT	41 (34.5)	24 (38.7)	17 (29.8)	1.047	2	0.592
	NT	66 (55.5)	32 (51.6)	34 (59.6)			
	ET	12 (10.1)	06 (09.7)	06 (10.5)			
Depression	Normal	49 (41.2)	26 (41.9)	23 (40.4)	0.031	1	0.861
	Depressed	70 (58.8)	36 (58.1)	34 (59.6)			

Table 3. The correlation between daytime sleepiness, fatigue, depression, and chronotype scale score

Variables	Mean ±SD	1	2	3	4
Fatigue	25.55 ±13.11	1			
Day time Sleepiness	8.02 ±4.03	0.451**	1		
Depression	17.84 ±8.92	0.332**	0.213**	1	
Chronotype	16.53 ±4.46	-0.124*	-0.105*	-0.198**	1

* Correlation is significant at the 0.05 level (2- tailed)

** Correlation is significant at the 0.01 level (2- tailed)

Table 4. Represented the multiple linear regression analysis where the dependent variable was Depression (CES-D) and the independent variables were chronotype, fatigue, and Daytime sleepiness (P<0.001)

Model Depression (CES-D)	Standardized coefficient (Beta)	t	p	R	R ²	F	Model p-value
Chronotype	-0.156	-3.522	0.000				
Fatigue	0.281	5.695	0.000	0.373	0.139	24.091	0.000
Day time Sleepiness	0.070	1.425	0.155				

In four categories according to the presence combination of ESS and fatigue, E⁻F⁻ (N=131, 29.1%) was low in participants with depression, and E⁺F⁻ (N=36, 8.0%) was related to participants with depression. E⁻F⁺ (N=34, 07.6%) and E⁺F⁺ (N=35, 07.8%) had higher participants with depression. These findings underscore a significant association (p<0.001) between depression and the presence of both excessive daytime sleepiness and fatigue.

3.3 Comparison of between E⁺F⁻ and E⁻F⁺

Table 2 depicts the participants' characteristics of the two groups, E⁺F⁻ (N = 62) and E⁻F⁺ (N = 57). Compared to E⁺F⁻, E⁻F⁺ was associated with subjects who preferred to be in the group (n= 46, 80.7%) (p <0.05). Gender, age, Education level, chronotype, depression, and other variables were not significantly associated with both groups.

Depression was found positively correlated (Table 3) with fatigue (r =0.332, p < 0.001) and daytime sleepiness (r=0.213, p<0.001). Chronotype was found negatively correlated with fatigue (r=-0.124, p<0.001), daytime sleepiness (r=-0.105, p<0.05), and depression (r=-0.198, p<0.001)

The multiple linear regression analysis (Table 4) demonstrated that fatigue (β= 0.281, t=5.695, p<0.001) and daytime sleepiness (β= 0.070, t= 1.425, p=0.155) were related to depression. However, there was no direct impact of

chronotype on depression (β= -0.156, t= -3.522, p<0.001).

4. DISCUSSION

The primary aim of this study was to estimate the degree of association between fatigue and excessive daytime sleepiness (EDS) with depression in the student population. Depression was significantly associated with E⁺F⁺, E⁻F⁺, and E⁺F⁻, compared with E⁻F⁻ as the reference category. E⁺F⁺ was the most closely related to depression. This potent association between depression and E⁺F⁺ is consistent with the result of a previous study [24]. EDS and fatigue share aspects in relation to depression. First, EDS and unexplained fatigue are some of the symptoms that precede depression [25]. Second, EDS and fatigue are a kind of somatic symptom caused by depression and tend to remain as a residual symptom even with pharmacological treatment through selective serotonin reuptake inhibitors [26]. Third, EDS and fatigue tend to cluster together rather than presenting as isolated symptoms. In women of the general population, fatigue and EDS were identified as the most critical risk factors for psychiatric diseases including depression [27]. As in our study, the distribution of EDS and fatigue tended to cluster together rather than to exist separately.

The outcomes, as depicted in Table 1, revealed significant distinctions (p=0.000) among the four delineated categories of sleepiness and fatigue, all of which were closely linked with depression

[28]. Notably, the E⁺F⁺ group exhibited a higher incidence of depression (D⁺) in comparison to the E⁻F⁺ and E⁺F⁻ cohorts, while the E⁻F⁻ subset displayed a reduced prevalence of D⁺ [28]. Furthermore, gender, age, preference, sleepiness, and fatigue emerged as factors significantly associated with depression. The findings suggested a notable elevation in depression among individuals experiencing symptoms of sleepiness and fatigue [29]. This observation raises the possibility of a bidirectional relationship between depression and these symptoms, wherein depression may exacerbate sleepiness and fatigue, or vice versa. It is conceivable that the presence of depression disrupts sleep patterns, thereby contributing to daytime sleepiness and fatigue during waking hours [30].

There is ample evidence for an association between daytime sleepiness and moderate to severe depression [31]. Given that insomnia, particularly when associated with impaired daytime function, plays a key role in the onset of depression [32], it is of significant clinical importance to unravel the link between EDS and depression. Furthermore, knowledge of the interrelation between EDS and mood disorders may have ramifications for the treatment of depression [31].

These findings underscore the substantial role played by depression in the manifestation of sleepiness and fatigue. Table 2 offered additional insights, indicating that individuals who preferred group settings (E⁺F⁻ and E⁻F⁺) exhibited higher levels of depression compared to those who lived alone. This trend may be attributed to a desire for social interaction to alleviate feelings of loneliness. Similar trends were observed in Table 3, which highlighted a positive correlation between sleepiness, fatigue, and depression. Table 4 further demonstrated that depression exhibited a stronger association with fatigue than with sleepiness, suggesting a closer relationship between depression and fatigue. Collectively, these results underscore the complex interplay between depression, sleepiness, and fatigue, emphasizing the need for comprehensive approaches to addressing these interconnected phenomena. However, several limitations warrant consideration. The study's cross-sectional design precludes causal inferences, emphasizing the importance of longitudinal research to understand the temporal relationship between these variables. Furthermore, using self-reported measures may introduce response

bias, and the study's sample may not be representative of the broader population.

5. CONCLUSION

This study sought to investigate the link between fatigue, excessive daytime sleepiness (EDS), and depression in students. The results show a significant relationship between depression and various combinations of sleepiness and fatigue, with the E⁺F⁺ group having the strongest association with depression. This aligns with prior research, suggesting a robust link between depression and the co-occurrence of sleepiness and fatigue. The study highlights several key insights. Firstly, EDS and fatigue may precede depression and persist as residual symptoms despite treatment, indicating a complex relationship wherein these symptoms may serve as both precursors and consequences of depression. Secondly, the clustering of EDS and fatigue suggests a shared underlying mechanism, further supporting their association with depression. This revelation demands a nuanced approach to mental health interventions, recognizing the shared underlying mechanisms and advocating for comprehensive strategies that address these intertwined facets of student well-being.

FUNDING

This work is supported by the grant funded by Indian Council of Medical Research (ICMR), New Delhi, India under grant number - SRF/05/2022/SBHSR of author SJ.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during the writing or editing of manuscripts.

CONSENT

As per international standards or university standards, Participants' written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

As per international standards or university standards written ethical approval has been collected and preserved by the author(s).

ACKNOWLEDGEMENT

The authors would like to express their gratitude towards all the participants who volunteered for this study and their contribution to data collection.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Matsangas P, Shattuck NL. Discriminating Between Fatigue and Sleepiness in the Naval Operational Environment. *Behavioral Sleep Medicine*. 2018;16(5):427-436.
DOI:10.1080/15402002.2016.1228645
2. Pigeon WR, Sateia MJ, Ferguson RJ. Distinguishing between excessive daytime sleepiness and fatigue. *J Psychosom Res*. 2003;54(1):61-69.
DOI: 10.1016/S0022-3999(02)00542-1
3. Van'T Leven M, Zielhuis GA, Van Der Meer JW, Verbeek AL, Bleijenberg G. Fatigue and chronic fatigue syndrome-like complaints in the general population. *Eur J Public Health*. 2010;20(3):251-257.
DOI: 10.1093/eurpub/ckp113
4. Kim CH, Shin HC, Won CW. Prevalence of Chronic Fatigue and Chronic Fatigue Syndrome in Korea: Community-Based Primary Care Study. *J Korean Med Sci*. 2005;20(4):529-534.
DOI:10.3346/jkms.2005.20.4.529
5. Sunwoo JS, Kim D, Chu MK, Yun CH, Yang KI. Fatigue is associated with depression independent of excessive daytime sleepiness in the general population. *Sleep and Breathing*. 2022;26(2):933-940.
DOI: 10.1007/s11325-021-02448-3
6. Sharafkhaneh A, Giray N, Richardson P, Young T, Hirshkowitz M. Association of Psychiatric Disorders and Sleep Apnea in a Large Cohort. *Sleep*. 2005;28(11):1405-1411.
DOI: 10.1093/sleep/28.11.1405
7. Bixler EO, Vgontzas AN, Lin HM, Calhoun SL, Vela-Bueno A, Kales A. Excessive daytime sleepiness in a general population sample: The role of sleep apnea, age, obesity, diabetes, and depression. *Journal of Clinical Endocrinology and Metabolism*. 2005;90(8):4510-4515.
DOI: 10.1210/jc.2005-0035
8. Dittner AJ, Wessely SC, Brown RG. The assessment of fatigue: A practical guide for clinicians and researchers. *J Psychosom Res*. 2004;56(2):157-170.
DOI: 10.1016/S0022-3999(03)00371-4
9. Krupp LB. The fatigue severity scale. *Arch Neurol*. 1989;46(10):1121-1123.
DOI:10.1001/archneur.1989.00520460115022
10. Chianeh, Yousef Rezaei, Ali Mansouri Mehrabani, Alireza Akhavan Kharazi, Mahsa Ghaffari, and Faride Asgari. Role of nutritional factors, mitochondrial dysfunction, altered body composition and various biological factors in chronic fatigue syndrome. *Journal of Advances in Medical and Pharmaceutical Sciences*. 2024;26(6): 23-36.
Available:https://doi.org/10.9734/jamps/2024/v26i6692.
11. Alapin I, Fichten CS, Libman E, Creti L, Bailes S, Wright J. How is good and poor sleep in older adults and college students related to daytime sleepiness, fatigue, and ability to concentrate? *Journal of Psychosomatic Research*. 2000 Nov 1;49(5):381-90.
12. Bener A, Yildirim E, Torun P, Çatan F, Bolat E, Aliç S, Akyel S, Griffiths MD. Internet addiction, fatigue, and sleep problems among adolescent students: A large-scale study. *International Journal of Mental Health and Addiction*. 2019 Aug 15;17:959-69.
13. Kluger BM, Krupp LB, Enoka RM. Fatigue and fatigability in neurologic illnesses: Proposal for a unified taxonomy. *Neurology*. 2013;80(4):409-416.
DOI:10.1212/WNL.0b013e31827f07be
14. Chen MK. The epidemiology of self-perceived fatigue among adults. *Prev Med (Baltim)*. 1986;15(1):74-81.
DOI: 10.1016/0091-7435(86)90037-X
15. Shahid A, Shen J, Shapiro CM. Measurements of sleepiness and fatigue. *J Psychosom Res*. 2010;69(1):81-89.
DOI: 10.1016/j.jpsychores.2010.04.001
16. Fava M. Daytime sleepiness and insomnia as correlates of depression. *J Clin Psychiatry*. 2004;65(16):27-32.
17. Phillips RO. A review of definitions of fatigue - And a step towards a whole definition. *Transp Res Part F Traffic Psychol Behav*. 2015;29:48-56.
DOI: 10.1016/j.trf.2015.01.003

18. Slater G, Steier J. Excessive daytime sleepiness in sleep disorders. *J Thorac Dis.* 2012;4(6):608-616. DOI:10.3978/j.issn.2072-1439.2012.10.07
19. Wirz-Justice A. Chronobiology and mood disorders. *Dialogues Clin Neurosci.* 2003; 5(4):315-325. DOI:10.31887/DCNS.2003.5.4/awirzjustice
20. Johns MW. A New Method for Measuring Daytime Sleepiness: The Epworth Sleepiness Scale. 1991;14. Available: <https://academic.oup.com/sleep/article-abstract/14/6/540/2742871>
21. Adan A, Almirall H. Horne & Östberg morningness-eveningness questionnaire: A reduced scale. *Pers Individ Dif.* 1991; 12(3):241-253. DOI: 10.1016/0191-8869(91)90110-W
22. Radloff LS. The CES-D Scale: A Self-Report Depression Scale for Research in the General Population. *Appl Psychol Meas.* 1977;1(3):385-401. DOI:10.1177/014662167700100306
23. Smarr KL, Keefer AL. Measures of depression and depressive symptoms: Beck Depression Inventory-II (BDI-II), Center for Epidemiologic Studies Depression Scale (CES-D), Geriatric Depression Scale (GDS), Hospital Anxiety and Depression Scale (HADS), and Patient Health Questionnaire-9 (PHQ-9). *Arthritis Care Res (Hoboken).* 2011;63(SUPPL. 11). DOI: 10.1002/acr.20556
24. Bosse-Henck A, Koch R, Wirtz H, Hinz A. Fatigue and Excessive Daytime Sleepiness in Sarcoidosis: Prevalence, predictors, and relationships between the two symptoms. *Respiration.* 2017;94(2): 186-197. doi:10.1159/000477352
25. Lauer CJ. In Quest of Identifying Vulnerability Markers for Psychiatric Disorders by All-Night Polysomnography. *Arch Gen Psychiatry.* 1995;52(2):145-153. DOI:10.1001/archpsyc.1995.03950140063009
26. Papakostas GI, Nutt DJ, Hallett LA, Tucker VL, Krishen A, Fava M. Resolution of Sleepiness and Fatigue in Major Depressive Disorder: A Comparison of Bupropion and the Selective Serotonin Reuptake Inhibitors. *Biol Psychiatry.* 2006; 60(12):1350-1355. DOI:10.1016/j.biopsych.2006.06.015
27. Theorell-Haglöw J, Lindberg E, Janson C. What are the Important Risk Factors for Daytime Sleepiness and Fatigue in Women? *Sleep.* 2006;29(6):751-757. DOI:10.1093/sleep/29.6.751
28. Carney CE, Moss TG, Lachowski AM, Atwood ME. Understanding mental and physical fatigue complaints in those with depression and insomnia. *Behavioral Sleep Medicine.* 2014;12(4):272-289. DOI: 10.1080/15402002.2013.801345
29. Corfield EC, Martin NG, Nyholt DR. Co-occurrence and symptomatology of fatigue and depression. *Compr Psychiatry.* 2016; 71:1-10. DOI: 10.1016/j.comppsy.2016.08.004
30. Chellappa SL, Schröder C, Cajochen C. Chronobiology, excessive daytime sleepiness and depression: Is there a link? *Sleep Med.* 2009;10(5):505-514. DOI: 10.1016/j.sleep.2008.05.010
31. Breslau N, Roth T, Rosenthal L, Andreski P. Daytime sleepiness: An epidemiological study of young adults. *Am J Public Health.* 1997;87(10):1649-1653. DOI:10.2105/AJPH.87.10.1649
32. Ohayon MM, Roth T. Place of chronic insomnia in the course of depressive and anxiety disorders. *J Psychiatr Res.* 2003;37(1):9-15. DOI:10.1016/S0022-3956(02)00052-3

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/119138>