



Potential Health Benefits of Conjugated Linoleic Acid: An Important Functional Dairy Ingredient

**Chaudhari Kalpeshkumar Virsangbhai¹, Ankit Goyal^{1*}, Beenu Tanwar²
and Manvesh Kumar Sihag¹**

¹*Department of Dairy Chemistry, Mansinhbhai Institute of Dairy and Food Technology, Mehsana-384002, Gujarat, India.*

²*Department of Dairy Technology, Mansinhbhai Institute of Dairy and Food Technology, Mehsana-384002, Gujarat, India.*

Authors' contributions

This work was carried out in collaboration among all authors. Authors CKV and AG wrote the first draft of the article, while, author BT prepared the figure 2 and table 2 and edited the manuscript. Author MKS managed the literature search and prepared the table 3. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/EJNFS/2019/v11i430162

Editor(s):

(1) Dr. Johnson Akinwumi Adejuyitan, Senior Lecturer, Department of Food Science and Engineering, Ladake Akintola University of Technology (LAUTECH), Nigeria.

Reviewers:

(1) Asaad Ahmed Ghanem, Mansoura University, Egypt.

(2) Jihan Seid Hussein, National Research Centre, Egypt.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/53785>

Review Article

Received 25 October 2019
Accepted 28 December 2019
Published 11 January 2020

ABSTRACT

Conjugated linoleic acid (CLA) refers to a class of positional and geometrical isomers of linoleic acid (cis-9, cis-12 octadecadienoic acid) having conjugate double bond system. CLA are synthesized in rumen of the ruminants by biohydrogenation of dietary fatty acids; and thus, can be obtained from dairy products as well as from the meat of sheep, lamb and other ruminants. Among the several isomers, c9, t11-CLA isomer is the most biologically active form and accounts approximately 80% of total isomers. A number of clinical and epidemiological studies have demonstrated the role of CLA as anti-atherogenic, anti-inflammatory, anti-oxidative, anti-carcinogenic, etc. Several researchers have suggested the positive association of CLA in weight management, hypercholesterolemia, immunomodulatory functions, and improved bone metabolism.

Keywords: *Conjugated linoleic acid; biohydrogenation; anti-carcinogenic activity; anti-atherogenic; anti-obesity.*

*Corresponding author: Email: ankit.ndri@gmail.com;

1. INTRODUCTION

Conjugated linoleic acid (CLA) refers to a mixture of positional and geometric isomers of linoleic acid (C18:2). CLA are naturally synthesized from linoleic acid in the rumen and intestine of pastured ruminants like cattle, sheep and goat. Conjugated linoleic acids have conjugated dual bond and consist a class of geometric and positional isomers of essential linoleic fatty acid (cis-9, cis-12) by biohydrogenation. CLA isomers may be present in various double bond positions (7–9, 8–10, 9–11, 10–12, 11–13) or geometric orientation (trans-cis, cis-trans, trans-trans & cis-cis). However, cis-9, trans-11 is the most predominant and biologically active form. The c9, t11 isomer has been reported to have anti-carcinogenic effects as well as regulating body mass. Other isomer of CLA such as t10, c12 has also been reported to confer several health benefits such as preventing artery thickening, protecting the heart disease, preventing from cancer, and modulating immune system. Technically, conjugated linoleic acid is not an essential fatty acid. Unlike other fatty acids, CLA cannot be synthesized by the human body, and thus, needs to be taken from diet such as meat and dairy products. It is important to note that CLA has two double bonds in conjugated form, in which one is present in cis and other is present in trans configuration. As per the definition given by CODEX, polyunsaturated fatty acids having isolated (non-conjugated) double bonds in trans configuration only are known as trans fatty acids [1]. Therefore, the isomers of CLA are exempted from the trans-fat labelling [1]. Moreover, the equivalent mixture of c9, t11 and t10, c12 CLA isomers (1:1) have been granted GRAS

(Generally recognized as safe) status by FDA in 2008.

1.1 Sources

The most common sources of natural CLA in the human diet are milk & milk products and meat of ruminant animals. The content of CLA in various products are given in below table (Table 1).

1.2 Global Market of CLA

CLA is a popular supplement for weight management worldwide. CLA is used for medical benefits like cancer fighting properties, muscle strengthening, improved metabolic functions. Conjugated Linoleic Acid (CLA) market is expected to grow 5.8% Compound Annual Growth Rate (CAGR) in terms of revenues. According to a recent report, the size of conjugated linoleic acid market is expected to reach US\$ 50 million up to 2024, from US\$ 36 million in 2019 [3]. China is the biggest producer of CLA supplements capturing 47.93% market share followed by North America (25.63%).

1.3 Biosynthesis of CLA

In ruminants, CLA is synthesized by two methods: 1) Ruminal Biohydrogenation (from linoleic acid), and 2) Endogenous synthesis (in tissues from *trans* Vaccenic acid). First the linoleic acid (C18:2, C9, C11) is converted into conjugated linoleic acid (C9, t11) in presence of linoleate isomerase. Thus, CLA formed goes to mammary glands and gets incorporated into the milk fat. Some of the portion of CLA is





Common Name	ω -Position (from -CH ₃ end)	Structure	Systemic name (from -COOH end)
Oleic acid	ω -9		(<i>cis</i> -9)-Octadecaenoic acid
Linoleic acid	ω -6		(<i>cis</i> -9, <i>cis</i> -12)-Octadecadienoic acid
9-CLA (Rumenic acid)	ω -7		(<i>cis</i> -9, <i>trans</i> -11)-Octadecadienoic acid
10-CLA	ω -6		(<i>trans</i> -10, <i>cis</i> -12)-Octadecadienoic acid

Fig. 1. Structure of CLA and other related fatty acids (Source: Benjamin et al. 2015 [2])

Table 1. Conjugated linoleic acid content (mg CLA/g fat) in various food products

Dairy products	Mg of CLA/g of fat	Meat products	Mg of CLA/g of fat
Milk	5.5	Veal	2.7
Condensed milk	7.0	Ground beef	4.3
Butter fat	6.1	Lamb	5.8
Cultured milk	5.4	Chicken	0.9
Sour cream	4.6	Pork	0.6
Butter	4.7	Ground turkey	2.6
Ice cream	3.6	Egg yolk	0.6
Yogurt (low fat)	4.4	Salmon	0.3
Plain yogurt	4.8		
Custard yogurt	4.8		
Medium cheddar	4.1		

Source: Chin et al., [4]

**Fig. 2. Biosynthesis of conjugated linoleic acid (Adapted from Bauman et al. 2000 [5])**

hydrogenated into trans-vaccenic acid (TVA) (t11) (Fig. 2). The trans-vaccenic acid is further hydrogenated into endogenous stearic acid. On the other hand, α -Linolenic acid (ALA) (C18:3) (c9, c11, c15) is converted into Octadecatrienoic acid (C18:3) (c9, t11, c15) in the presence of linoleate isomerase. Hereafter, further conversion is shown in Fig. 2.

2. ROLE OF RUMEN BACTERIA IN CLA SYNTHESIS

For CLA synthesis, ruminants depend on microbial fermentation of feed and additional forage in rumen. Ruminant diet includes some polyunsaturated fatty acids (PUFAs) which are toxic to many of rumen microorganisms [6]. Nevertheless, in order to protect them from toxic effects, rumen microorganisms have pathways for hydrolysis and biohydrogenation of dietary lipid. Rumen bacteria play an important role in biohydrogenation. In biohydrogenation of lipids in

rumen, >250 bacterial strains are involved. The most common are:

- *Butyrovibrio*
- *Micrococcus*
- *Lactobacillus*
- *Ruminococcus*
- *Enterococcus*
- *Propionibacterium*

3. HEALTH BENEFITS OF CLA

Several recent clinical and epidemiological studies have reported the important role of CLA as anticarcinogenic, antiatherogenic, immunomodulating, anti-diabetic and lean body mass enhancing, etc., which are discussed below:

Anticarcinogenic activity: Though no concrete evidences for the anticarcinogen activity of CLA have been documented yet, possible mechanisms could be attributed to antioxidant

mechanisms, pro-oxidant cytotoxicity, suppression of nucleotide biosynthesis, decrease in proliferative activity and inhibition of carcinogen activation [7,8,9,10,1,11,12]. CLA is reported to show prooxidant activity in which they inhibit the oxidation causing substances [13]. A few studies have demonstrated that CLA decreases the proliferative activity and suppress the multiplication of free radical formation [14,15]. In another study, it is suggested that CLA suppresses the nucleotide synthesis and thus, inhibit the DNA replication in cells which prevent the formation of tumor in the body. Recent clinical and epidemiological studies on the anti-carcinogenic properties of CLA are shown in Table 2.

Antioxidant and anti-inflammatory properties:

CLA is a more potent antioxidant than other antioxidants. It is more effective than butylated hydroxytoluene (BHT) & alpha-tocopherol. Furthermore, it is reported to be more potent than vitamin E and butylated hydroxy anisole (BHA) in suppressing the formation of thio-barbituric acid reactive substances (TBARS). CLA is one of the biomarkers which is also used to assess oxidation status in biological systems. Environmental mutagens or carcinogens act as an inhibitor and they block the carcinogenesis and show their antioxidant properties [16]. Superoxide dismutase and catalase are the antioxidative enzymes involved into the removal of toxic free radicals and thus, lead to amelioration of oxidative damage in cells. Glutathione-S transferase is involved in biotransformation of carcinogens, detoxification of xenobiotics, peroxides and free radicals by conjugating toxic component with GSH (Glutathione reduced) and due to this they ultimately protecting organ and cells against inducing toxicity [17]. For this reason, increasing this enzyme by synthetic or natural way, which results in the inhibition of hepatic cancer-causing substances; and CLA behaves like chemo-preventive agent, inhibits the oxidative stress & reduce the levels of detoxifying enzymes [18]. It is reported that CLA at the level of 0.25% or more in diet inhibits the mammary carcinogenesis and decreases the formation of TBARS into mammary tissue [19]. Kathirvelan [20] observed that CLA fortified *ghee* (clarified butter) increased the antioxidative activity and reduced the toxic causing enzymes in mammary gland and liver than that of soybean oil. Several clinical and epidemiological studies on the anti-inflammatory and immunomodulatory properties of CLA are presented in Table 3.

Cholesterol-lowering and anti-obesity effects:

Several recent animal studies have demonstrated a positive role of CLA in prevention of cardiovascular diseases, hypercholesterolemia and obesity [21,22,23], which are presented in Table 4. Lee et al. [24] reported that CLA was effective against different stage of cancer like progression and initiation and found anti-atherogenic effect of CLA (at 0.5 g/d) in rabbits in a study of 22 days. On the other hand, Nicolosi et al., [25] reported that the hamsters fed together with CLA had significantly decrease the amount of total plasma cholesterol when compared to control. Kathirvelan [20] observed that the high and low CLA intakes reduced the total cholesterol, and triglycerides, and increased HDL cholesterol when compared with control (soya oil fed rats). Kathirvelan [20] demonstrated that the antiatherogenic effects of CLA were shown by decrease in triglycerides, total cholesterol, atherogenic index & LDL cholesterol and a rise in HDL-cholesterol into blood plasma.

CLA and Bone metabolism: Watkins et al. [26] have reported a high level of bone formation in butterfly-fed chickens, which proposed to be likely due to higher CLA consumption. Dietary CLA contributed to variations in CLA abundance in bone marrow, specific tissue and organs, and periosteum having the highest amount of CLA and the low amount in the heart. CLA has been reported to increase cis-9, trans-11 in tissue lipid and also involved in bone biomarkers and development.

Methods to increase CLA concentration in dairy products:

There are two methods of CLA enrichment: 1) Dietary modification of animal feed, and, 2) Direct enrichment of milk and dairy products. In dietary modification of animal feed, feed can be supplemented with rapeseed, cottonseed, soybean, corn, peanut, sunflower, canola, safflower and linseed or their oils. Oils can be added into the diet in the form of protected oils, free oils, processed oilseeds or whole oilseeds (extruded, crushed, roasted or ground). These seeds/oils are high in linoleic or linolenic acid, which are precursors of CLA. Oil to be added in diet can be protected in the form of fatty acyl amides, calcium salts, a lipid encapsulation or formaldehyde-protein protection matrix.

Food applications: CLA-rich oils can be used as functional ingredients in a variety of foods such as milk, *yoghurt/dahi*, cheese and *paneer*,

Table 2. Major findings of clinical and animal studies based on anti-carcinogenic activity of CLA

Subject	Duration of feeding	CLA concentration in diet	Results	Reference
Mice	NR	1 g CLA/100 g diet	Dietary CLA supplementation reduced disease activity, decreased colitis, and prevented adenocarcinoma formation in cancer-induced mice. Additionally, a significant decrease was also observed in the percentages of macrophages in lymph nodes and the gene expression for colonic tumor necrosis factor- α .	Evans et al., [8]
Rectal cancer patients (n=34)	6 weeks	3gCLA/d	On comparison with placebo-group, CLA-group showed significant reductions in the pro-inflammatory markers [TNF- α (P = 0.04), hsCRP (P = 0.03)], and biomarkers of angiogenesis (MMP-9). However, no significant change was observed in the level of IL-6 than that of control.	Mohammadzadeh et al., [10]
Rats	NA	NA	Polymeric nanoparticles containing linoleic acid conjugate (NPs) were developed. Results revealed that CLA-NPs were effective in growth inhibition of human colon cancer cells. Another important observation was that CLA-NPs could avoid the phagocytosis by macrophages and promote the uptake by cancer cell, suggesting the NPs could be promising candidates to treat colorectal cancer.	Cheng et al., [12]
Human colon cancer cell lines (Caco-2, HT-29 and DLD-1) (<i>In vitro</i>)	72 h	200 μ M CLA	Cell proliferation of human colon cancer cell lines was inhibited by all the isomers of conjugated linoleic acid. The maximum inhibition was shown by t9,t11-CLA, followed by t10,c12-CLA, c9,c11-CLA and c9,t11-CLA, respectively. The extent of apoptosis was also shown by all the isomers, maximum by t9,t11-CLA. The other isomers exhibited much lower apoptosis effect on Caco-2 cells.	Beppu et al., [29]
Mice and colon cancer cell lines (<i>in vitro</i>)	4 weeks for <i>in vivo</i>	0.1% c9, t11 or t 10,c 12 CLA	<i>In vitro</i> : c9,t11 CLA significantly inhibited cancerous cell growth (P<0.05), whereas t10,c12 CLA isomer had no effect on cell migration. <i>In vivo</i> : c9,t11 CLA isomer significantly (P<0.05) inhibited the activity of matrix metalloproteinases (MMPs), which are actively involved and generated in angiogenesis and metastasis. However, t10,c12 CLA isomer did not inhibit the activity of MMPs. Furthermore, both the isomers were equally effective in inhibiting the colon cancer cell metastasis <i>in vivo</i> .	Soel et al., [7]
Colon cancer	72 h	25 and 50 μ M CLA	Data suggested that t10,c12 CLA exhibited cytotoxic effect through	Pierre et al., [11]

Subject	Duration of feeding	CLA concentration in diet	Results	Reference
cells (<i>in vitro</i>)			reactive oxygen species (ROS) generation and a subsequent endoplasmic reticulum stress-dependent apoptosis in colon cancer cells. Overall, t10,c12 CLA showed an inhibition effect on the proliferation of colon cancer cells. It was also included that t10,c12 CLA was more efficient pro-apoptotic fatty acid than c9,t11 CLA.	
MCF-7 breast cancer cells (<i>in vitro</i>)	24 h	50 μ M CLA	Data exhibited that t9, t11 was the most efficient CLA isomer which decreased the proliferation and migration, and induced the apoptosis of MCF-7 breast cancer cells after 24 h of treatment. Another important observation was that the t9, t11 CLA treatment reduced the intracellular and membrane-associated cholesterol levels.	El Roz et al., [30]
Human colorectal adenocarcinoma cells (<i>in vitro</i>)	24 h	25, 50 and 100 μ M CLA	Results revealed that t10, c12-CLA induced the expression of ATF3 mRNA and luciferase activity (of ATF3 promoter), which are the main factors associated with apoptosis in colorectal cancer. Overall, it was suggested that t10, c12 CLA isomer treatment can induce the apoptosis of colon cancer cells, but not the most common CLA isomer i.e. c9, t11 as reported by Lee et al., (2006) in a similar study.	Kim et al., [1]

Table 3. Major findings of clinical and animal studies based on anti-inflammatory and immunomodulatory activities of CLA

Subject	Duration	CLA concentration in diet	Major findings	Reference
Mouse and <i>in vitro</i>	NR	-	Decreased pro-inflammatory cytokines and inhibited leukocyte recruitment <i>in vivo</i> . While, <i>in vitro</i> , attenuated NF- κ B-dependent gene expression, decrease pro-inflammatory cytokine production and up-regulated Nrf2-regulated proteins	Villacorta et al., [31]
<i>In vitro</i> (bovine mammary epithelial cells: BME-UV1)	-	-	Bovine mammary epithelial cells treated with CLA showed significantly lower levels of ROS when compared with other cells treated with ALA, γ -linolenic acid or linoleic acid. Reduced gene expression was observed for the production of pro-inflammatory cytokines (TNF- α , IL-1 β , IL-6 and IL-10). Whereas, greater expression was also observed for PPAR- γ (anti-inflammatory).	Dipasquale et al., [32]
Mice	4 weeks	0, 0.5, and 1.5% DHA in presence and absence of 0.5% CLA	Dietary CLA lowered EPA-, DHA-, and ALA-derived epoxides, PGF1 α , PGF2 α , and F2-isoprostanes. Overall, it was concluded that CLA elevated proinflammatory oxylipins, while DHA increased anti-inflammatory oxylipins and weakened the effect of CLA-induced pro-inflammatory oxylipins in adipose tissue.	Adkins et al., [33]
Human patients with Crohn's disease (n=13)	12 weeks	6g/d (orally)	Orally CLA intake suppressed peripheral blood T cells to produce pro-inflammatory cytokines (IFN-g, TNF-a and IL-17), reduced disease activity and increased the quality of life of patients with Crohn's disease. Oral administration of CLA did not show any negative effects and was well tolerated.	Bassaganya-Riera et al., [34]
Human (n=90)	2 months	3 g/d	The level of high sensitivity C-reactive protein (hsCRP), IL-6 and malondialdehyde (MDA), which are the markers of inflammation and oxidative stress, decreased significantly from 7.48 to 5.95 mg/ml, 16.13 to 12.95 pg/ml and 3.7 to 2.4 mol/l, respectively in CLA group after 8 weeks. However, glutathione peroxidase (GPx) increased from 125 \pm 46.06 (week 0) to 171.4 \pm 68.90 (week 8) mmol/ml/min.	Eftekhari et al., [35]
Human (n=29) Healthy adults	8 weeks	20 g/day of butter enriched with CLA (1020 \pm 167 mg CLA/day)	Intake of CLA-enriched butter resulted in increased levels of anti-inflammatory IL-10 and reduced levels of pro inflammatory components (TNF α , IL-2, IL-8 and transcription factor NF κ B).	Penedo et al., [36]

Subject	Duration	CLA concentration in diet	Major findings	Reference
			However, there was no significant effect on adiponectin, body composition, C-reactive protein, and IL-4 when compared to control group.	
Human (n=10)	10 weeks	200 g/week of cheese enriched with CLA	Dietary supplementation of CLA enriched cheese exhibited significant ($p < 0.05$) reduction in pro-inflammatory markers such as IL-6 (4.58 ± 0.94 vs. 8.08 ± 1.57 pg/mL), IL-8 (28.59 ± 2.64 vs. 45.02 ± 5.82 pg/mL), and TNF- α (32.09 ± 17.42 vs. 53.58 ± 25.67 pg/mL). Moreover, a substantial reduction was also observed in the extent of platelet aggregation (77.7 ± 3.56 vs. $87.8 \pm 1.76\%$).	Sofi et al., [37]
Mice	2 weeks	0.5- 1% CLA-supplemented diet	Data revealed that wound healing rate was significantly ($P < 0.05$) faster in group fed 1% CLA supplemented diet as compared to control and/or group fed 0.5% CLA-diet. However, no significant difference was observed between the wound healing rate between 0.5% CLA group and control. Overall, results showed that CLA supplementation reduced the levels of oxidative stress and inflammatory markers in experimental animals.	Park et al., [38]
Human cells (Human THP-1 monocytes) (In vitro)	NR	NR	Suppressed levels of pro-inflammatory markers such as macrophage phenotype; reduced expression of cyclooxygenase (COX)-2 and cytosolic phospholipase-A2 (cPLA2) and monocyte chemoattractant protein-1 (MCP-1); reduced levels of prostaglandin E2 (PGE2) and matrix metalloprotease (MMP)-9. Overall, it was concluded that CLA reduce the inflammatory outputs of macrophages.	McClelland et al., [39]
Mice	3 weeks	1% CLA supplemented diet	When colitis-induced mice were fed with CLA-based diet, a significant suppression was observed in weight loss, signs of colitis and inflammatory infiltration than that of control. Contrarily, when the mice with no colitis fed with CLA-based diet, a condition 'steatosis' was developed in which lipid metabolism by liver gets impaired resulting abnormal accumulation of fats in cells and organs. Results suggested that CLA is safe for use during gut inflammation but not at steady state conditions.	Moreira et al., [40]

Table 4. Major findings of clinical and animal studies based on cholesterol-lowering and anti-obesity effects of CLA

Subject	Duration of feeding	CLA Concentration in diet	Major findings	Reference
Cholesterol-lowering effects				
Male, female	9 weeks	20 g/day (oleic acid: CLA::80:20)	High intakes of an 80:20 mixture of cis-9, trans-11 and trans-10, cis-12 CLA raised the total- to HDL-cholesterol ratio in healthy volunteers.	Wander et al., 2010 [41]
Rats	6 weeks	0.5% trans-10, cis-12 CLA	Resveratrol and CLA significantly reduced body fat but did not do so when combined.	Arise et al.,2011 [42]
Rats	100 days	0.5% trans–trans CLA isomers	The trans–trans CLA-rich soy oil lowered the serum cholesterol and low-density lipoprotein–cholesterol levels by 41 and 50%, respectively, also lowered the liver lipid content and decreased the liver weight in the obese rats.	Gilbert et al.,2011 [43]
Rats	21 days	1% CLA and 63.2% of fructose	CLA in high-fructose diet, decreases serum LDL + VLDL and TG and plasma MDA concentrations as well as liver weight and liver cholesterol.	Kostogrys et al., 2010 [44]
Mice	4 weeks	0.4 g of CLS mixed with 1 kg of hyperlipidemic feed or 5 g every day	Decreased serum total cholesterol (TC), serum triacylglycerols (TAGs), serum low-density lipoprotein cholesterol (LDL-C), atherogenic index (AI), liver weight (LW), liver index (LI), liver TC, and TAGs of mice.	Li, R et al., 2010 [45]
Anti-obesity effects				
Children (6 to 10 year)	2 weeks	3 g/day (50:50 cis-9, trans-11 and trans-10, cis-12 isomers)	CLA supplementation decreased body fatness in 6–10 years old children who were overweight or obese	Racine et al., 2010 [46]
Mice	14 days	1% trans-10, cis-12-CLA	Trans-10, cis-12 isomer of conjugated linoleic acid (CLA) caused a rapid reduction of body and adipose mass in mice.	Ashwell et al., 2010 [47]
Male, female (human)	12 weeks	1.7 g CLA in 200 ml sterilized milk	CLA significantly decreased the body weight, body mass index, body fat mass, fat percentage, subcutaneous fat mass, and waist-to-hip ratio in overweight.	Chen et al., 2012 [48]
Rat	8 weeks	2% of the 50:50 (c-9, t-11: t-10, c-12) mixture of CLA and 2% phytosterols	Diet supplementations with CLA, phytosterols or their combination for 65 days were effective in reducing body fat, adipose tissue and feed consumption, and CLA, but not phytosterols, modulated the action of leptin in obesity.	Furlan et al., 2013 [49]

Table 5. Intended food uses and maximum use-levels of CLA in various food products (USDA, 2006 [50])

Food Category	Intended food use	Maximum CLA level (g/ serving)	Maximum Use-level (%)
Beverages & beverage bases	Specific soy milk beverages	1.5	0.625
Grain products and pasta	Meal replacement bars	1.5	3.750
Milk and milk products	Flavored milk products	1.5	0.625
	Milk	1.5	0.625
	Specific yoghurt products	1.5	0.667
Processed fruits and fruit juices	Specific fruit juice products	1.5	0.625

milk-based fermented beverages, ice-creams, grains and pasta products, processed fruits and fruit juices, etc. Campbell et al., [27] fortified CLA oil (Clairinol G-80) in skim milk at the concentration of 1 & 2%. After HTST pasteurization, the concentration of cis-9/trans 11 isomers remains stable by 2 weeks of refrigerated storage. Sensory analysis revealed that in fortified milks, low intensities of a "grassy/vegetable oil" odor was perceived. Rodriguez-Alcala and Fontecha [28] supplemented milk powder, milk, yoghurt, fermented milk, fresh cheese and milk-juice blend with Tonalin-80 (CLA-oil, 80% CLA). Major of CLA isomers were not affected by thermal treatment and processing but the total CLA content decreased in fresh samples after 10 weeks of low temperature storage. Kim et al. [1] has reported that the consumption of CLA up to 6 g/day for 1 year or 3.4 g/day for up to 2 years is currently being considered safe based on the outcomes of previous clinical studies.

Commercially available CLA-rich dietary supplements SafflorinTM

Safflorin is suitable as ingredient in dietary supplements in various forms, including softgels, capsules, and supplement bars and similar products. Chemically, SafflorinTM is a mixture of the cis-9, trans-11 [(better known as conjugated linoleic acid (CLA))] and trans-10, cis-12 isomers of octadecadienoic acid in a 40:60 ratio. Depending on the application, SafflorinTM can be supplied as free fatty acids or esterified in glyceride form. The latter form is tasteless and therefore suitable for use in dietary supplement bars and similar products. Physically, SafflorinTM is a clear, colorless to pale yellow liquid at ambient temperature, free from foreign odors or off flavors. It contains

≥74% CLA (cis-9, trans-12 & trans-10, cis-12) isomers.

Tonalin® TG80 and ClairinolTM G -80

CLA-Rich Oil is a food grade preparation derived from processed safflower oil. It consists of approximately 78% total conjugated linoleic acid (CLA) isomers and 74% of an approximately 50:50 mixture of cis-9, trans-11 and trans-10, cis-12 CLA isomers. CLA-rich oil is intended to be added to certain specified foods within the general categories of soy milk, meal replacement beverages and bars, milk products and fruit juices. CLA-rich oil would be added to these foods at a level of 1.5 g per serving.

Fortification of CLA-rich oils in foods

CLA-Rich Oil is intended for use in specific foods within the following general food categories:

- Beverages and beverage bases
- Grain and pasta products
- Milk and milk products
- Processed fruits and fruit juices

4. CONCLUSION

Conjugated linoleic acid has unique property of anti-carcinogenic activity.. Diverse physiological functions of CLA in preventing the risks of cancer, obesity, hypercholesterolemia, atherosclerosis, etc. studied in animal model are little convincing and require further intense research. However, a few attempts have been made to confirm the positive impacts of CLA on human; nevertheless, it is not clear whether CLA promotes all of these advantages to humans as well. Further research is required to establish the biological benefits of CLA's activity, to come about efficient strategies of intake and recommended amount for human safety.

5. LIMITATIONS

As far as CLA is concerned, most of the studies have been conducted on animals (rodents, pigs, dogs, etc.) to establish the health effects of CLA. A very few studies have been conducted on human beings. No clear/concrete mechanisms are established yet for its antioxidative, anticarcinogenic and weight-control activities. No RDA is given by any regulatory agency yet. Toxicological effects of CLA are lacking due to insufficient data and research. Further clinical studies are required to establish safe limit or dietary intake of CLA along with the health benefits.

ACKNOWLEDGEMENT

The authors are thankful to the Executive Director, Mansinhbhai Institute of Dairy and Food Technology (MIDFT) for his continuous support for writing the manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kim KJ, Lee J, Park Y, Lee SH. ATF3 mediates anti-cancer activity of trans-10, cis-12-conjugated linoleic acid in human colon cancer cells. *Biomol Ther.* 2015;23(2): 134. Available: <https://dx.doi.org/10.4062%2Fbio molther.2014.107>
2. Benjamin S, Prakasan P, Sreedharan S, Wright AD, Spener F. Pros and cons of CLA consumption: An insight from clinical evidences. *Nutr Metab.* 2015;12(1):1-20. Available: <https://doi.org/10.1186/1743-7075-12-4>.
3. Market watch global Conjugated Linoleic Acid (CLA) Market Growth 2019-2024; 2019. Available: <https://www.marketwatch.com/press-release/conjugated-linoleic-acid-cla-market-2019-industry-growth-size-share-global-forecasts-analysis-company-profiles-competitive-landscape-and-key-regions-analysis-research-report-2019-04-17> Last accessed 20/12/2019
4. Chin SF, Liu W, Storkson JM, Ha YL, Pariza MW. Dietary sources of conjugated dienoic isomers of linoleic acid, a newly recognized class of anticarcinogens. *J Food Comp Anal.* 1992;5(3):185-197. Available: [https://doi.org/10.1016/0889-1575\(92\)90037-K](https://doi.org/10.1016/0889-1575(92)90037-K)
5. Bauman DE, Baumgard LH, Corl BA, Griinari DJ. Biosynthesis of conjugated linoleic acid in ruminants. *J Anim Sci.* 2000; 77:1–15. Available: <https://doi.org/10.2527/jas2000.77E-Suppl1f>
6. Harfoot GC, Hazlewood GP. Lipid metabolism in the rumen. In: Hobson PN, Stewart CS (Eds.) *The Rumen Microbial Ecosystem* London: Blackie Academic and Professional. 1997;382–426.
7. Soel SM, Choi OS, Bang MH, Park JHY, Kim WK. Influence of conjugated linoleic acid isomers on the metastasis of colon cancer cells in vitro and in vivo. *J Nutr Biochem.* 2007;18(10):650-657. Available: <https://doi.org/10.1016/j.jnutbio.2006.10.011>
8. Evans NP, Misyak SA, Schmelz EM., Guri AJ, Hontecillas R, Bassaganya-Riera J. Conjugated linoleic acid ameliorates inflammation-induced colorectal cancer in mice through activation of PPAR γ . *J Nutr* 2010;140(3):515-521. Available: <https://doi.org/10.3945/jn.109.115642>
9. Sharma V, Goyal A, Sihag M. Role of omega-3 fatty acids, conjugated linoleic acid and phytosterols in improving functionality of dairy products. *Lecture compendium on national training on innovative trends in dairy and food products formulation.* National Dairy Research Institute, India. 2012;10:126-30.
10. Mohammadzadeh M, Faramarzi E, Mahdavi R, Nasirimotlagh B, Asghari Jafarabadi M. Effect of conjugated linoleic acid supplementation on inflammatory factors and matrix metalloproteinase enzymes in rectal cancer patients undergoing chemoradiotherapy. *Integr Cancer Ther* 12(6):496-502. Available: <https://doi.org/10.1177%2F1534735413485417>
11. Pierre AS, Minville-Walz M, Fèvre C, Hichami A, Gresti J, Pichon L, Bellenger S, Bellenger J, Ghiringhelli F, Narce M, Riolland M. Trans-10, cis-12 conjugated linoleic acid induced cell death in human colon cancer cells through reactive oxygen species-mediated ER stress. *BBA Mol Cell.* 2013;1831(4):759-768. Available: <https://doi.org/10.1016/j.bbailip.2013.01.005>

12. Cheng G, Zhang X, Chen Y, Lee RJ, Wang J, Yao J, Zhang Y, Zhang C, Wang K and Yu B. Anticancer activity of polymeric nanoparticles containing linoleic acid-SN38 (LA-SN38) conjugate in a murine model of colorectal cancer. *Colloids Surf B Bio-interfaces*. 2019;181:822-829. Available:<https://doi.org/10.1016/j.colsurfb.2019.06.020>
13. Devery R, Miller A, Stanton C. Conjugated linoleic acid and oxidative behaviour in cancer cells. *Biochem Soc Trans*. 2001; 29(2):341-345. Available:<https://doi.org/10.1042/bst0290341>
14. Coakley M, Johnson MC, McGrath E, Rahman S, Ross RP, Fitzgerald GF, Devery R, Stanton C. Intestinal bifidobacteria that produce trans-9, trans-11 conjugated linoleic acid: A fatty acid with anti-proliferative activity against human colon SW480 and HT-29 cancer cells. *Nutr Cancer*. 2006; 56(1):95-102. Available:https://doi.org/10.1207/s15327914nc5601_13
15. Kadirareddy RH, Mathakala VANI, Palempalli UD. Therapeutic activity of conjugated linoleic acids synthesized by *Lactobacillus plantarum*. *Int J Pharma Bio Sci*. 2016;7(4):215-23. Available:<https://doi.org/10.22376/ijpbs.2016.7.4.b215-223>
16. Cunningham ML, Lokesh BR. Superoxide anion generated by potassium superoxide in cytotoxic and mutagenic to Chinese Hamster Ovary cells. *Mutat Res*. 1983; 121:299-304. Available:[https://doi.org/10.1016/0165-7992\(83\)90218-X](https://doi.org/10.1016/0165-7992(83)90218-X)
17. Basiricò L, Morera P, Dipasquale D, Tröscher A, Serra A, Mele M, Bernabucci U. Conjugated linoleic acid isomers strongly improve the redox status of bovine mammary epithelial cells (BME-UV1). *J Dairy Sci*. 2015;98(10):7071-7082. Available:<https://doi.org/10.3168/jds.2015-9787>
18. Ip CS, Banni E, Angioni G, Carta J, Macginley HJ, Thompson D, Barbano P, Bauman D. Conjugated Linoleic Acid enriched butter fat alters mammary gland morphogenesis and reduces cancer risk in rats. *J Nutr*. 1999;129:2135-2142. Available:<https://doi.org/10.1093/jn/129.12.2135>
19. Kritchevsky D. Conjugated Linoleic Acid. In *handbook of functional dairy products* Ed. Colette Shortt, John O'Brien. 2016;173-186. CRC Press.
20. Kathirvelan C. Influence of mustard cake feeding on conjugated linoleic acid content of buffalo milk and its anticarcinogenic potency. Ph. D thesis, National Dairy Research Institute (NDRI) Karnal India; 2007.
21. Wannamethee SG, Jefferis BJ, Lennon L, Papacosta O, Whincup PH, Hingorani AD Serum conjugated linoleic acid and risk of incident heart failure in older men: The British regional heart study. *Am Heart J Assoc*. 2018;7(1):006653. Available:<https://doi.org/10.1161/JAHA.117.006653>
22. Gray C, Vickers MH, Segovia SA, Zhang XD, Reynolds CM. A maternal high fat diet programmes endothelial function and cardiovascular status in adult male offspring independent of body weight, which is reversed by maternal conjugated linoleic acid (CLA) supplementation. *PLoS One* 2015;10(2):0115994. Available:<https://doi.org/10.1371/journal.pone.0115994>
23. Limongi F, Noale M, Marseglia A, Gesmundo A, Mele M, Banni S, Crepaldi G, Maggi S. Impact of cheese rich in Conjugated Linoleic Acid on low density lipoproteins cholesterol: Dietary Intervention in Older People (CLADIS Study). *J Food Nutr Res*. 2018;6(1):1-7.
24. Lee KN, Kritchevsky D, Pariza MW. Conjugated Linoleic Acid and atherosclerosis in rabbits. *Atherosclerosis* 1994; 108:19-25. Available:[https://doi.org/10.1016/0021-9150\(94\)90034-5](https://doi.org/10.1016/0021-9150(94)90034-5)
25. Nicolosi RJ, Rodgers EJ, Kitchevsky D, Scimeca JA, Huth PJ. Dietary Conjugated Linoleic Acid reduces plasma lipoproteins and early aortic atherosclerosis in hypercholesterolaemic hamsters. *Artery*. 1997; 22(5):266-277.
26. Watkins BA, Li Y, Seifert MF. Bone metabolism and dietary Conjugate Linoleic Acid. In: *Advances in CLA Research*. Ed. MP Yurawecz, MM Mossoba, JKG Kramer, MW Pariza, Nelson GJ. AOCS Press Champaign IL. 1999;1:253-255.
27. Campbell W, Drake MA, Larick DK. The impact of fortification with conjugated linoleic acid (CLA) on the quality of fluid milk. *J Dairy Sci*. 2003;86(1):43-51. Available:[https://doi.org/10.3168/jds.S0022-0302\(03\)73582-6](https://doi.org/10.3168/jds.S0022-0302(03)73582-6)

28. Rodríguez-Alcalá LM, Fontecha J. Fatty acid and Conjugated Linoleic Acid (CLA) isomer composition of commercial CLA-fortified dairy products: Evaluation after processing and storage. *J Dairy Sci* 2007;90(5):2083-2090. Available: <https://doi.org/10.3168/jds.2006-693>
29. Beppu F, Hosokawa M, Tanaka L, Kohno H, Tanaka T, Miyashita K. Potent inhibitory effect of trans9, trans11 isomer of conjugated linoleic acid on the growth of human colon cancer cells. *J Nutr Biochem* 2006;17(12):830-836. Available: <https://doi.org/10.1016/j.jnutbio.2006.01.007>
30. El Roz A, Bard JM, Huvelin JM, Nazih H. The anti-proliferative and pro-apoptotic effects of the trans9, trans11 conjugated linoleic acid isomer on MCF-7 breast cancer cells are associated with LXR activation. *Prostag Leukotr Ess (PLEFA)*. 2013; 88(4):265-272. Available: <https://doi.org/10.1016/j.plefa.2012.12.006>
31. Villacorta L, Minarrieta L, Salvatore SR, Khoo NK, Rom O, Gao Z, Berman RC, Jobbagy S, Li L, Woodcock SR and Chen YE. *In situ* generation, metabolism and immunomodulatory signaling actions of nitro-conjugated linoleic acid in a murine model of inflammation. *Redox Biol*. 2018; 15:522-531. Available: <https://doi.org/10.1016/j.redox.2018.01.005>
32. Dipasquale D, Basirico L, Morera P, Primi R, Tröscher A, Bernabucci U. Anti-inflammatory effects of conjugated linoleic acid isomers and essential fatty acids in bovine mammary epithelial cells. *Animal* 2018;12(10):2108-2114. Available: <https://doi.org/10.1017/S1751731117003676>
33. Adkins Y, Belda BJ, Pedersen TL, Fedor DM, Mackey BE, Newman J W, Kelley DS. Dietary docosahexaenoic acid and trans-10, cis-12-conjugated linoleic acid differentially alter oxylipin profiles in mouse periuterine adipose tissue. *Lipids*. 2017; 52(5):399-413. Available: <https://doi.org/10.1007/s11745-017-4252-3>
34. Bassaganya-Riera J, Hontecillas R, Horne WT, Sandridge M, Herfarth HH, Bloomfeld R, Isaacs KL. Conjugated Linoleic Acid modulates immune responses in patients with mild to moderately active Crohn's disease. *Clin Nutr*. 2012;31(5):721-727. Available: <https://doi.org/10.1016/j.clnu.2012.03.002>
35. Eftekhari MH, Aliasghari F, Babaei-Beigi MA, Hasanzadeh J. Effect of conjugated linoleic acid and omega-3 fatty acid supplementation on inflammatory and oxidative stress markers in atherosclerotic patients. *ARYA Atherosclerosis*. 2013; 9(6):311.
36. Penedo LA, Nunes JC, Gama MAS, Leite PEC, Quirico-Santos TF, Torres AG. Intake of butter naturally enriched with cis9, trans11 conjugated linoleic acid reduces systemic inflammatory mediators in healthy young adults. *J Nutr Biochem*. 2013; 24(12):2144-2151. Available: <https://doi.org/10.1016/j.jnutbio.2013.08.006>
37. Sofi F, Buccioni A, Cesari F, Gori AM, Minieri S, Mannini L, Casini A, Gensini GF, Abbate R, Antongiovanni M. Effects of a dairy product (pecorino cheese) naturally rich in cis-9, trans-11 conjugated linoleic acid on lipid, inflammatory and haemorheological variables: A dietary intervention study. *Nutr Metab Cardiovasc Dis*. 2010;20(2):117-124. Available: <https://doi.org/10.1016/j.numecd.2009.03.004>
38. Park NY, Valacchi G, Lim Y. Effect of dietary conjugated linoleic acid supplementation on early inflammatory responses during cutaneous wound healing. *Mediat Inflamm*. 2010;1-8. Available: <http://dx.doi.org/10.1155/2010/342328>
39. McClelland S, Cox C, O Connor R, de Gaetano M, McCarthy C, Cryan L, Fitzgerald D, Belton O. Conjugated Linoleic Acid suppresses the migratory and inflammatory phenotype of the monocyte/macrophage cell. *Atherosclerosis*. 2010; 211(1):96-102. Available: <https://doi.org/10.1016/j.atherosclerosis.2010.02.003>
40. Moreira TG, Gomes-Santos AC, Horta LS, Goncalves MC, Santiago AF, Lauar JG, dos Reis DS, Castro-Junior AB, Lemos L, Guimarães M, Aguilar EC. Consumption of conjugated linoleic acid (CLA)-supplemented diet during colitis development ameliorates gut inflammation without causing steatosis in mice. *J Nutr Biochem*. 2018;57: 238-245.

- Available:<https://doi.org/10.1016/j.jnutbio.2018.04.003>
41. Wanders AJ, Brouwer IA, Siebelink E, Katan MB. Effect of a high intake of conjugated linoleic acid on lipoprotein levels in healthy human subjects. *PLoS One*. 2010;5(2):9000. Available:<https://doi.org/10.1371/journal.pone.0009000>
 42. Arias N, Macarulla, MT, Aguirre L, Martínez-Castaño MG, Gómez-Zorita S, Miranda J, Martínez JA, Portillo MP. The combination of resveratrol and conjugated linoleic acid is not useful in preventing obesity. *J Physiol Biochem*. 2011;67(3):471-477. Available:<https://doi.org/10.1007/s13105-011-0086-2>
 43. Gilbert W, Gadang V, Proctor A, Jain V, Devareddy L. Trans-trans Conjugated linoleic acid enriched soybean oil reduces fatty liver and lowers serum cholesterol in obese Zucker rats. *Lipids*. 2011;46(10):961-968. Available:<https://doi.org/10.1007/s11745-011-3585-6>
 44. Kostogrys RB, Pisulewski PM. Effect of conjugated linoleic acid (CLA) on lipid profile and liver histology in laboratory rats fed high-fructose diet. *Environ Toxicol Pharmacol*. 2010;30(3):245-250. Available:<https://doi.org/10.1016/j.etap.2010.06.006>
 45. Li R, Jia CS, Yue L, Zhang XM, Xia QY, Zhao SL, Feng B, Zhong F, Chen WJ. Lipase-catalyzed synthesis of conjugated linoleyl β -sitosterol and its cholesterol-lowering properties in mice. *J Agri Food Chem* 2010;58(3):1898-1902. Available: <https://doi.org/10.1021/jf902943y>
 46. Racine NM, Watras AC, Carrel AL, Allen DB, McVean JJ, Clark RR, O'Brien AR, O'Shea M, Scott CE, Schoeller DA. Effect of conjugated linoleic acid on body fat accretion in overweight or obese children. *Am J Clin Nutr*. 2010;91(5):1157-1164. Available:<https://doi.org/10.3945/ajcn.2009.28404>
 47. Ashwell MS, Ceddia RP, House RL, Cassady JP, Eisen EJ, Eling TE, Collins JB, Grissom S F, Odle J. Trans-10, cis-12-conjugated linoleic acid alters hepatic gene expression in a polygenic obese line of mice displaying hepatic lipodosis. *J Nutr Biochem*. 2010;21(9):848-855. Available:<https://doi.org/10.1016/j.jnutbio.2009.06.013>
 48. Chen SC, Lin YH, Huang HP, Hsu WL, Houg JY, Huang CK. Effect of conjugated linoleic acid supplementation on weight loss and body fat composition in a Chinese population. *Nutr*. 2012;28(5):559-565. Available:<https://doi.org/10.1016/j.nut.2011.09.008>
 49. Furlan CPB, da Silva Marineli R, Júnior MRM. Conjugated linoleic acid and phytosterols counteract obesity induced by high-fat diet. *Food Res Int*. 2013;51(1):429-435. Available:<https://doi.org/10.1016/j.foodres.2012.12.023>
 50. USDA. What We Eat In America: National Health and Nutrition Examination Survey (NHANES): 2003-2004 U.S Department of Agriculture (USDA); Riverdale, Maryland; 2006. Available:http://www.ars.usda.gov/research/publications/publications.htm?seq_no_115=199902 Last accessed on 01/01/2020

© 2019 Virsangbhai et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/53785>