



# **Studies on the Response of Hybrid Maize to IPNSS on Productivity, Nutrient Uptake and Post Harvest Soil Status**

**R. Jayasoorya<sup>a\*</sup>, P. Stalin<sup>a</sup>, P. Anandan<sup>a</sup>, C. Prabakar<sup>a</sup>, R. Gobi<sup>a</sup>,  
G. Murugan<sup>a</sup> and P. Naveen Prasath<sup>a</sup>**

<sup>a</sup> *Department of Agronomy, Faculty of Agriculture, Annamalai University, Annamalai Nagar, Tamil Nadu-608002, India.*

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/IJECC/2022/v12i111416

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/93285>

**Received 29 August 2022**

**Accepted 31 October 2022**

**Published 04 November 2022**

**Original Research Article**

## **ABSTRACT**

A field study was carried out at farmer's field, Kalappanaickenpatti village, Sendamangalam block, Namakkal district of Tamil Nadu during July - October 2021 to study the effect of Integrated plant nutrient supply system (IPNSS) for sustaining yield of hybrid maize variety, KAVERI SUPER – 244 in clay loam soil. There are ten different treatments with three replications each were carried out in the plot RBD design. IPNSS practice including inorganic fertilizers, organic manures and biofertilizers in which treatment (T<sub>8</sub>) 100 per cent RDF + CPM @ 3 t ha<sup>-1</sup> + Micronutrient mixture (25 kg ha<sup>-1</sup>) + Azophos (20 packets) showed its best results with respect to soil properties and yield parameters like cob length, cob diameter, number of grain cob<sup>-1</sup>, test weight and harvest index.

**Keywords:** *Maize; IPNSS; composted poultry manure (CPM).*

## **1. INTRODUCTION**

“Globally, Maize is the third most important food crop after rice and wheat. Maize is one of the

most versatile crop and is grown in more than 166 countries across the world, including tropical, subtropical and temperate regions. India produced 30 million tonnes in an area of 9.9

\*Corresponding author: E-mail: jayasoorya1998@gmail.com;

million hectares in 2020-21" [1]. Maize is less water demanding and gives higher yield per hectare. By growing maize farmers can save 90 per cent of water, 70 per cent of power compared to Paddy.

In addition to staple food for human being and quality feed for animals, maize serves as a basic raw material to thousands of industrial products that includes starch, oil, protein, alcoholic beverages, food sweeteners, pharmaceutical, cosmetic, film, textile, gum, package, paper industries etc. However, over application of inorganic fertilizers causes inefficient use, large losses and imbalances of nutrients. It also leads to environmental contamination in a number of areas in developed world.

Hence, it is important to reduce the usage of chemical fertilizers and supplementing the nutrient need same through organic manures such as CPM, FYM, pressmud compost, vermicompost and biofertilizers have become necessary to attain sustainable productivity, profitability and to maintain soil health.

Furthermore, the fertilizer management is one of the most important factors that influence the growth and yield of maize crop. Maize is considered as most exhaustive crop after sugar cane and requires both micro and macro nutrients to obtain high growth and yield potentials. In fact, organic nutrients not only provide plant with nutrients but also improve and or sustain the soil health. The micronutrients content in organic manure may be sufficient enough to meet the crop production requirement.

## 2. MATERIAL AND METHODS

A field experiment entitled "Integrated plant nutrient supply system for sustaining yield of hybrid maize" was conducted during *kharif* season at farmer's field, Kalappanaickenpatti village, Sendamangalam block, Namakkal district of Tamil Nadu during July - October 2021.

The soil of experimental site was sandy clay loam with pH of 7.1, Electrical conductivity 0.38 dS m<sup>-1</sup>, medium in organic carbon (0.57 per cent), low in available nitrogen (181 kg ha<sup>-1</sup>) and medium in phosphorus (18.5 kg ha<sup>-1</sup>) and medium in potassium (230 kg ha<sup>-1</sup>).

The experiment was laid out in a randomized block design for maize during *kharif* 2021 with 10

treatments consisting of T<sub>1</sub> - Control (No application), T<sub>2</sub> - 100% Recommended Dose of Fertilizer (RDF), T<sub>3</sub> - 100% RDF + Vermicompost @ 3 t ha<sup>-1</sup>, T<sub>4</sub> - 100% RDF + CPM @ 3 t ha<sup>-1</sup>, T<sub>5</sub> - 100% RDF + Pressmud compost @ 3 t ha<sup>-1</sup>, T<sub>6</sub> - 100% RDF + Farm yard manure @ 10 t ha<sup>-1</sup>, T<sub>7</sub> - 100% RDF + Vermicompost @ 3 t ha<sup>-1</sup> + Micronutrient mixture + Azophos, T<sub>8</sub> - 100% RDF + CPM @ 3 t ha<sup>-1</sup> + Micronutrient mixture + Azophos, T<sub>9</sub> - 100% RDF + Pressmud compost @ 3 t ha<sup>-1</sup> + Micronutrient mixture + Azophos, T<sub>10</sub> - 100% RDF + Farm yard manure @ 10 t ha<sup>-1</sup> + Micronutrient mixture + Azophos.

The hybrid maize Kaveri super - 244 was sown at spacing of 60 cm x 25 cm. Organic manures were applied as per the treatment and incorporated into the soil two weeks before sowing. The fertilizers were applied to the experimental field as per the recommended fertilizer schedule of 250:75:75 kg N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O ha<sup>-1</sup>. The nitrogen source was given in three splits in the form of urea, phosphorous as single super phosphate and potassium as muriate of potash. A half dose of N and full dose of P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O were applied basally and the remaining half doses of N were applied as two splits on 25 and 45 days after sowing. In addition, as per the treatment schedule the 20 packets of Azophos were applied initially in the soil.

The weather at Kalappanaickenpatti village, Namakkal is moderately warm with hot summer months. Namakkal district comes under north western agro climatic zone and western agro climatic zone. The weekly maximum temperature mean is about 32.6°C during the cropping year of 2021. The weekly minimum temperature mean is about 24.4°C during the cropping year of 2021. The average relative humidity was 80 per cent during 2021.

## 3. RESULTS AND DISCUSSION

### 3.1 Yield Parameters

The yield potential of maize is determined by the resultant values of yield components which are greatly influenced by the growth parameters. This is well reflected in the present investigation also. Almost all yield attributing characters *viz.*, cob length, cob diameter, number of grains cob<sup>-1</sup>, test weight and harvest index were remarkably influenced by the practice of integrated plant nutrient supply system. Among the various treatments imposed in the study of integrated

application of 100 percent RDF + CPM @ 3 t ha<sup>-1</sup> + Micronutrient mixture + Azophos (T<sub>8</sub>) registered an increased in yield parameters.

“The number of grains cob<sup>-1</sup> of maize obtained from plants that received poultry manure was because of more nutrients were released which mineralized rapidly for plant uptake and utilization. It could also be attributed to reduction in plant diseases or other soil-borne diseases which decrease yield, or to increased microbial activities which favour yield increases” as reported by Olanikan [2] and Enujoke et al. [3]. These reports are generally in harmony with the work and recommendations of Directorate of Information and Publications of Agriculture [4] and Fagimi and Odebode [5]. The higher yield attributes under combined application of organics and inorganics was due to higher levels of inorganic fertilizer that have increased the activity of photosynthesis and enzymes which is responsible for transformation of energy, carbohydrates, fat metabolism and respiration of the plant. Number of grains cob<sup>-1</sup> and test weight increased linearly with the application of poultry manure. Similar results are reported by Sudhagar Rao et al. [6].

### 3.2 Nutrient Uptake and Post Harvest Soil Status

It was quite clear from the data that the application of 100 percent RDF + CPM @ 3 t ha<sup>-1</sup> + Micronutrient mixture + Azophos (T<sub>8</sub>) recorded the maximum nitrogen, phosphorous and potassium in uptake by the plant and post harvest available soil nutrient status after harvest.

The uptake of phosphorus might be attributed to the reason of the positive effect of organic sources viz., poultry manure and Azophos which chelates micronutrients (Ca and Mg) and preventing them from fixing phosphorus in insoluble compounds and thereby increased the phosphorus uptake.

The potassium uptake in all fertilizers and manurial treatments increased significantly over control plot. This might be due to more availability of the nutrient potassium from added fertilizers and to the solubilizing action on potassium by organic acids produced during degradation of organic manures. Similar findings were quoted by Altaf et al. [7].

**Table 1. Effect of IPNSS on cob length (cm), cob diameter (cm), number of grain cob<sup>-1</sup>, test weight (g) and harvest index of hybrid maize**

Treatments	Cob length	Cob diameter	Number of grain cob <sup>-1</sup>	Test weight	Harvest index
T <sub>1</sub> - Control	13.28	5.18	175.10	24.83	31.06
T <sub>2</sub> - 100% RDF	15.75	5.42	295.63	25.27	38.05
T <sub>3</sub> - 100% RDF + Vermicompost @ 3 t ha <sup>-1</sup>	19.64	6.06	370.01	25.79	39.99
T <sub>4</sub> - 100% RDF + CPM @ 3 t ha <sup>-1</sup>	20.47	6.28	410.24	25.91	41.39
T <sub>5</sub> - 100%RDF + Pressmud compost @ 3 t ha <sup>-1</sup>	19.13	5.97	360.66	25.48	39.67
T <sub>6</sub> - 100% RDF + FYM @ 10 t ha <sup>-1</sup>	17.64	5.73	330.44	25.34	38.48
T <sub>7</sub> - 100%RDF + Vermicompost @ 3 t ha <sup>-1</sup> + Micronutrient mixture + Azophos	23.17	6.9	438.99	26.23	42.87
T <sub>8</sub> - 100% RDF +CPM @ 3 t ha <sup>-1</sup> + Micronutrient mixture +Azophos	24.29	7.20	454.11	26.34	43.30
T <sub>9</sub> - 100% RDF+ Pressmud compost @ 3 t ha <sup>-1</sup> + Micronutrient mixture + Azophos	22.72	6.78	431.48	26.15	42.63
T <sub>10</sub> - 100% RDF+ FYM @ 10 t ha <sup>-1</sup> + Micronutrient mixture + Azophos	21.83	6.53	418.35	26.02	41.79
SEm±	0.18	0.06	3.90	0.005	0.12
CD (P=0.05)	0.55	0.16	11.93	NS	0.34

**Table 2. Effect of IPNSS on nutrient uptake and Post harvest nutrient status of soil (kg ha<sup>-1</sup>) of hybrid maize**

Treatments	Nutrient uptake			Post harvest nutrient status of soil		
	N	P	K	N	P	K
T <sub>1</sub> - Control	99.80	23.70	101.70	156.40	15.90	210.15
T <sub>2</sub> - 100% RDF	122.47	29.96	125.63	164.66	17.66	215.50
T <sub>3</sub> - 100% RDF + Vermicompost @ 3 t ha <sup>-1</sup>	157.03	41.45	145.03	170.94	19.65	224.33
T <sub>4</sub> - 100% RDF + CPM @ 3 t ha <sup>-1</sup>	169.64	44.85	152.43	173.64	20.39	228.43
T <sub>5</sub> - 100%RDF + Pressmud compost @ 3 t ha <sup>-1</sup>	151.63	40.13	141.23	170.14	19.36	223.52
T <sub>6</sub> - 100% RDF + FYM @ 10 t ha <sup>-1</sup>	137.47	34.73	134.33	166.99	18.50	219.31
T <sub>7</sub> - 100%RDF + Vermicompost @ 3 t ha <sup>-1</sup> + Micronutrient mixture + Azophos	198..99	54.47	171.40	180.21	21.84	237.14
T <sub>8</sub> - 100% RDF +CPM @ 3 t ha <sup>-1</sup> + Micronutrient mixture +Azophos	210.12	57.80	178.50	183.12	22.50	240.00
T <sub>9</sub> - 100% RDF+ Pressmud compost @ 3 t ha <sup>-1</sup> + Micronutrient mixture + Azophos	194.89	53.45	168.29	179.31	21.65	236.13
T <sub>10</sub> - 100% RDF+ FYM @ 10 t ha <sup>-1</sup> + Micronutrient mixture + Azophos	182.67	49.35	160.63	176.30	20.98	233.04
SEm±	2.76	0.62	1.83	0.50	0.14	0.66
CD (P=0.05)	8.24	1.86	5.47	1.51	0.44	1.98

“The presence of nitrogen fertilizer and poultry manure helps the soil to maintain higher rate of applied urea intact or in the form of ammonium ions for a longer time, resulting in higher nitrogen uptake efficiency”. The above results are in accordance with the earlier findings of Naiji et al. [8].

Application of inorganic fertilizers and poultry manure increased the nutrient uptake over control or in combination improves the availability of major nutrients to the maize. This was due to the addition of organic manure which improved the soil properties, hence, enhancing nitrogen, phosphorus and potassium uptake. This result is in agreement with Vidyavathi et al. [9].

Combined application of inorganic mineral fertilizer and poultry manure has significantly higher nitrogen, phosphorus and potassium uptake values of maize than the sole organic and inorganic fertilizers. A similar result was also documented by Quansah [10].

The increased nitrogen content in this treatment could attribute to the addition of organic material viz., poultry manure with balanced chemical fertilizers. These results are in close conformity with the findings of Sarangthem et al. [11].

An increase in soil available phosphorus in poultry manure treatments may be due to

mineralisation of organic phosphorus and production of organic acids that makes soil phosphorus more available and reduces phosphorus fixation. This result corroborates the findings of Islam et al. [12], who explained that “soils treated with either organic fertilisers or, inorganic fertilisers or combination of these fertilizers have higher values of available phosphorus will be obtained compared to the control treatments”.

The increase in soil exchangeable potassium in combined application of inorganic and organic fertilizer treatments may be due to the direct supplementation of potassium to soil which increases the potassium pool in the soil. This result is supported by Islam et al. [12] who reported that soil exchangeable potassium increased when organic or both organic and chemical fertilizers were applied. Ahmad et al. [13] also reported that “the highest soil OM, total nitrogen, available phosphorus and exchangeable potassium, after maize was harvested”.

#### 4. CONCLUSION

Based on the results the integration of composted poultry manure, inorganic fertilizer and biofertilizer (T<sub>8</sub>) 100% RDF + CPM @ 3 t ha<sup>-1</sup> + Micronutrient mixture + Azophos improved all the yield parameters and soil properties. The

combined uses of organic and inorganic source of nutrients have the improvement of crop yield parameters and soil physio-chemical properties. This kind of nutrients integration will produce the crop in sustainable manner. Whereas, the lowest values were recorded in the treatment (T<sub>1</sub>) which is control.

## ACKNOWLEDGEMENT

The authors wish to acknowledge the Annamalai University, Annamalai Nagar- 608002, Tamil Nadu.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. USDA. Foreign agricultural services office of global analysis report: USA; 2022. Available:<https://www.firs.usda.gov/data/world-agricultural-production>.
2. Olanikan, PC. Organic manures as soil amendment in eroded tropical soil of South Western Nigeria. *J. Trop. Agric.* 2006; 5:11-18.
3. Enujeke EC, Ojeifo IM, Nnaji GU. Residual effects of organic manure an inorganic fertilizer on maize grain weight and some soil properties in Asaba area of Delta State. *Int. J. Adv. Biol. Biomed. Res.* 2013; 3(3):433-442.
4. Directorate of Information and Publications of Agriculture. Handbook of Agriculture: facts and figures for farmers, students and all interested in farming. Indian Council of Agricultural Research, New Delhi. 2006;435.
5. Fagimi AA, Odebode CA. Effect of Poultry Manure on Pepper Veinal Mottle Virus (PVMV), yield and agronomic parameters of Pepper (*Capsicum annum*) in Nigeria. *East Afr. J. Sci.* 2007;1(2): 104-111.
6. Sudhagar Rao GB, Rex Immanuel R, Suthin Raj T, Pushpanathan KR. Analysis of growth and development of maize in response to integration of poultry manure with chemical fertilizers in maize (*Zea mays* L.). *J. Emerg. Technol. Innov. Res.* 2017;4(7):278-283.
7. Altaf MA, Shahid R, Altaf MA, Ren MX, Tan K, Xiang WQ, Qadir A, Shakoor A Altaf MM. Effect of NPK, organic manure and their combination on growth, yield and nutrient uptake of Chilli (*Capsicum annum* L.). *Hortic. Int j.* 2019;3(5):217-222.
8. Najji M, Sourji MK. Nutritional value and mineral concentrations of sweet basil under organic compared to chemical fertilization. *Acta Sci. Pol. Hortorum Cultus.* 2018;17(2):167-175.
9. Vidyavathi V, Dasog G, Babalad H, Hebsur NS, Gali SK, Patil SG, Alagawadi AR. Nutrient status of soil under different nutrient and crop management practices. *Karnataka J. Agric. Sci.* 2012; 25(2):1-9.
10. Quansah GW. Effect of organic and inorganic fertilizers and their combinations on the growth and yield of maize in the semi-deciduous forest zone of Ghana, M.Sc Thesis, Submitted to the Department of Crop and Soil Sciences, College of Agriculture and Natural Resources, Kwame Nkrumah University of Science And Technology, Kumasi, Ghana; 2010.
11. Sarangthem I, Mishra AD, Chakraborty Y. Cabbage productivity, nutrient uptake and soil fertility as affected by organic and bio-sources. *Agric. Sci. Digest.* 2011;31: 260-264.
12. Islam R, Sultana T, Haque A, Hossain I, Sabrin N, Islam R. Growth and yield of chilli influenced by nitrogen and phosphorus. *J. Agric. Vet. Sci.* 2018;11(5):54-68.
13. Ahmad W, Shah Z, Khan F, Ali S, Malik W. Maize yield and soil properties as influenced by integrated use of organic, inorganic and bio fertilizers in a low fertility soil. *Soil Environ.* 2013; 32(2):121-129.

© 2022 Jayasoorya et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/93285>