



Stellar Velocity Dispersion of a Massive Quenching Galaxy at $z = 4.01$

Masayuki Tanaka^{1,2}, Francesco Valentino^{3,4}, Sune Toft^{3,4}, Masato Onodera^{2,5}, Rhythm Shimakawa⁵, Daniel Ceverino⁶, Andreas L. Faisst⁷, Anna Gallazzi⁸, Carlos Gómez-Guijarro^{3,4,9}, Mariko Kubo¹, Georgios E. Magdis^{3,4,10}, Charles L. Steinhardt^{3,4}, Mikkel Stockmann^{3,4}, Kiyoto Yabe¹¹, and Johannes Zabl¹²

¹ National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan; masayuki.tanaka@nao.ac.jp

² Department of Astronomical Science, The Graduate University for Advanced Studies, SOKENDAI, Mishima, 411-8540 Japan

³ Cosmic Dawn Center (DAWN), Denmark

⁴ Niels Bohr Institute, University of Copenhagen, Lyngbyvej 2, DK-2100 Copenhagen, Denmark

⁵ Subaru Telescope, National Astronomical Observatory of Japan, 650 N. Aohoku Place, Hilo, HI 96720, USA

⁶ Universidad Autónoma de Madrid, Ciudad Universitaria de Cantoblanco, E-28049 Madrid, Spain

⁷ IPAC, California Institute of Technology, 1200 East California Boulevard, Pasadena, CA 91125, USA

⁸ INAF-Osservatorio Astrofisico di Arcetri, Largo Enrico Fermi 5, I-50125 Firenze, Italy

⁹ Laboratoire AIM-Paris-Saclay, CEA/DSM-CNRS-Universite Paris Diderot, Irfu/Service d'Astrophysique, CEA Saclay, Orme des Merisiers, F-91191 Gif-sur-Yvette, France

¹⁰ DTU Space, National Space Institute, Technical University of Denmark, Elektrovej 327, DK-2800 Kgs. Lyngby, Denmark

¹¹ Kavli Institute for the Physics and Mathematics of the Universe (Kavli IPMU, WPI), University of Tokyo, Chiba 277-8582, Japan

¹² Univ. Lyon, Univ. Lyon1, Ens de Lyon, CNRS, Centre de Recherche Astrophysique de Lyon UMR5574, F-69230 Saint-Genis-Laval, France

Received 2019 September 23; revised 2019 October 15; accepted 2019 October 22; published 2019 November 6

Abstract

We present the first stellar velocity dispersion measurement of a massive quenching galaxy at $z = 4$. The galaxy is first identified as a massive $z \geq 4$ galaxy with suppressed star formation from photometric redshifts based on deep multiband data. A follow-up spectroscopic observation with MOSFIRE on Keck revealed strong multiple absorption features, which are identified as Balmer lines, giving a secure redshift of $z = 4.01$. This is the most distant quiescent galaxy known to date. Thanks to the high S/N of the spectrum, we are able to estimate the stellar velocity dispersion, $\sigma = 268 \pm 59 \text{ km s}^{-1}$, making a significant leap from the previous highest redshift measurement at $z = 2.8$. Interestingly, we find that the velocity dispersion is consistent with that of massive galaxies today, implying no significant evolution in velocity dispersion over the last 12 Gyr. Based on a stringent upper limit on its physical size from deep optical images ($r_{\text{eff}} < 1.3 \text{ kpc}$), we find that its dynamical mass is consistent with the stellar mass inferred from photometry. Furthermore, the galaxy is located on the mass fundamental plane extrapolated from lower redshift galaxies. The observed no strong evolution in σ suggests that the mass in the core of massive galaxies does not evolve significantly, while most of the mass growth occurs in the outskirts of the galaxies, which also increases the size. This picture is consistent with a two-phase formation scenario in which mass and size growth is due to accretion in the outskirts of galaxies via mergers. Our results imply that the first phase may be completed as early as $z \sim 4$.

Unified Astronomy Thesaurus concepts: Galaxy formation (595); Galaxy evolution (594); Elliptical galaxies (456); Galaxy dynamics (591); High-redshift galaxies (734)

1. Introduction

The majority of massive galaxies in the local universe have not been actively forming stars for $\gtrsim 10$ billion years. Their spectroscopic properties are consistent with an intense starburst that occurred in the early universe followed by passive evolution (e.g., Gallazzi et al. 2005; Thomas et al. 2005; Renzini 2006). The recent advent of sensitive near-IR spectrographs has allowed us to reach close to their primary formation epoch. There are a handful of massive quiescent galaxies confirmed out to $z = 3.7$ (Glazebrook et al. 2017; Schreiber et al. 2018). Detailed photometric analyses suggest that they indeed seem to form in a short and intense starburst, followed by rapid quenching (Schreiber et al. 2018). However, the physics of this entire process still remains unclear.

A key parameter for characterizing quiescent galaxies is the stellar velocity dispersion, which is an integrated motion of stars along the line of sight. It exhibits tight correlations with other fundamental properties of galaxies (e.g., the fundamental plane; Djorgovski & Davis 1987). It is also the best predictor of stellar age with no residual correlation between size and age at fixed velocity dispersion (Graves et al. 2009; van der Wel et al. 2009).

The velocity dispersion measurement is observationally challenging due to the demanding S/N of rest-frame optical spectra. It has been measured out to $z = 2.7$ with the help of gravitational lensing effects (Hill et al. 2016). Although quiescent galaxies have been confirmed at higher redshifts, their dynamical properties remain unknown. This is unfortunate because dynamical information may also hold a key to understanding how these galaxies form.

This paper presents the spectroscopic confirmation of a massive quenching galaxy at $z = 4.01$, the most distant galaxy with suppressed star formation rate (SFR) known to date. The paper further presents the measurement of its stellar velocity dispersion, opening a new window to explore dynamical properties of massive galaxies at $z \sim 4$. A companion paper (Valentino et al. 2019) discusses star formation histories and progenitors of this and other galaxies at $z \sim 4$ in detail. We first summarize our observation in Section 2, and then present spectral analyses in Section 3. Measurements of the physical size and discussions of its dynamical properties are given in Sections 4 and 5, respectively. Finally, we discuss implications of our results and conclude the paper in Section 6. We adopt




Figure 1. Left: MOSFIRE K -band spectrum. The top panel shows the 2D spectrum. The A and B nods are in white and the combined spectrum is in black. The bottom panel shows the extracted 1D spectrum. The black and blue spectra are the object and noise spectra binned over approximately the resolution element, respectively. The red spectrum is the best-fit model spectrum from `ppxf`. Right: broadband SED of the object. The red circles and blue squares are the observed and model photometry, respectively. The spectrum is the best-fitting model spectrum from the photometric redshift code (Tanaka 2015).

$H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M = 0.3$, and $\Omega_\Lambda = 0.7$, unless otherwise stated. Magnitudes are in the AB system.

2. Observation

2.1. Target Selection

We select massive quiescent galaxy candidates located at $z \sim 4$ using deep multiwavelength data available in Subaru/*XMM-Newton* Deep Field. We have compiled *uBVRizJHK*[3.6][4.5][5.8][8.0] photometry measured in a consistent manner and applied a photometric redshift code from Tanaka (2015) to infer redshifts as well as SFR and stellar mass. Further details are given in Kubo et al. (2018). We define galaxies with 1σ upper limit of specific SFR below $10^{-9.5} \text{ yr}^{-1}$ as quiescent galaxies. These candidates are typically $K \sim 24$. Among them, there is one outstandingly bright galaxy with $K = 21.9$, which is the subject of the paper. There is no nearby galaxy or galaxy cluster, and this object is not strongly lensed. Figure 1 (right panel) shows the SED of the object. It is a massive galaxy at $z_{\text{phot}} \sim 4$ exhibiting a prominent Balmer break, which is indicative of a recent starburst a few hundred million years ago. The SFR inferred from the fit is low for its stellar mass; $\text{SFR} = 24.0^{+21.7}_{-22.7}$ and $M_* = 1.15^{+0.11}_{-0.10} \times 10^{11} M_\odot$, giving a specific SFR of $\sim 10^{-10} \text{ yr}^{-1}$. The object is about 1 dex below the sequence of star-forming galaxies on an SFR versus M_* diagram and it is likely that this galaxy recently quenched (see Valentino et al. 2019 for further discussions).

2.2. Observation and Data Reduction

We were allocated the first half nights of the 2018 December 20–21 to follow-up the target with Keck/MOSFIRE. The observing conditions were good and the seeing was around

$0''.7$. Each exposure was 180 s long and the classical ABBA nodding was applied along the slit. The total integration time was 7.75 hr. A bright star was put in the mask, which allowed us to keep track of changes in the observing conditions. 7% of the exposures with low fluxes from the bright star were excluded due to poor seeing and/or poor alignment. The data were processed in a standard manner using the MOSFIRE DRP 2018 release. AOV stars were observed as spectrophotometric standards twice per night. The flux calibration vectors derived from those stars were consistent at a few percent level on both nights. The vectors were averaged and the mean flux calibration vector was applied to the extracted 1D spectra.

Figure 1 shows the spectrum. There are four prominent absorption lines, which were identified as $H\gamma$ to $H\zeta$. There is also $H\eta$ on the edge. We confirm the redshift to be $z = 4.012 \pm 0.001$. This proves the accuracy of our photometric redshift estimate ($z_{\text{phot}} = 4.12$). There is no emission line in the spectrum, and 1σ upper limit on (unobscured) SFR from $H\gamma$ is $\lesssim 8 M_\odot \text{ yr}^{-1}$. There is no IR detection either. All this confirms the quiescent nature of the galaxy (see Valentino et al. 2019).

3. Spectral Fit

We fit the observed MOSFIRE spectrum using `ppxf` (Cappellari & Emsellem 2004; Cappellari 2017). We use the simple stellar population models from Vazdekis et al. (2010) for the fit. We exclude all models that are older than the age of the universe at the redshift of the object. We also exclude models with subsolar metallicity as we are focusing on a massive galaxy. Since the rest-frame spectral resolution of the MOSFIRE spectrum is slightly higher than that of the library, we apply Gaussian smoothing to match the resolution. We use




Figure 2. Size from HSC i -band plotted against size from *HST* F160W. The plotted objects have i -band magnitudes similar to that of the $z = 4$ object. The solid line shows $r_{\text{eff, HSC}} = r_{\text{eff, HST}}$. The left and right panels show objects at $z < 1.5$ and $z > 1.5$, respectively. The systematic offset and scatter around it are shown in the figures.

an additive correction function of order 1 (i.e., linear) and no multiplicative correction. Our results are not sensitive to the choices here.

The best-fit spectrum is shown in red in Figure 1. All observed Balmer absorption lines are fit very well, and the overall fit has $\chi^2_\nu = 1.2$. In addition, thanks to the high S/N of the spectrum ($S/N \sim 5$ per resolution element), we measure a velocity dispersion of $268 \pm 59 \text{ km s}^{-1}$. The uncertainty here is based on a Monte-Carlo simulation; we perturb the observed spectrum using the noise spectrum and repeat the fits. The quoted uncertainty is the 68th percentile of the distribution from the Monte-Carlo fits. If we use the Indo-US stellar spectral library (Valdes et al. 2004) instead of Vazdekis so that we do not need to smooth the MOSFIRE spectrum, we obtain a consistent stellar velocity dispersion ($\sigma = 252 \pm 77 \text{ km s}^{-1}$). We also have confirmed that emission line in-filling does not affect our measurement either; we repeat the fits by excluding $H\gamma$ and $H\delta$ and obtain a fully consistent measurement, $\sigma = 277 \pm 58 \text{ km s}^{-1}$.

4. Physical Size

In order to fully utilize the stellar velocity dispersion, we are interested in structural properties of the $z = 4$ galaxy, in particular its size. The typical rest-frame optical size of $z \sim 4$ galaxies is very small ($r_{\text{eff}} \sim 0.5 \text{ kpc}$; Kubo et al. 2018), and adaptive optics assisted observations or high spatial resolution imaging with the *Hubble Space Telescope* (*HST*) are an ideal way to measure the size of the $z = 4$ galaxy. Unfortunately, there is no existing *HST* data for the galaxy and Kubo et al. (2018) did not observe the galaxy with AO.

However, a useful upper limit on the size can still be obtained from the ground-based data. We use deep optical data from Hyper Suprime-Cam Subaru Strategic Program (HSC-SSP; Aihara et al. 2018, 2019) and demonstrate how well we

can reproduce the sizes measured from *HST*. The i -band image from HSC-SSP processed with the pipeline designed for LSST (Jurić et al. 2017; Bosch et al. 2018; Ivezić et al. 2019) is used here to measure sizes due to superior seeing than the other bands. We retrieve objects with similar i -band magnitudes to the $z = 4$ galaxy ($i \sim 24.3$) in UltraDeep-SXDS, which is the same field as UDS, in which the $z = 4$ galaxy is located, and run `galfit` (Peng et al. 2002, 2010) to measure their effective radii adopting the Sérsic profile. We use the variance image to generate a sigma map and use the “coadd PSF” (Bosch et al. 2018) as an input PSF image. The PSF has $\text{FWHM} = 0''.65$. We perform a Monte-Carlo run by slightly perturbing the initial centroid, position angle, effective radius and brightness. We fix the Sérsic index to a randomly drawn value between 0.5 and 4 in each run. We use the fit with the smallest χ^2 as the best estimate and $\Delta\chi^2 < 1$ as the 68th percentile interval.

Figure 2 compares the size measurements between HSC and *HST*. The *HST* sizes are taken from van der Wel et al. (2014) and are measured in the WFC3/F160W filter. We split the sample into low and high-redshift ranges using the photo- z described in Section 2.1. These two panels give a quantitative estimate of the rest-frame wavelength dependence of the size measurements. Interestingly, we observe equally good correlation between the two measurements down to $\sim 0''.1$ in both plots. If we define outliers as $|r_{\text{eff, HSC}} - r_{\text{eff, HST}}| > 0''.2$, the outlier rate is about 4% in both plots. This good correlation is likely due to the depth of the HSC data (the 5σ depth is $\sim 27 \text{ mag}$) and also to the good PSF model. Other studies also have shown that sizes can be measured to $\sim 0.2 \text{ FWHM}$ (van der Wel et al. 2014) for high S/N and adequate spatial sampling. It is encouraging that there seems to be no major evidence for large morphological k -corrections in the size measurements. We note that van der Wel et al. (2014) also find that the wavelength dependence of size is not very strong for quiescent galaxies.




Figure 3. Panels show the *i*-band image from HSC, the best-fit *galfit* model, and residual, from left to right. The flux scales are the same in all the panels.




Figure 4. Left: size plotted against stellar mass. The $z = 4$ galaxy is shown in red. The black points are quiescent galaxies at $z > 1.5$ drawn from the literature; compilation by van de Sande et al. (2013), which includes van Dokkum et al. (2009), Onodera et al. (2012), Toft et al. (2012), Bezanson et al. (2013), and newer data from Belli et al. (2017) and Stockmann et al. (2019). Their mean redshift is around 2. The shades show data from the LEGA-C survey, which covers $0.6 \lesssim z \lesssim 1.0$. The dashed line shows the local relation from Shen et al. (2003) based on data from the Sloan Digital Sky Survey (SDSS; York et al. 2000; Strauss et al. 2002). The solid and dotted lines in the bottom right corner show evolutionary tracks with $r_{\text{eff}} \propto M_*$ and $r_{\text{eff}} \propto M_*^2$, respectively. They represent the major and minor merger tracks. Right: stellar velocity dispersion plotted against stellar mass. The meaning of the symbols are the same as in the left figure. The blue dashed curve is a fit to SDSS galaxies (Zahid et al. 2016).

Adopting this size measurement procedure for the $z = 4$ galaxy, we find the best size estimate of $0.11 \pm 0.03''$ (0.76 ± 0.20 kpc). This size is fully consistent with the typical size of $z \sim 4$ massive quiescent galaxies from Kubo et al. (2018). Figure 3 shows the object, best-fit model, and residual images, demonstrating an excellent fit. Figures 2 and 3 suggest that our estimate here is likely reasonable, but to be fully conservative, we primarily use the upper limit on the size in what follows; $0.76 + 0.20$ (random) $+0.35$ (systematic corresponding to $0.05''$ observed scatter) $= 1.3$ kpc.

The left panel of Figure 4 shows r_{eff} against stellar mass. In addition to the $z = 4$ galaxy, the figure also includes lower redshift galaxies for comparison; quiescent galaxies at $z \sim 2$ from a compilation of literature and $0.6 \lesssim z \lesssim 1.0$ galaxies from the LEGA-C survey (van der Wel et al. 2016; Straatman et al. 2018) cross-matched with the size measurements from van der Wel et al. (2014). Quiescent galaxies among the LEGA-C galaxies are selected using the multiband classification from Laigle et al. (2016). The $z = 4.01$ galaxy is compact

for its stellar mass, and it is physically smaller than $z \sim 2$ and other lower redshift galaxies. This clear redshift trend is consistent with Kubo et al. (2018).

In contrast, the stellar velocity dispersion of the $z = 4$ galaxy plotted in the right panel is largely consistent with those of the lower redshift galaxies. There is a possible hint that low redshift galaxies from LEGA-C and SDSS have a slightly lower velocity dispersion. However, the difference, if any, is fairly modest. This indicates that the stellar velocity dispersion has not significantly evolved over the last 12 Gyr. This is an intriguing result because the size and mass are known to evolve significantly over this time period.

5. Dynamical Analysis

Using the size from the previous section, we estimate the galaxy’s dynamical mass as

$$M_{\text{dyn}} = \frac{\beta(n) \sigma^2 r_{\text{eff}}}{G}, \quad (1)$$




Figure 5. Left: stellar mass plotted against dynamical mass. The dotted line shows $M_{*} = M_{\text{dyn}}$. The black points are the lower redshift objects ($z \sim 2$) as in Figure 4. The shades are quiescent galaxies from LEGA-C, and the dashed curve is the running median of the distribution of quiescent galaxies with $\text{sSFR} < 10^{-10} M_{\odot} \text{ yr}^{-1}$ drawn from SDSS DR15 (Aguado et al. 2019). SFR and stellar mass are from Granada FSPS fits (Ahn et al. 2014). The red point is the $z = 4$ object. The pink points are the expected evolutionary track assuming constant velocity dispersion (see the text for details). Right: mass fundamental plane. The dotted line is a fit from Bezanson et al. (2013). The other symbols are the same as those in the left panel.

where β is $\beta(n) = 8.87 - 0.831n + 0.024n^2$ (Cappellari et al. 2006). As discussed in the same paper, a constant $\beta = 5.0$ also works well. Given the unconstrained n for our object, we adopt $\beta = 5$ here, which yields $M_{\text{dyn}} = 1.3 \times 10^{11} M_{\odot}$ with an upper limit of $2.1 \times 10^{11} M_{\odot}$.

Figure 5 (left panel) compares the stellar mass with the dynamical mass estimated here. The stellar mass of the $z = 4$ object is fully consistent with the dynamical mass. We use the Chabrier IMF here (Chabrier 2003). This consistency is encouraging because this confirms the accuracy of our photometry-based estimate of stellar mass, which then implies that assumptions employed such as the initial mass function are reasonable. The $z = 4$ galaxy is expected to increase its size and stellar mass with time through mergers. Marchesini et al. (2014) estimated the likely stellar mass growth from abundance matching, and Kubo et al. (2018) followed the size growth of these galaxies on the stellar mass evolutionary track from Marchesini et al. (2014). From these results, we can make a prediction for how the galaxy will evolve in the M_{*} versus M_{dyn} plane. If we assume that the velocity dispersion does not change with time, which is the assumption also taken by, e.g., Belli et al. (2017), the expected location of the descendant of the $z = 4$ galaxy is shown as the pink points (Figure 5). The expected evolutionary track shows an increase in the dynamical mass with a smaller increase in the stellar mass. If minor mergers entirely drive the evolution and velocity dispersion does not change over time, we expect $M_{*} \propto r_{\text{eff}}^{0.5} \propto M_{\text{dyn}}^{0.5}$. As the size growth observed by Kubo et al. (2018) is close to the maximum growth rate expected from minor mergers, the evolutionary track is indeed close to $M_{*} \propto M_{\text{dyn}}^{0.5}$.

In the local universe, early-type galaxies are known to show a tight relationship between effective radius, mean intensity,

and velocity dispersion (i.e., fundamental plane; Djorgovski & Davis 1987). The evolution of the fundamental plane is mostly due to the evolving stellar mass to luminosity ratio due to stellar aging. The mass fundamental plane, which replaces luminosity with stellar mass, has been suggested in the literature to largely remove that effect (Bezanson et al. 2013). Figure 5 (right panel) shows the mass fundamental plane.

The $z = 4$ galaxy is located at the bottom left part of the distribution of the lower redshift galaxies as expected from its small size and high stellar density. The evolutionary track with constant velocity dispersion is along the location of the lower redshift objects and the galaxy will likely be among the most massive galaxies at each epoch. This in turn implies that the velocity dispersion does not evolve significantly. If it had changed by a factor of, e.g., 2 by $z = 1$, the expected location of the descendant galaxy would be inconsistent with the observed $z \sim 1$ galaxies.

We also briefly discuss the classical fundamental plane. The interpretation of its evolution is not straightforward, but assuming that the evolution in the fundamental plane is entirely due to the luminosity evolution for simplicity, we find that the $z = 4$ galaxy is $1.70_{-0.46}^{+\infty}$ dex off in $\log I_{\text{eff}}$ in the rest-frame g -band from the local relation (recall that we have only an upper limit on size). A simple stellar population model from Bruzual & Charlot (2003) predict a luminosity evolution of a factor of ~ 80 , or equivalently ~ 1.9 dex, adopting the luminosity-weighted age of ~ 0.2 Gyr (Valentino et al. 2019) for the galaxy. The fundamental plane evolution therefore is consistent with the pure luminosity evolution, although the uncertainty here is rather large and we do not discuss further at this point.

6. Summary and Discussion

We have presented spectroscopic confirmation of a massive galaxy being quenched at $z = 4.01$, which is the most distant example known to date. Thanks to the high S/N of the spectrum, we are able to measure its stellar velocity dispersion, $268 \pm 59 \text{ km s}^{-1}$. The size estimate based on the deep optical data is $0.76 \pm 0.20 \text{ kpc}$ with a stringent upper limit of $<1.3 \text{ kpc}$, which is consistent with the typical size of massive quiescent galaxies from Kubo et al. (2018). Combining the velocity dispersion and size, we find that the dynamical mass is consistent with the stellar mass inferred from photometry. Also, the galaxy is on the mass fundamental plane and the expected evolutionary path of the galaxy is consistent with the massive quiescent galaxies at lower redshifts.

The most striking finding of this work is that the stellar velocity dispersion of the massive galaxy at $z = 4.01$ is consistent with that of massive galaxies at lower redshifts. This is in contrast to the very large velocity dispersion of a $z = 2.2$ galaxy reported by van Dokkum et al. (2009), $510^{+165}_{-95} \text{ km s}^{-1}$. The $z = 4$ galaxy is expected to increase its mass by a factor of $2.1^{+2.2}_{-1.0}$ (Marchesini et al. 2014) and size by a factor of $10.8^{+5.3}_{-3.6}$ (Kubo et al. 2018) by $z = 1$. However, the stellar velocity dispersion evolution is significantly weaker; Figure 4 (right) shows that even the most massive galaxies at lower redshifts have only slightly larger velocity dispersion. This finding indicates that the size and mass evolution does not significantly increase the total mass contained in the core of these massive galaxies. This has significant implications for how galaxies increase their size and mass; galaxies do not increase mass equally at all radii, and instead most of the mass growth occurs in the outer parts perhaps through minor mergers, which also increases the effective radius. This is fully consistent with the two-phase formation scenario (Naab et al. 2007; Oser et al. 2010). In fact, simulations show only a mild evolution in velocity dispersion (Oser et al. 2012), in agreement with our observation here. An interesting implication here is that the first of the two phases, namely the formation of the dense core, may be completed as early as $z \sim 4$.

It is, however, not clear whether the $z = 4$ galaxy is a dispersion-dominated system. Recent work finds that massive galaxies at $z \sim 2$ exhibit significant rotational motion (Toft et al. 2017; Newman et al. 2018b). Simulations predict a relatively small V/σ if massive galaxies at high redshifts form in starbursts triggered by dissipative mergers (Wuyts et al. 2010). The reported large rotational motion favors formation through disk instabilities. It is possible that the $z = 4$ galaxy also rotates, potentially complicating the interpretation above. But, given the absence of sufficient spatial resolution to resolve any rotational motion from the MOSFIRE data nor even an ellipticity measurement from the imaging, we have to await further observations to reveal V/σ of the $z = 4$ galaxy. The large rotational motion, if present, has to be damped at lower redshifts because the majority of the most massive galaxies in the local universe are slow rotators (e.g., Veale et al. 2017). Mergers may be able to do the job (Lagos et al. 2018).

Prior to this work, the most distant velocity dispersion measurements were made at $z = 2.6\text{--}2.8$ with the help of a gravitational lensing effect (Hill et al. 2016; Newman et al. 2018a). This work makes a major leap in redshift without lensing and demonstrates that the current facilities have the ability to confirm redshifts and measure the stellar velocity

dispersion of the brightest galaxies at $z > 4$. Massive galaxies at such redshifts are rare, but ongoing/upcoming massive imaging surveys will be able to construct a significantly larger sample, which can then be followed up spectroscopically to further extend the work presented here.

This work is supported by JSPS KAKENHI grant Nos. JP23740144 and JP15K17617. F.V. and G.E.M. acknowledge the Villum Fonden research grant 13160 ‘‘Gas to stars, stars to dust: tracing star formation across cosmic time.’’ F.V. acknowledges the Carlsberg Fonden research grant CF18-0388 ‘‘Galaxies: Rise And Death.’’ The Cosmic Dawn Center (DAWN) is funded by the Danish National Research Foundation under grant No. 140. S.T. and G.E.M. acknowledge support from the European Research Council (ERC) Consolidator Grant funding scheme (project ConText, grant No. 648179). M.O. acknowledges support by KAKENHI JP17K14257. K.Y. acknowledges support by JSPS KAKENHI grant No. JP18K13578. We thank the anonymous referee for a useful report, which helped improve the paper.











The HSC collaboration includes the astronomical communities of Japan and Taiwan, and Princeton University. The HSC instrumentation and software were developed by NAOJ, Kavli IPMU, the University of Tokyo, KEK, ASIAA, and Princeton University. Funding was contributed by the FIRST program from Japanese Cabinet Office, the Ministry of Education, Culture, Sports, Science and Technology, the Japan Society for the Promotion of Science, Japan Science and Technology Agency, the Toray Science Foundation, NAOJ, Kavli IPMU, KEK, ASIAA, and Princeton University. This paper makes use of software developed for LSST. We thank the LSST Project for making their code available as free software at <http://dm.lsst.org>. This paper is based in part on data collected at the Subaru Telescope and retrieved from the HSC data archive system, which is operated by Subaru Telescope and ADC at NAOJ. Data analysis was in part carried out with the cooperation of CfCA and NAOJ.

Funding for the SDSS and SDSS-II has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, the U.S. Department of Energy, the National Aeronautics and Space Administration, the Japanese Monbukagakusho, the Max Planck Society, and the Higher Education Funding Council for England. The SDSS website is <http://www.sdss.org/>.

The SDSS is managed by the Astrophysical Research Consortium for the Participating Institutions. The Participating Institutions are the American Museum of Natural History, Astrophysical Institute Potsdam, University of Basel, University of Cambridge, Case Western Reserve University, University of Chicago, Drexel University, Fermilab, the Institute for Advanced Study, the Japan Participation Group, Johns Hopkins University, the Joint Institute for Nuclear Astrophysics, the Kavli Institute for Particle Astrophysics and Cosmology, the Korean Scientist Group, the Chinese Academy of Sciences (LAMOST), Los Alamos National Laboratory, the Max-Planck-Institute for Astronomy (MPIA), the Max-Planck-Institute for Astrophysics (MPA), New Mexico State University, Ohio State University, University of Pittsburgh, University of Portsmouth, Princeton University, the United States Naval Observatory, and the University of Washington.

Facility: Keck(MOSFIRE).

ORCID iDs

Masayuki Tanaka  <https://orcid.org/0000-0002-5011-5178>
 Francesco Valentino  <https://orcid.org/0000-0001-6477-4011>
 Masato Onodera  <https://orcid.org/0000-0003-3228-7264>
 Rhythm Shimakawa  <https://orcid.org/0000-0003-4442-2750>
 Andreas L. Faisst  <https://orcid.org/0000-0002-9382-9832>
 Anna Gallazzi  <https://orcid.org/0000-0002-9656-1800>
 Mariko Kubo  <https://orcid.org/0000-0002-7598-5292>
 Charles L. Steinhardt  <https://orcid.org/0000-0003-3780-6801>
 Mikkel Stockmann  <https://orcid.org/0000-0001-5983-6273>
 Kiyoto Yabe  <https://orcid.org/0000-0001-6229-4858>

References

- Aguado, D. S., Ahumada, R., Almeida, A., et al. 2019, *ApJS*, 240, 23
 Ahn, C. P., Alexandroff, R., Allende Prieto, C., et al. 2014, *ApJS*, 211, 17
 Aihara, H., AIsayyad, Y., Ando, M., et al. 2019, arXiv:1905.12221
 Aihara, H., Arimoto, N., Armstrong, R., et al. 2018, *PASJ*, 70, S4
 Belli, S., Newman, A. B., & Ellis, R. S. 2017, *ApJ*, 834, 18
 Bezanson, R., van Dokkum, P., van de Sande, J., et al. 2013, *ApJL*, 764, L8
 Bezanson, R., van Dokkum, P. G., van de Sande, J., et al. 2013, *ApJL*, 779, L21
 Bosch, J., Armstrong, R., Bickerton, S., et al. 2018, *PASJ*, 70, S5
 Bruzual, G., & Charlot, S. 2003, *MNRAS*, 344, 1000
 Cappellari, M. 2017, *MNRAS*, 466, 798
 Cappellari, M., Bacon, R., Bureau, M., et al. 2006, *MNRAS*, 366, 1126
 Cappellari, M., & Emsellem, E. 2004, *PASP*, 116, 138
 Chabrier, G. 2003, *PASP*, 115, 763
 Djorgovski, S., & Davis, M. 1987, *ApJ*, 313, 59
 Gallazzi, A., Charlot, S., Brinchmann, J., et al. 2005, *MNRAS*, 362, 41
 Glazebrook, K., Schreiber, C., Labbé, I., et al. 2017, *Natur*, 544, 71
 Graves, G. J., Faber, S. M., & Schiavon, R. P. 2009, *ApJ*, 698, 1590
 Hill, A. R., Muzzin, A., Franx, M., et al. 2016, *ApJ*, 819, 74
 Ivezić, Ž., Kahn, S. M., Tyson, J. A., et al. 2019, *ApJ*, 873, 111
 Jurić, M., Kantor, J., Lim, K.-T., et al. 2017, in ASP Conf. Ser. 512, Astronomical Data Analysis Software and Systems XXV, ed. N. P. F. Lorente, K. Shorridge, R. Wayth et al. (San Francisco, CA: ASP), 279
 Kubo, M., Tanaka, M., Yabe, K., et al. 2018, *ApJ*, 867, 1
 Lagos, C. del P., Stevens, A. R. H., Bower, R. G., et al. 2018, *MNRAS*, 473, 4956
 Laigle, C., McCracken, H. J., Ilbert, O., et al. 2016, *ApJS*, 224, 24
 Marchesini, D., Muzzin, A., Stefanon, M., et al. 2014, *ApJ*, 794, 65
 Naab, T., Johansson, P. H., Ostriker, J. P., et al. 2007, *ApJ*, 658, 710
 Newman, A. B., Belli, S., Ellis, R. S., et al. 2018a, *ApJ*, 862, 125
 Newman, A. B., Belli, S., Ellis, R. S., et al. 2018b, *ApJ*, 862, 126
 Onodera, M., Renzini, A., Carollo, M., et al. 2012, *ApJ*, 755, 26
 Oser, L., Naab, T., Ostriker, J. P., et al. 2012, *ApJ*, 744, 63
 Oser, L., Ostriker, J. P., Naab, T., et al. 2010, *ApJ*, 725, 2312
 Peng, C. Y., Ho, L. C., Impey, C. D., et al. 2002, *AJ*, 124, 266
 Peng, C. Y., Ho, L. C., Impey, C. D., et al. 2010, *AJ*, 139, 2097
 Renzini, A. 2006, *ARA&A*, 44, 141
 Schreiber, C., Glazebrook, K., Nanayakkara, T., et al. 2018, *A&A*, 618, A85
 Shen, S., Mo, H. J., White, S. D. M., et al. 2003, *MNRAS*, 343, 978
 Stockmann, M., Toft, S., Gallazzi, A., et al. 2019, *ApJ*, submitted
 Straatman, C. M. S., van der Wel, A., Bezanson, R., et al. 2018, *ApJS*, 239, 27
 Strauss, M. A., Weinberg, D. H., Lupton, R. H., et al. 2002, *AJ*, 124, 1810
 Tanaka, M. 2015, *ApJ*, 801, 20
 Thomas, D., Maraston, C., Bender, R., et al. 2005, *ApJ*, 621, 673
 Toft, S., Gallazzi, A., Zirm, A., et al. 2012, *ApJ*, 754, 3
 Toft, S., Zabl, J., Richard, J., et al. 2017, *Natur*, 546, 510
 Valdes, F., Gupta, R., Rose, J. A., et al. 2004, *ApJS*, 152, 251
 Valentino, F., Tanaka, M., Davidzon, I., et al. 2019, arXiv:1909.10540
 van de Sande, J., Kriek, M., Franx, M., et al. 2013, *ApJ*, 771, 85
 van der Wel, A., Bell, E. F., van den Bosch, F. C., et al. 2009, *ApJ*, 698, 1232
 van der Wel, A., Franx, M., van Dokkum, P. G., et al. 2014, *ApJ*, 788, 28
 van der Wel, A., Noeske, K., Bezanson, R., et al. 2016, *ApJS*, 223, 29
 van Dokkum, P. G., Kriek, M., & Franx, M. 2009, *Natur*, 460, 717
 Vazdekis, A., Sánchez-Blázquez, P., Falcón-Barroso, J., et al. 2010, *MNRAS*, 404, 1639
 Veale, M., Ma, C.-P., Greene, J. E., et al. 2017, *MNRAS*, 471, 1428
 Wuyts, S., Cox, T. J., Hayward, C. C., et al. 2010, *ApJ*, 722, 1666
 York, D. G., Adelman, J., Anderson, J. E., et al. 2000, *AJ*, 120, 1579
 Zahid, H. J., Geller, M. J., Fabricant, D. G., et al. 2016, *ApJ*, 832, 203



Erratum: “Stellar Velocity Dispersion of a Massive Quenching Galaxy at $z = 4.01$ ” (2019, *ApJL*, 885, L34)

Masayuki Tanaka^{1,2}, Francesco Valentino^{3,4}, Sune Toft^{3,4}, Masato Onodera^{2,5}, Rhythm Shimakawa⁵,
Daniel Ceverino⁶, Andreas L. Faisst⁷, Anna Gallazzi⁸, Carlos Gómez-Guijarro^{3,4,9}, Mariko Kubo¹,
Georgios E. Magdis^{3,4,10}, Charles L. Steinhardt^{3,4}, Mikkel Stockmann^{3,4}, Kiyoto Yabe¹¹, and Johannes Zabl¹²

¹ National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan

² Department of Astronomical Science, The Graduate University for Advanced Studies, SOKENDAI, Mishima, 411-8540 Japan

³ Cosmic Dawn Center (DAWN), Denmark

⁴ Niels Bohr Institute, University of Copenhagen, Lyngbyvej 2, DK-2100 Copenhagen, Denmark

⁵ Subaru Telescope, National Astronomical Observatory of Japan, 650 N Aohoku Place, Hilo, HI 96720, USA

⁶ Universidad Autónoma de Madrid, Ciudad Universitaria de Cantoblanco, E-28049 Madrid, Spain

⁷ IPAC, California Institute of Technology, 1200 East California Boulevard, Pasadena, CA 91125, USA

⁸ INAF-Osservatorio Astrofisico di Arcetri, Largo Enrico Fermi 5, I-50125 Firenze, Italy

⁹ Laboratoire AIM-Paris-Saclay, CEA/DSM-CNRS-Universite Paris Diderot, Irfu/Service d’Astrophysique, CEA Saclay, Orme des Merisiers, F-91191 Gif-sur-Yvette, France

¹⁰ DTU Space, National Space Institute, Technical University of Denmark, Elektrovej 327, DK-2800 Kgs. Lyngby, Denmark

¹¹ Kavli Institute for the Physics and Mathematics of the Universe (Kavli IPMU, WPI), University of Tokyo, Chiba 277-8582, Japan

¹² Univ. Lyon, Univ. Lyon1, Ens de Lyon, CNRS, Centre de Recherche Astrophysique de Lyon UMR5574, F-69230 Saint-Genis-Laval, France

Received 2020 April 14; published 2020 May 5

1. Introduction

We have reported on the stellar velocity dispersion measurement of a massive galaxy at $z = 4.01$ in the published article (Tanaka et al. 2019). The dynamical mass in the paper was quoted erroneously about a factor of 2 higher. The dynamical mass quoted in Section 5 should have been $M_{\text{dyn}} = 6.3 \times 10^{10} M_{\odot}$ with an upper limit of $1.2 \times 10^{11} M_{\odot}$. The same error was also in the dynamical mass estimates for the LEGA-C and SDSS samples. An updated version of Figure 5 (left) can be found below. This error does not change the conclusions of the paper at all.







Figure 5. Left: stellar mass plotted against dynamical mass. The meaning of the symbols are the same as in the published article.

ORCID iDs

Masayuki Tanaka  <https://orcid.org/0000-0002-5011-5178>
Francesco Valentino  <https://orcid.org/0000-0001-6477-4011>

Sune Toft  <https://orcid.org/0000-0003-3631-7176>

Masato Onodera  <https://orcid.org/0000-0003-3228-7264>

Rhythm Shimakawa  <https://orcid.org/0000-0003-4442-2750>

Daniel Ceverino  <https://orcid.org/0000-0002-8680-248X>

Andreas L. Faisst  <https://orcid.org/0000-0002-9382-9832>


Anna Gallazzi  <https://orcid.org/0000-0002-9656-1800>

Mariko Kubo  <https://orcid.org/0000-0002-7598-5292>

Georgios E. Magdis  <https://orcid.org/0000-0002-4872-2294>

Charles L. Steinhardt  <https://orcid.org/0000-0003-3780-6801>

Mikkel Stockmann  <https://orcid.org/0000-0001-5983-6273>

Kiyoto Yabe  <https://orcid.org/0000-0001-6229-4858>

Reference

Tanaka, M., Valentino, F., Toft, S., et al. 2019, *ApJL*, 885, L34