



Myocarditis and Autoimmune Disease Development Following Bariatric Surgery: A Possible Nutritional Deficiency Link

**Zainab Al Bahrani^{a,b}, Abdulaziz Sarawi^{a,b,c},
Hani Abu Shanab^{a,b}, Ayman F. Soliman^{a,b,c*}
and Ali A. Al Qarni^{a,b,c}**

^a King Abdulaziz Hospital, MNGHA, Al Ahsa, Saudi Arabia.

^b King Abdullah International Medical Research Center (KAIMRC), Al Ahsa, Saudi Arabia.

^c King Saud bin Abdulaziz University for Health Sciences, COAMS-A, Al Ahsa, Saudi Arabia.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJMAH/2023/v21i9856

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/101774>

Case Report

Received: 12/04/2023

Accepted: 15/06/2023

Published: 20/06/2023

ABSTRACT

Post bariatric surgery induced myocarditis in setting of autoimmune disease is rarely encountered in clinical practice. However, till date only few data of bariatric surgery induced myocarditis in setting of connective tissue disease (CTD) have been available. The following report illustrates the case of a 39-year-old female suffering from myositis who exhibited a nutritional deficiency myocarditis following biliopancreatic diversion surgery (BPD). The mechanism of myocarditis in patients suffering from CTD is incompletely understood. In this case we are going to try to found a relation between bariatric surgery, myocarditis and connective tissue disease and urges cautious

*Corresponding author: E-mail: Solimana@ngha.med.sa, SolimanA@mngha.med.sa;

action before surgery performance in the setting of suspected nutritional deficiencies and in connective tissue disease (CTD) as injudicious act might increase the risk of deleterious myocarditis and increase the mortality.

Keywords: Myocarditis; autoimmune disease; biliopancreatic diversion; nutritional deficiency.

1. INTRODUCTION

Acute myocarditis is defined as an acute inflammatory disease of the myocardium, caused by a variety of infectious (e.g., viral, bacterial) and noninfectious conditions (including cardio toxins, hypersensitivity reactions, systemic disorders, and radiation) [1]. Myocarditis is an inflammatory disease of the heart with major public health impact. Thorough understanding of its immune pathogenesis is crucial for accurate diagnosis and effective treatment. Myocarditis refers to an inflammatory process in the heart that can be initiated by various factors. The most common cause of myocarditis is viral infection [2].

Bariatric surgery is an effective option for weight loss in obese patients. It is an operation in which part of the stomach is resected, creating a smaller stomach to which the distal part of the small intestine is then connected, bypassing the duodenum and jejunum. It represents a mal absorptive procedure [3].

There are few cases report an autoimmune diseases post bariatric surgery however none of them prove a causal relationship between both of them [4]. The autoimmune diseases can cause myocarditis, because the immune system may target any organ in the body, including the heart, producing inflammation. Although a nutritional cardiomyopathy was observed [5], myocarditis and post-bariatric surgery were not linked.

2. CASE REPORT

A 39-year-old female patient known case of diabetes mellitus, hypothyroidism, post biliopancreatic diversion (BPD) with duodenal switch, Laparoscopic cholecystectomy 2008, Post Abdominal liposuction with pre umbilical hernia repair 2019, chronic Multiple nutritional deficiency and depression. Presented to emergency department with complaints of; shortness of breath at rest with orthopnoea& paroxysmal nocturnal dyspnoea for 14 days, along with some chest discomfort & occasional palpitations and lower limbs oedema. She also was complaining of lower limb weakness. On physical examination her chest showed bilateral

fine basal crepitation. She was tachycardia with low blood pressure, and generalized weakness. Blood test showed elevated cardiac enzymes with increasing CK which is not correlating to the CKMB, elevated BNP, hypomagnesemia and hypocalcemia. The significant laboratory findings included, fluctuating Zinc level and selenium was low normal.

She was admitted to the cardiac care unit as a case of acute heart failure, and managed accordingly. She was received IV replacement of multi- vitamins and micronutrients. Cardiac work up showed RBBB with wide spread T wave inversion in the ECG. The Echocardiogram was found to be dilated cardiomyopathy with sever biventricular dysfunction and low ejection fraction (25-30%) with severe global impairment of systolic function with grade III diastolic dysfunction, left atrium moderately dilated, Moderate MR, Moderate TR, RVSP 65, Trivial pericardial effusion. Cardiovascular MRI suggestive of sub-acute myocarditis. The end myocardial tissue biopsy showed; Myocyte hypertrophy with focal interstitial fibrosis. Focal myocyte vascular changes consistent with myocyte injury. Negative for myocarditis, vasculitis and granulomas. Negative for amyloid and iron. The histological features are nonspecific and can be seen in a wide range of pathological conditions including cardiomyopathies and post myocarditis.

The patient was referred to rheumatology team and neurology team for high CK and skeletal myopathy. Nerve conduction study done, showed sever myopathy change. Rheumatology investigation showed positive ANA, anti-SSB, Ro2-52 Ab positive Femur/Thigh MRI was done showed Signs suggestive of interval regression of polymyositis in both thighs. With diagnosis of SLE / Sjogren disease with inflammatory myositis

The Rheumatology team treat the patient by prednisone. The patient heart failure condition was improved clinically.

Her condition was slowly improved. She was discharged home after one month of multidisciplinary treatment team.

3 weeks later patient was admitted with worsening heart failure patient stabilized and transfer to advance heart failure centre for end myocardial tissue biopsy and for further management. At the advance heart failure centre she developed cardiogenic shock requiring intubation and mechanical ventilation. Patient treated in Intensive care unit accordingly. Rheumatology team started Azathioprine, IVIG, and cyclosporine with steroids.

Patient condition was gradually improved and discharged after four month. During admission cardiac transplant team involved for work up however she was not eligible for heart transplant due to the presence of anti-myocyte AB.

Four months later she presented again to emergency department with body ache, pain, dyspnea, fever and cough for 4 days, upon presentation she was found hypotensive admitted under internal medicine in stepdown but later on the same day her condition worsened when she required intensive care admission. Upon admission to intensive care she became more hypotensive, requiring high doses of vasopressors (norepinephrine and vasopressin) with signs of low perfusion so she was intubated and mechanically ventilated, and connected to continuous noninvasive cardiac monitoring which revealed significant drop of cardiac output (EF < 10%) Patient showed worsening course as she developed refractory shock status on mega doses of vasopressors also she developed multi organ failure and disseminated intravascular coagulopathy. Next day she grew gram negative bacilli in the blood (she was started already on wide spectrum antibiotics).

Patient developed refractory septic / cardiogenic shock requiring maximum dose of inotropes and vasopressors, unfortunately her condition not improved and she passed away.

3. DISCUSSION

This patient is a known case of post biliopancreatic diversion, which is a malabsorptive procedure for obesity management. This procedure's well-known side effects include nutritional deficiencies of micronutrients, trace elements, and albumin. A few Similar cases have been reported [6,7]. The etiology includes autoimmune and inflammatory conditions and exposure to toxic agents. However, complications such as heart failure or arrhythmias may worsen, especially in the early

postoperative period [8]. There is emerging evidence for immunosuppressive therapy in some instances that are considered related in this respect. Despite our diligent search, we found few cases of novel immunosuppressant drug-induced myocarditis in the setting of autoimmune disease in the published literature resulting from an overly aggressive immune system [9]. The nutritional deficiency is associated with an immunocompromised state which contributes to cardiac dysfunction as myocarditis; however, 11% reduction in coronary events was observed in subjects taking selenium replacement [10].

In this case the reversal of the primary procedure (BPD) would improve the patient current nutritional deficiencies as well as the maximum absorptive benefit of the immunosuppressive medication; however, No solid evidence is there that this patient cardiac disease is related to micronutrient deficiencies as consequences of the malabsorption effects of BPD and no guarantee for improvement of her cardiac conditions after reversal of BPD. Therefore, the timing of the reversal procedure should be set after reaching maximum optimization of her current conditions. Moreover, 33% of myocarditis patients required mechanical circulatory support (MCS) or cardiac transplantation [11]. However, this is not the case in this patient because there is HLA incompatibility for heart transplantation with the risk of rejection and the fear of malabsorption to medications. Furthermore, the autoimmune nature of systemic disease would have a high chance of recurrence after transplant and failure of the procedure.

Despite immunological dysregulation that made the patient more vulnerable to opportunistic infections, documented cases typically result in myocarditis, according to the literature [12]. 50-80% of myocarditis cases has no cause is ever found, and infectious is reported in 1-5% of patients as compared to nutritional deficiency which is caused by 17-35% [13]. However, Virus-induced abnormal immunomodulation in cardiac patients is believed to be a key factor for the development of carditis during concomitant immunosuppressive therapy [14,15]. Infection causes the innate immune system in the heart to become more active in viral myocarditis. Chronic inflammation that is exacerbated by excessive or ongoing activation can cause myocardial remodeling and damage, which ultimately results in cardiac dysfunction. This, in this scenario, could emerge the myocarditis. Furthermore,

Carsten et al. describe that the persistence of infection and development of dilated cardiomyopathy have been linked to the patient's genetic background [15]. From this perspective, this might have played a role in the development of myocarditis in this patient.

In this case we could not know which is triggering the myocarditis, is it malabsorption post BPD leading to malnutrition, autoimmune disease could be the cause however autoimmune disease itself could be caused by malnutrition, or just a simple viral infection causing myocarditis. Furthermore, to our knowledge in the literature, no similar case has been described before.

4. CONCLUSION

The current case report suggests that before doing biliopancreatic diversion surgery (BPD) surgery the screening for nutritional deficiencies like vitamins and minerals as well as the screening for connective tissue disease (CTD) might decrease the detrimental consequences and mortality post BPD surgery in patients with hypovitaminosis and connective tissue disease.

This might include looking into the underlying autoimmune inflammatory myositis in routine clinical practice before undergoing biliopancreatic diversion surgery (BPD).

Further research is required to determine the relationship between weight reduction surgery, autoimmune disease, and heart disease specially the myocarditis.

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

CONSENT

As per international standard or university standard, patient(s) written consent has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Bruestle K, Hackner K, Kreye G, Heidecker B. Autoimmunity in acute myocarditis: how

- immunopathogenesis steers new directions for diagnosis and treatment. *Current Cardiology Reports*. 2020 May;22:1-2.
2. Moimas S, Zacchigna S, Merlo M, Buiatti A, Anzini M, Dreas L, et al. Idiopathic dilated cardiomyopathy and persistent viral infection: Lack of association in a controlled study using a quantitative assay. *Heart Lung Circ*. 2012;21(12):787–793. DOI: 10.1016/j.hlc.2012.07.013
3. *Metabolism and Pathophysiology of Bariatric Surgery. Nutrition, Procedures, Outcomes and Adverse Effects*. 1st Edition - November 26, 2016. Editors: Victor R Preedy, Rajkumar Rajendram, Colin R Martin; 2016.
4. Cañas CA, Echeverri AF, Ospina FE, Suso J-P, Agualimpia A, Echeverri A, Bonilla-Abadía F, Tobón GJ. Is bariatric surgery a trigger factor for systemic autoimmune diseases? *J Clin Rheumatol*. 2016;22:89–91. Available: <https://doi.org/10.1097/RHU.00000>
5. Amin Gasmi, Geir Bjørklund, Pavan Kumar Mujawdiya, Yuliya Semenova, Alexandru Dosa, Salva Piscopo, et al. Dietary supplements and bariatric surgery, *Critical Reviews in Food Science and Nutrition*; 2022. DOI: 10.1080/10408398.2022.2046542
6. Yahalom M. *Cardiomyopathy Associated with Zinc Deficiency after Bariatric Surgery*. Thieme Medical Publishers. 2019;333 .
7. Khorgami Z. Early cardiac complications after bariatric surgery: does the type of procedure matter? *Surg Obes Relat Dis*; 2019. PMID: 31151890. DOI: 10.1016/j.soard.2019.03.030
8. Kuno T. *Effects of Bariatric Surgery on Cardiovascular Disease: A Concise Update of Recent Advances*. *Front Cardiovasc Med*; 2019. DOI: 10.3389/fcvm.2019.00094
9. Karina Bruestle. Autoimmunity in Acute Myocarditis: How Immunopathogenesis Steers New Directions for Diagnosis and Treatment. *Curr Cardiol Rep*; 2020. DOI: 10.1007/s11886-020-01278-1
10. Joseph J, Loscalzo J. Selenistasis: Epistatic effects of selenium on cardiovascular phenotype. *Nutrients*. 2013; 5(02):340–358
11. ElAmm CA. Characteristics and Outcomes of Patients With Myocarditis Listed for

- Heart Transplantation. Circulation: Heart Failure; 2016.
Available: <https://doi.org/10.1161/CIRCHEARTFAILURE.116.003259>
12. Bracamonte-Baran W, Čiháková D. Cardiac Autoimmunity: Myocarditis. Adv Exp Med Biol. 2017;1003:187-221.
DOI: 10.1007/978-3-319-57613-8_10
PMID: 28667560; PMCID: PMC5706653.
13. Kang M, An J. Viral Myocarditis. [Updated 2022 Jan 5]. In: Stat Pearls [Internet]. Treasure Island (FL): Stat Pearls Publishing; 2022.
Available: <https://www.ncbi.nlm.nih.gov/books/NBK459259/>
14. Lampejo T. Acute myocarditis: aetiology, diagnosis and management. Clinical Medicine. 2021;21(5):e505–10.
15. Carsten Tschöpe. Myocarditis and inflammatory cardiomyopathy: Current evidence and future directions. Nat Rev Cardiol. 2021;18(3):169–193.

© 2023 Al Bahrani et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/101774>