



Case Study on Production and Marketing Constraints of Sugarcane Farmers in Southern Tamil Nadu, India

Jayanthi, M. ^{a++*}, Mareeswaran, P. ^{b++}, Vijay Prabha, V. ^{a++}
and M. Shree Charan ^{c#}

^a Department of Agriculture, KSAH, KARE, Krishnankoil, India.

^b College of Agriculture, Sri Venkateswara University, Ettayapuram, Kovilpatti, India.

^c KSAH, KARE, Krishnankoil, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJAEES/2023/v41i102170

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/104960>

Original Research Article

Received: 01/07/2023

Accepted: 05/09/2023

Published: 13/09/2023

ABSTRACT

This study assesses the production and marketing constraints faced by sugarcane farmers in Southern Tamil Nadu, India. Sugarcane is an important commercial crop in India. India has emerged as one of the largest producers of sugarcane and sugar in the world with the highest number of sugar factories. India ranks second in Sugarcane production and Tamil Nadu ranks fourth. However, despite its economic importance, farmers encounter various challenges that impede their productivity and profitability. The production constraints primarily revolve around water availability, fluctuating weather patterns and pest infestations. Furthermore, pest attacks, such as

⁺⁺ Assistant Professor;

[#] B.Sc. (Hons) Agriculture Student;

^{*} Corresponding author: E-mail: m.jayanthi@klu.ac.in;

the sugarcane whitefly and red rot disease, lead to yield losses and increased production costs due to the need for pest control measures. The marketing constraints predominantly pertain to inadequate infrastructure, inconsistent pricing and limited access to markets.

Keywords: Constraints; production technology; sugarcane growers.

1. INTRODUCTION

Sugarcane (*Saccharum officinarum*) is a species of tall, perennial grass that is used for sugar production. Sugarcane is an important commercial crop in India [1]. India has emerged as one of the largest producers of sugarcane and sugar in the world with the highest number of sugar factories. India ranks second position in Sugarcane production and Tamil Nadu ranks fourth [2]. The plants are 3-7m tall with stout, jointed, fibrous stalks that are rich in sucrose, which accumulates in the stalk internodes. Sugarcane is propagated by the plantings of cuttings. The planted cane setts should have two or three buds. Sugarcane is a multipurpose crop and has a rich source of food (Sucrose, jaggery and syrups), fiber (cellulose), fodder (green top, bagasse, molasses) fuel and chemicals (Bagasse molasses & alcohol) [3]. During the process of sugar production, the main by-product of the cane sugar industry is Bagasse, Molasses and Press mud. In India, sugarcane is grown as a Kharif crop, with temperature of 21° to 27° C and rainfall of 75-150cm [4]. India ranks second in sugarcane production with 281,170 tonnes next to Brazil. The largest producer of sugarcane in India is Uttar Pradesh, which produced over 177 million tonnes of sugarcane in 2021 [5]. Top Sugarcane Producing States are Uttar Pradesh, Maharashtra, Karnataka, Tamil Nadu and Bihar. Tamil Nadu occupies the fourth position with an average of 37.5 MT annually [6]. Tamil Nadu maintains the record of 107 tonnes/ha and produces an average of 98 tonnes/ha [7]. The major sugarcane-producing districts in Tamil Nadu are Coimbatore, Erode, Tiruppur, Trichy, Theni, Dindugal, Madurai and Tirunelveli. There are 42 sugarcane mills in Tamil Nadu [8,9].

In other countries sugar is a by-product, but in India sugar is the main product produced by sugar mills. India is producing large quantities of sugar because of the country's needs, but nowadays only sugar production is not profitable for sugar mills. The major sugarcane-producing states are Uttar Pradesh (36%), Maharashtra (24%), Karnataka (11%), Tamil Nadu (11%),

Gujarat (4%), Andhra Pradesh (5%), Bihar (3%), Uttarakhand (2%), Haryana (2%) and Punjab (2%). Total sugar production is 100000 quintals and the recovery rate is 11%. The total cost of operation is 9677 lakh rupees (Mishra, 2017). The total profit is 5677 lakh rupees. Current price of sugarcane is 325 rupees per quintal. The objective of this study to identify major constraints faced by sugarcane producers in the production and marketing of sugarcane in selected places in the Tenkasi districts of Tamil Nadu [10].

2. METHODOLOGY

In southern Tamil Nadu, from Tenkasi district, Vasudevanallur Block was purposively selected for this study. In Vasudevanallur Block of the Tenkasi district, the major crop being cultivated are lemon, sugarcane and paddy. Increased sugarcane cultivation is encouraged by the sugar factory in the study region. Six villages from vasudevanullur block were selected based on sugarcane production and primary data are gathered randomly from 30 sugarcane growing farmers in selected villages.

Sugarcane is an irrigated crop, where it is cultivated around 3000 hectares in area. In Vasudevanallur, the planting system followed is Ridges and Furrows. It has been planted in February and harvested in December. Next to lemon, sugarcane is being cultivated in Rayagiri (80 %), Sivagiri (75 %), Arulachi, Ramanathapuram, Ullar Thalavaipuram, Viawanathaperi, Devipattinam, and Thenamlai with 50% of total land area production and most cresers are found in Rayagiri for jaggary production. Predominant varieties used for cultivation are Co 86032, Theni Sevala and Thiruvanamalai Sevala.

3. RESULTS AND DISCUSSION

The Table 1 shows the production and marketing constraints of sugarcane farmers in selected study area.

Table 1. Constraints faced by sugarcane farmers

S. No.	Particulars	No. of Farmers (n=30)	Per cent (%)
1	Climatic factor	25	83.33
2	Pest and Diseases of sugarcane	19	63.33
3	Weed problem in sugarcane field	22	73.33
4	Lack of awareness and scientific knowledge	20	66.67
5	Marketing constraints	15	50

3.1 Climatic Factor

Predominant varieties used for cultivation are Co 86032, Theni Sevala and Thiruvanamalai Sevala which are used for Jaggery production which differ from Rasthali variety. Improper ploughing may cause soil termites that attack the stalk. An excess wind causes the breakage of the stalk. Due to excess rain, Lepidopteran pest spreads through the stalk causing damage of the stalk and yield loss. Due to lack of availability of farm labours and also there is no proper irrigation, there will be possibility of termite growth in the soil. That soil termites affect the sugarcane stalk and leads to stalk death. Sometimes unfavourable conditions like drought, disease and pest attack lessens the weight of sugarcane and decreases the profit. Extreme changes in rainfall have an impact on sucrose yield.

3.2 Pest and Diseases of Sugarcane

In the villages of Rayagiri and Arulachi, small-scale jaggery production faces several challenges, with pests and diseases being among the most significant constraints. These threats attack the sugarcane stalks and leaves, leading to yield losses and the potential spread of infections from one plant to another. Common pests like Internode borer, Top shoot borer, Termites, and Wolly aphids, as well as diseases such as red rot, Wilt, Smut, and yellow leaf disease, pose a continuous risk to the crop's health and overall profitability. Lepidopteran stalk borers cause the splitting of stalk on sugarcane which causes reduced yield and market price. Web formation in stalk and leaves of sugarcane causes stunt growth which is the reason for declining in market price.

3.3 Weed Problem in Sugarcane Field

In addition to these agronomic challenges, the presence of major weeds such as *Cyperus rotendus*, *Cynodon dactylon*, and *Amaranthus viridis* exacerbates the situation by absorbing soil moisture and competing with the sugarcane for essential nutrients, further reducing yields.

Moreover, the scarcity of labour during crucial activities like planting and weeding adds to the farmers' woes, hindering timely and efficient crop management.

3.4 Lack of Awareness and Scientific Knowledge

The lack of knowledge and awareness about the Sustainable Sugarcane Initiative (SSI) has become a significant constraint for farmers in the villages of Rayagiri and Arulachi. SSI as an innovative and eco-friendly approach to sugarcane cultivation has the potential to yield an impressive 70 tonnes per acre, which is more than 30% higher than traditional cultivation practices. However, due to the absence of meetings, workshops, and awareness programs, many farmers in the region are unaware of the benefits and techniques associated with SSI. As a result, they continue to rely on conventional methods, missing out on the opportunity to improve yields, optimize resource use and contribute to sustainable farming practices in the community.

3.5 Marketing Constraints

In response to the lack of a sugar factory in their locality, farmers in the villages of Rayagiri and Arulachi have embraced small-scale jaggery production as a means to utilize their sugarcane yield efficiently. With an impressive yield of 40 tonnes of sugarcane per acre, they have found a sustainable way to extract maximum value from their agricultural efforts.

The process of jaggery production begins with crushing one ton of fresh sugarcane to yield approximately 300 liters of sugarcane juice. To refine the juice and remove impurities, they employ a simple processing technique by adding soda, which helps to excrete sugarcane waste materials like muds, molasses, and dry leaves. This traditional approach, passed down through generations, enables them to create a natural and wholesome jaggery product.




Fig. 1. Production and Marketing constrains of sugarcane farmers

Table 2. SWOT analysis

<p>Strength</p> <ul style="list-style-type: none"> Well grown sugarcane due to soil type and other favourable conditions. Small scale industry like by-products of sugarcane production is being processing in villages. 	<p>Weakness</p> <ul style="list-style-type: none"> Transportation cost is high. Pale colour of the jaggery may reduce the price.
<p>Opportunity</p> <ul style="list-style-type: none"> Own their cresers for production of jaggery which facilitates farm earning, makes better employment to women and facilitates the reach of their by-products to near-by markets. 	<p>Threats</p> <ul style="list-style-type: none"> Damage caused by wild pigs. Scarcity of labour at the time of planting and weeding. Due to lack of sugar industries, farmers face major problems while transportation where they now transport to rajasree (theni)

However, despite their dedication and hard work, the farmers face a significant challenge with the appearance of the jaggery. The final product tends to have a pale colour, which decreases its market price and demand. Consumers often associate darker shades with higher quality and rich flavors in jaggery, leading to an unfair perception of the locally produced jaggery.

4. CONCLUSION

Despite these constraints, sugarcane farming remains important for sugar and ethanol production. Addressing these challenges requires sustainable practices, technological advancements, and better market linkages to ensure the viability and sustainability of sugarcane farming in the long run. Governments, agricultural experts, and stakeholders must collaborate to find innovative solutions and support farmers in overcoming these constraints.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Hoang NV, Furtado A, Botha FC, Simmons BA, Henry RJ. Potential for genetic improvement of sugarcane as a source of biomass for biofuels. *Frontiers in Bioengineering and Biotechnology*. 2015;3: 182.
- Solomon S. The Indian sugar industry: an overview. *Sugar Tech*. 2011 Dec;13:255-65.
- Mishra S. A case study of Akbarpur Sugar mill; 2017. Available:Lwww.academia.edu. Available:https://www.academia.edu/82170602/A_case_study_of_Akbarpur_Sugar_mill?f_ri=75130

4. Papini-Terzi, Flávia S, Rocha, Flávia R, Vêncio, Ricardo ZN, Felix, Juliana M, Branco, Diana S, Waclawovsky, Alessandro J, Del Bem, Luiz EV, Lembke, Carolina G, Costa, Maximiller DL, Nishiyama, Milton Y, Vicentini, Renato. Sugarcane genes associated with sucrose content". BMC Genomics. 2009;10(1):120. DOI:10.1186/1471-2164-10-120. ISSN 1471-2164. PMC 2666766. PMID 19302712.
5. Bhatti SS, Kumar V, Singh N, Sambyal V, Singh J, Katnoria JK, Nagpal AK. Physico-chemical properties and heavy metal contents of soils and kharif crops of Punjab, India. Procedia Environmental Sciences. 2016;35:801-8.
6. Shivashenkaramurthy M, Nayak GV, Channabasappa KS, Patil SB, Rajakumar GR. Effect of nutrient management and cultivars on sugarcane yield and economics. Journal of Pharmacognosy and Phytochemistry. 2021;10(2S): 157-61.
7. Shukla SK, Sharma L, Awasthi SK, Pathak AD. Sugarcane in India. Package of practices for different agro-climatic zones, All Indian Coordinated Research Project on Sugarcane, IISR Lucknow, Uttar Pradesh. 2017 Mar:1-64.
8. Gopakumar K. Indian aquaculture. Journal of Applied Aquaculture. 2003;13(1-2):1-0.
9. Sujatha K, Bajpai AK, Singh RB, NAMDEV A. Direct and indirect effects of production related characters on sugarcane production (*Saccharum spp.*). Trends in Biosciences. 2017;10(42):8803-7.
10. Durai AA, Mahadevaiah C, Gopinath K. Identification of early and mid-late maturing sugarcane varieties for western region of Tamil Nadu. Journal of Sugarcane Research. 2020;10(1).

© 2023 Jayanthy et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/104960>