



Evaluation of Anxiety and Locomotor Behaviour in High Salt-Fed Mice Treated with L-Arginine and Losartan

Justin Atiang Beshel ^{a*}, Justina Andornimye Ashipu ^a,
Linns-Paul Utionkpan ^a, Favour Nyoh Beshel ^a
and Gabriel Otu Ujong ^b

^a Department of Physiology, Faculty of Basic Medical Sciences, College of Medical Sciences, University of Calabar, Calabar, Nigeria.

^b Department of Physiology, Faculty of Basic Medical Sciences, Cross River State University, Calabar, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JABB/2023/v26i10659

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/108632>

Original Research Article

Received: 13/09/2023

Accepted: 18/11/2023

Published: 27/11/2023

ABSTRACT

Salt consumption above dietary guidelines is detrimental to health, it causes impaired cognitive, cardiovascular and neurodegenerative functions. This study aimed at evaluating anxiety and locomotor behaviour in high salt-fed mice treated with L-arginine and losartan. Forty (40) mice weighing between 27-37g were categorized into four (4) parts of ten (10) members. Group I (Control) was served ordinary mice chow and water, group II was given 8% sodium chloride in chow and 1% NaCl in drinkable water (High Salt Diet), group III was given same diet as group II in addition to L-arginine on the 43rd day, group IV was given same as group II in addition to Losartan on the 43rd day. L-arginine and losartan administration lasted for 14 days. The total duration for drug administration and feeding lasted for one month and three weeks. Light-Dark Transition Box

(LDTB) and Elevated+ Maze (EPM) were used to assess anxiety and locomotor behaviour. The results showed a significant ($p < 0.001$) decrease in line crossing, transition frequency, and light box duration, with a corresponding significant ($p < 0.001$) increase in stretch attend posture, grooming frequency duration, close arm entry, dark box duration frequency in the high salt fed group compared with the control. Both L-arginine and Losartan significantly ($p < 0.001$) reversed these changes towards normal. Thus, a high salt diet is implicated in increased anxiety, whereas L-arginine, a nitric oxide precursor, and losartan, an ACE blocker reduced anxiety which in turn improves locomotor behaviour in the mice.

Keywords: Locomotor behavior; losartan decreased anxiety.

1. INTRODUCTION

L-arginine is an amino acid and constitutes the building blocks of proteins. Amino acids are grouped into Non-essential amino acids that are produced in the body and Essential amino acids which is taken externally in daily meal, as such, they must be provided through dietary intake [1,2] opined that L-arginine is sometimes referred to as semi-essential because it becomes essential under certain conditions like pregnancy, infancy, critical illness, and trauma. However, it is a precursor for nitric oxide (signalling molecule) required for carrying out different body functions and processes and includes regulation of blood flow, mitochondrial, and cellular communication [3]. L-arginine has numerous in the body's function and deficiency can disrupt cellular function as well as some organs leading to serious health challenges. L-arginine is usually synthesized from the breakdown of body proteins or produced from dietary proteins. Certain protein-rich foods like meat, poultry, dairy, nuts, soy products, and fish are concentrated with L-arginine. The daily intake of L-arginine from foods is reported to average 4–6 grams [4].

L-arginine has versatile metabolic roles to produce several active biological intermediate substances such as nitric oxide (NO), polyamines, and creatine [2]. However, some research implicates L-arginine as a mediator of the central nervous system and some of its activities are not yet fully understood. Therefore, is possible L-arginine might also affect behaviours such as anxiety and locomotor activities. Nitric oxide has been recognized to play an important role in many behavioural, cognitive and emotional processes [5].

Anxiety and Fear are used mutually, fear refers to alertness in the presence of danger whereas, Anxiety is the unpleasant emotions evoked when fear is stimulated [6]. Anxiety could present as a serious mental ailment where patients are unable

to function normally and worried by every little occurrence (such as cell phone ringing or door slamming). An estimated population of 20-30% in Nigeria is assumed to suffer from mental disorders. A Journal of Nutritional Neuroscience (2005) revealed that a combination of L-lysine and L-arginine ameliorate the effects of stress in subjects with high traits of anxiety.

The impact of excessive dietary additives like NaCl is implicated in negative blood pressure and a decrease in nitric oxide. The high amount of NaCl may reduce L-arginine or decrease its cellular uptake [7] and subsequently cause deficiency of nitric oxide Synthase cofactors [8] and alteration in signalling pathway [9]. Inhibition or uncoupling of nitric oxide synthase will result in reduced nitric oxide production [10]. In hypertension, any combination of these may occur and this mal-function of the nitric oxide pathway serves to increase oxidative stress and perpetual loss of available nitric oxide.

This research therefore aimed at evaluating anxiety and locomotor behaviour in high salt fed mice treated with L-arginine and losartan.

2. MATERIALS AND METHODS

Some materials used include; a light and dark transition box, elevated plus maze (EPM), measuring cylinder, weighing balance, stop watch, L-arginine & losartan, NaCl etc.

2.1 Experimental Animals and Protocol

A total of forty (40) Mice weighing 27-37g were used in the study. They were allowed access to twelve hours of light and feed ad libitum. The Mice were divided into four groups with 10 rats per group. Group I as the control, was given distilled water and normal rat chow. Group II was given 8% NaCl in chow, and 1% NaCl in drinkable water. Group III was fed same as group II, but L-arginine was administered on the 43rd

day. Group IV was fed the same as Group II, but Losartan was administered on the 43rd day of the experiment. L-arginine was administered for nine days and Losartan for four days and was done sequentially. The total duration for feeding and drug administration was 56 days.

At the end of drug administration, the Light-dark transition box and Elevated plus maze were used to assess anxiety and locomotor behaviour.

3. RESULTS

3.1 Frequency of Head Dips in the Elevated Plus Maze Among the Groups

The mean values for head dips in the Control, salt-fed, salt + L-arginine and salt + Losartan group were 10.00 ± 0.17 , 6.40 ± 0.75 , 11.00 ± 0.84 and 9.20 ± 0.86 .

The results show a significant decrease ($P < 0.001$) in head dips in the salt-fed group. Administration of L-arginine and Losartan

resulted in increased head dips and tended towards the normal (control), Fig. 1.

3.2 Frequency of Stretch Attend Posture (SAP) Among the Groups

The mean values for frequency of SAP in the control, salt-fed, salt +L-arginine and salt + losartan group were 4.20 ± 0.58 , 18.80 ± 1.36 , 5.60 ± 0.71 and 8.60 ± 0.93 . The results show a significant increase ($p > 0.001$) in SAP in salt fed group. Administration of L-arginine and losartan improved SAP and tended towards the normal, Fig. 2.

3.3 Grooming Duration (GD) Among the Groups

The GD in the control, salt-fed, salt + L-arginine and salt + Losartan group were 7.60 ± 1.50 , 24.80 ± 0.24 , 3.20 ± 0.58 and 3.40 ± 0.51 respectively. The result showed a longer duration in GD of the salt fed group when compared with the control. Administration of L-arginine and Losartan helped to reduce GD that tends towards the normal control, Fig. 3.




Fig. 1. Frequency of head dip in the control and test groups during the elevated plus maze test
 Values are expressed as mean +SEM, n = 5, ** = $p < 0.0$ vs control, a = $p < 0.05$, c = $p < 0.001$ salt fed




Fig. 2. Stretch attend posture in the control and tests groups during the lelevated plus maze test
Values are expressed as mean +SEM, n = 5, ** = $p < 0.01$, *** = $p < 0.001$ vs control, c = $p < 0.001$ salt fed, x = $p < 0.05$ salt + L-Arginine




Fig. 3. Grooming duration in the control and test groups during the elevated plus maze test
Values are expressed as mean +SEM, n = 5, * = $p < 0.05$, *** = $p < 0.001$ vs control, c = $p < 0.001$ salt fed

3.4 Frequency of Close Arm Entry (CAE) Among the Group

The mean values of CAE in the control, salt-fed, salt + L-arginine and salt + Losartan group were 5.60 ± 1.29 , 6.40 ± 0.89 , 5.00 ± 1.14 and 4.00 ± 0.45 respectively. There was no significant difference among the groups. Perhaps, the salt + Losartan group reduced CAE which indicates some specific role of Losartan, Fig. 4.

3.5 Close Arm Duration (CAD) Among the Groups

The mean values of CAD in the control, salt-fed, salt + L-arginine and salt + Losartan group were 201.60 ± 4.91 , 232.80 ± 9.47 , 201.00 ± 5.97 and 199.40 ± 6.57 respectively. The result showed a significant ($p < 0.001$) increase in CAD in the salt-fed group compared with the control. Administration of L-arginine and Losartan resulted in reduced CAD, salt + Losartan group showed further reduction which may be due to specific actions of Losartan in hypertension. Fig. 5.

3.6 Line Crossing Frequency in Light/Dark Box

The mean values of line crossing frequency in the control, salt-fed, salt + L-arginine and salt + Losartan group were 98.20 ± 3.46 , 66.80 ± 4.09 , 98.40 ± 5.10 and 99.60 ± 2.66 respectively. The result showed a significant decrease ($p < 0.001$) in the salt-fed group compared with the control. Administration of L-arginine and Losartan as observed in other groups increased line crossing frequency, Fig. 6.

3.7 Stretch Attend Posture (SAP) Frequency in Light/Dark Box

The mean values of SAP frequency in the Control, salt-fed, salt + L-arginine and salt + Losartan group were 3.40 ± 0.51 , 13.20 ± 1.20 , 6.80 ± 0.58 and 7.40 ± 0.51 respectively. The result showed a significant increase in SAP frequency in the salt-fed group and was reduced in control. Administration of L-arginine and Losartan in other groups revealed a decrease in SAP frequency towards normal, Fig. 7.




Fig. 4. Frequency of close arm entry in the control and test groups during the elevated plus maze test.

Values are expressed as mean +SEM, n = 5, No significant differences among groups




Fig. 5. Close arm duration in the control and test groups during the light dark transition box test

Values are expressed as mean +SEM, n = 5, ** = p<0.01 vs control, b = p<0.01 salt fed




Fig. 6. Line crossing in the control and test groups during the light dark transition box test

Values are expressed as mean +SEM, n = 5, *** = p<0.001 vs control, c = p<0.001 salt fed




Fig. 7. Stretch attend posture in the control and tests groups during the light dark transition box test

Values are expressed as mean +SEM, n = 5, ** = p<0.01, *** = p<0.001 vs control, c = p<0.001 salt fed

3.8 Lightbox Duration (LBD) in the Light/Dark Box Test

The mean values of LBD in the control, salt-fed, salt +L-arginine and salt + Losartan group were 200.40±5.89, 262.40±12.09, 212.80±3.65 and 219.40±6.02 respectively. The results showed no difference compared with dark box duration (DBD), Fig. 8.

3.9 Freezing Frequency (FF) in the Open Field Maze

The mean values of FF in the control, salt-fed, salt + L-arginine and salt + Losartan group were 0.40±0.24, 1.60±0.24, 0.20±0.20 and 0.40±0.24 respectively. The result showed a significant increase (p<0.001) in FF in the salt-fed group compared to the control. Administration of L-arginine and Losartan restored the FF to normal. Fig. 9.

3.10 Freezing Duration (FD) in the Open Field Maze

The mean values of FD in the control, salt-fed, salt + L-arginine and salt + Losartan group were 0.80±0.49, 9.00±0.71, 0.40±0.40 and 0.40±0.24

respectively. The result showed a significant increase (p<0.001) in the salt-fed group compared to the control. Administration of L-arginine and Losartan decreased FD towards normal. Fig. 10.

4. DISCUSSION AND CONCLUSION

This study evaluates anxiety and locomotor behaviour in high-salt-fed Mice treated with L-arginine and Losartan. High salt intake has been suspected as a contributor to many neuropsychiatric illnesses including; depression and anxiety disorder, and reduces locomotor activities [11,12] demonstrated that excess salt induces anxiety, unusual behaviour and memory impairment in the brain of aging rats. The High energy in synapses, along with increased output of reactive oxygen species is detrimental during increased stress, especially in conditions such as neurodegenerative disorders like Alzheimer's and Parkinson's diseases [13].

Anxiety and locomotive behaviours were assessed using the Elevated+ Maze (EPM) and Light/Dark transition box (LDTB) test. Anxiety behaviour in the mice such as grooming, freezing, stretch attend posture (SAP), line crossing, head dipping, closed arm activity, and

duration of stay in light/dark transition box was studied within the experimental environment. The Elevated+ Maze test is an apparatus which consists of two "open arms and closed arms" (X). The open arms are aversive to mice because they are open and the maze is elevated [14]. The closed arms provide a sense of safety because they are enclosed. Specifically, this research work aimed to evaluate if high salt-fed mice treated with L-arginine had any improvement in behaviour during the experiment. Behaviour like transition frequency from open to closed arms, the grooming duration, urination and defecation were observed. The light-dark transition box (LDTB) is an unconditioned anxiety behaviour test, it involves exploring an environment while avoiding bright light [15]. The box is divided into two compartments, light and dark. Generally, the dark compartment is smaller and is considered

"safe" for mice. Behaviour like the number of line crossing, head dipping and frequency, stretch attend posture, etc. were used to measure anxiety and locomotor behaviour. A low frequency of these behaviours indicates a high anxiety level.

In the elevated plus maze, the frequency of stretch attend posture (SAP), grooming frequency (GF), Grooming duration (GD), Close arm entry (CAE) and Close arm duration (CAD) increased in the salt group. In the L-arginine group, SAP, GF, GD, CAE and CAD were reduced. In the Losartan group, SAP, GF, GD, CAE and CAD were reduced. The EPM test indicates that high salt-fed mice are observed with increased anxiety while L-arginine and Losartan showed decreased anxiety and improved behaviour.




Fig. 8. Light box duration in the control and test groups during the light dark transition box test

Values are expressed as mean +SEM, n = 5, *** = p<0.001 vs control, c = p<0.001 salt fed




Fig. 9. Freezing frequency in the control and test groups during the open field maze test
Values are expressed as mean +SEM, $n = 5$, ** = $p < 0.01$ vs control, $b = p < 0.01$ salt fed




Fig. 10. Freezing duration in the control and test groups during the open field maze test
Values are expressed as mean +SEM, $n = 5$, *** = $p < 0.001$ vs control, $c = p < 0.001$ salt fed

In the Light-dark test box (LDTB); SAP, GF, GD and DBD (dark box duration) are increased in the salt group. In light box duration (LBD); line crossing (LC) and transition frequency is also reduced in the salt group than in the Control. In the L-arginine group, LBD and transition frequency are increased while SAP, GF, GD, and DBD decrease compared to the salt group. In the Losartan group, LBD and transition frequency increased while SAP, GF and DBD decreased compared to the salt group. It is observed from LDBT that high salt increased anxiety and reduced locomotor activities while L-arginine and Losartan decreased anxiety and improved locomotor behaviour.

This study is similar to earlier report [16] which opined that L-arginine helps to reduce anxiety and negative effects of stress, improves cognitive function and exploration in mice [17,18] revealed that L-arginine improves locomotor activities in rats.

Observations from this work revealed that both L-arginine and Losartan reduce anxiety, as well as improve locomotor behaviour in mice.

ETHICAL APPROVAL

Animal Ethics Committee approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kumar KV, Das UN. Are free radicals involved in the pathobiology of human essential hypertension? *Free Radic Res Commun.* 2013;19:59–66.
2. Sagar S, Kallo IJ, Kaul N, Ganguly NK, Sharma BK. Oxygen free radicals in essential hypertension. *Mol Cell Biochem.* 2011;111:103–108.
3. Beckman JS, Beckman TW, Chen J, Marshall PA, Freeman BA. Apparent hydroxyl radical production by peroxynitrite: Implications for endothelial injury from nitric oxide and superoxide. *Proc Natl Acad Sci USA.* 2017; 87:1620–1624.
4. Kimura G, Murakami K, Kojima S. The association between salt sensitivity of blood pressure and family history of hypertension. *Clin Exp Pharmacol Physiol Suppl.* 2017;20:61–63.
5. Gilman TL, Mitchell NC, Dawns LC, Toney GM. Neuro inflammation contributes to high salt intake-augmented neuronal activation and active coping responses to acute stress. *Int J Neuro psychopharmacol.* 2019;22:137–142.
6. Beck AT, Emery G, Greenberg RL. *Anxiety disorders & phobia: A cognitive perspective* Basic Books/Hachette Book Group. 2005;10.
7. Mrug S, Orihuela C, Mrug M, Sanders P. Sodium and potassium excretion predict increased depression in urban adolescents. *Physiol Rep.* 2029;7:e14213.
8. Alp NJ, Channon KM. Regulation of endothelial nitric oxide synthase by tetrahydrobiopterin in vascular disease. *Arterioscler Thromb Vasc Biol.* 2014;24:413–420.
9. Zecchin HG, Bezerra RM, Carnevali JB. Insulin signaling pathways in aorta and muscle from two animal models of insulin resistance – the obese middle-aged and the spontaneously hypertensive rats. *Diabetologia.* 2003;46:479–91.
10. Zalba G, Beaumont FJ, San-Jose G, Fortuno A, Fortuno MA, Diez J. Is the balance between nitric oxide and superoxide altered in spontaneously hypertensive rats with endothelial dysfunction? *Nephrol Dial Transplant.* 2001;23:2–5.
11. Mitchell NC, Gilman TL, Daws LC, Toney GM. High salt intake enhances swim stress-induced PVN vasopressin cell activation and active stress coping. *Psycho Neuro Endocrinology.* 2018; 93:29–38.
12. Chugh G, Asghar M, Patki G, Bohat R, Jafri F, Allam F. A high-salt diet further impairs age-associated declines in cognitive behavioral and cardiovascular functions in male Fischer brown Norway rats. *J Nutr.* 2013;143:1406–1413.
13. Chen J, Kuhlencordt P, Urano F, Ichinose H, Astern J, Huang PL. Effects of chronic treatment with L-arginine on atherosclerosis in apoE knockout and apoE/inducible NO synthase double-knockout mice. *Arterioscler Thromb Vasc Biol.* 2003;23:97–103.
14. Lister R. Ethologically-based animal models of anxiety disorders. *Pharmacol Ther.* 1990; 46:321–340.

15. Maren S. Seeking a spotless mind: extinction deconsolidation and erasure of fear memory. *Neuron*. 2013;70:830–845.
16. Pervin M, Unno K, Konishi T, Nakamura Y. L-Arginine Exerts Excellent Anti-Stress Effects on Stress-Induced Shortened Lifespan Cognitive Decline and Depression. *Int J Mol Sci*. 2021; 22(2):508.
17. Pique JM, Whittle BJ, Esplugues JV. The vasodilator role of endogenous nitric oxide in the rat gastric microcirculation. *Eur J Pharmacol*. 2009;174(2–3):293-296.
18. Le Doux JE. *Anxious: Using the Brain to Understand and Treat Fear and Anxiety*. 1st edition, Penguin Group (USA) LLC. 2015;50.

© 2023 Beshel et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/108632>