



Physiochemical Properties of Different Oil Media during Deep Fat Frying of Akara Chips

G. C. Nneji^{a*} and S. I. Oluka^a

^a *Department of Agricultural and Bioresource Engineering, Enugu State University of Science and Technology, Agbani, Enugu, Nigeria.*

Authors' contributions

This work was carried out in collaboration between the two authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJB2T/2024/v10i1194

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/96500>

Original Research Article

Received: 18/12/2022

Accepted: 22/02/2023

Published: 15/01/2024

ABSTRACT

The specific objective is to determine the physiochemical characteristics of frying and fresh palm oil, groundnut oil and olive oil under deep fat frying condition at a temperature of 120⁰ Celsius in the production of akara chips derived from cowpea paste. Standard methods were used in determining physiochemical properties of the different oil samples. The research was carried out in the Department of Agricultural and Bioresources Engineering, Enugu State University of Science and Technology, Agbani Enugu State, Nigeria. The results showed that there was a significant ($p < 0.05$) variation among the three fresh (palm oil, groundnut oil and olive oil) and frying (palm oil, groundnut oil and olive oil) oil media respectively on the solidification point, iodine concentration, moisture content, melting point, fat levels and viscometric values. Frying reduced the moisture content and iodine value of palm, groundnut and olive oil than the non-frying, Fresh palm oil, groundnut oil and olive oil were high in viscometric and saponification values respectively.

*Corresponding author: E-mail: eejike43@yahoo.com;

Keywords: Deep fat frying; akara chips; palm oil; groundnut; oil; olive oil.

1. INTRODUCTION

Fried akara chips is a popular food vended delicacy in the streets of Sub-Sahara Africa. It is a fast conveniently made food products that meet time requirement for a developing country where people eat outdoors. Thus, it is consumed as a snack. Most of the fried akara chips consumed in Nigeria are fried in groundnut oil. Therefore, efforts to replace groundnut oil with palm oil and olive oil in the frying process for commercial purposes is of a great practical interest. Frying also initiates different physical and chemical changes which results in extensive degradation of oil. During this process, the fried food absorbs some amount of oil and as a result, proportions of degraded products from the oil are accumulated by the fried food. [1 and 2] reported that “frequent frying of vegetable oils leads to deterioration and lipid oxidation results in the formation of peroxides which are responsible for primary oxidation in the oil. Degradation of vegetable oils during frying result in loss of nutritive value and causes damaging health effects due to the toxic substances produced”. [3] also reported that “repeated usage of vegetable oil for frying lowers the smoke point which makes the oil to smoke on heating at a lower temperature”. There are physical and chemical qualities that can be used to identify oil on or after frying. Some of these characteristics are melting point, solidification point, viscometric studies, moisture content, iodine value, saponification value, peroxide value and acidity. The main objective of this study is to determine the physiochemical characteristics of different oil media use in frying akara chips. The specific objective is to determine the physiochemical characteristics of frying and fresh palm oil, groundnut oil and olive oil under deep fat frying condition in the production of akara chips derived from cowpea paste.

2. MATERIALS AND METHODS

2.1 Materials

Cowpea was purchased at the Enugu State Agricultural Development Programme office in the city of Enugu (Fig. 1a). Palm oil, groundnut oil and olive oil samples were purchased from local market in Enugu (Fig. 1b). The cowpea samples were selected and sorted taking into account the size and that they did not present damages by manipulation.



Fig. 1a. Cowpea samples



Fig. 1b. Oil samples

2.2 Akara Chips Preparation

Cowpea samples were sorted and soaked in water for three minutes before mashing to remove chaff. It was then grounded into paste, weighed and the paste made in form of balls ready for frying. The fryer tank was parallelepiped-shaped and made of stainless steel. Its dimensions were: length 300×10^{-3} m; width 150×10^{-3} m; height 200×10^{-3} m. The bath had a volume of five liters. The temperature of the frying bath was determined by simultaneously measuring the temperatures with attached thermometer.

2.3 Deep Fat Frying

Frying was carried out using processed palm oil, olive oil and groundnut oil. Each of the oil

samples weighed 1189 g and poured in the fryer and heated to 120^o Celsius. A total of 121 g of each of the pre-treated cowpea samples was weighed. The oil was allowed to heat up to first temperature of 120^oC at three to seven minutes before the samples were introduced. The full fried samples of chips were recovered after ten minutes of frying. Akara balls were placed on absorbent paper to remove the oil remaining on the surface.

2.4 Determination of Physicochemical Properties.

2.4.1 Moisture content

The percent loss in weight of oil sample heated in a drying oven at 105^oC for 2 h and weighed at an interval of 30 min till a constant weight was obtained. Initial weight of oil sample = a gm, final weight of oil sample after drying = b gm, the dry weight percent = (a/b)*100. Moisture content (x %) = (1-(a/b))*100.

2.4.2 Solidification point

The oil in a capillary tube was allowed to flat in watch glass on a water trough. Blocks of ice were added continuously until the oil solidified and the temperature recorded.

2.4.3 Viscometric studies

“The time taken for the oil to fall between the two graduation marks on Oswalds viscometer at room temperature was recorded” [4].

2.4.4 Acid value determination [5]

“Two grams of the oil sample was dissolved in 25ml diethyl ether with 25ml ethanol was titrated with 0.1M NaOH solution and 1ml of phenolphthalein indicator until a faint pink colour persisted for 15 seconds”.

$$\text{Acid value} = \frac{\text{titration (ml)} \times 5.61}{\text{weight of oil sample used}} \quad (1)$$

2.4.5 Determination of saponification value

Two grams of the oil was refluxed with 25 mL of alcoholic potassium hydroxide solution (0.5 M) for 1 h with frequent shaking. The excess alkali was titrated with 0.5 M hydrochloric acid and 1mL of phenolphthalein indicator. A blank titration was carried out alongside and the

$$\text{Saponification value} = \frac{(\text{blank-titration value}) \times 28.05}{\text{weight of oil sample (g)}} \quad \text{thus} \quad (2)$$

2.4.6 Determination of iodine value

Twenty-five milliliters of iodine monochloride was added to 1.01 g of the oil, stoppered and left to stand in the dark alongside a blank without the oil sample and 10 mL of chloroform added instead for 1 h. The flask was rinsed with 50 mL of distilled water and 10 ml of 10% KI solution was added. The liberated iodine was immediately titrated with 0.1 M Na₂S₂O₃ until the iodine solution was brownish yellow then 1 mL of starch solution indicator was added. The titration was continued until the developed blue color disappeared.

$$\text{The volume of the Na}_2\text{S}_2\text{O}_3 \text{ was used to calculate the Iodine value} = \frac{(\text{blank-titration}) \times \text{molarities of Na}_2\text{S}_2\text{O}_3 \times 12.69}{\text{weight of oil sample (g)}} \quad (3)$$

2.3 Data Analyses

Data collected was subjected to analysis of variance (ANOVA) test for completely randomized block design as outlined by [6]. Significant means was separated using Fisher's least significant difference (F-LSD) at 5% probability level and t-test was to compare the differences between means. Statistical analysis was executed using [7] Statistical Software.

3. RESULTS AND DISCUSSION

The results presented in Table 1a revealed the physicochemical composition of fresh and frying oil during deep fat frying of akara chips. The results showed that there was a significant (p< 0.05) variation among the three fresh (palm oil, groundnut oil and olive oil) and frying (palm oil, groundnut oil and olive oil) oil media respectively on the solidification point. Fresh palm oil had significantly (p< 0.05) the highest solidification point followed by fresh groundnut oil and last was recorded in fresh olive oil. The solidification point of the frying olive oil was significantly (p< 0.05) the highest than the frying palm oil and frying groundnut oil respectively. Frying palm oil, groundnut oil, and olive oil solidified more than the fresh palm oil, groundnut oil, and olive oil respectively.

Table 1a. Physiochemical composition of fresh and frying oil during deep fat frying of akara chips

Samples	Solidification point			Melting point (°C)			Moisture content (%)			Saponification value		
	Fresh	Frying	t-test	Fresh	Frying	t-test	Fresh	Frying	t-test	Fresh	Frying	t-test
Palm oil	19	20	0.4	38	39	0.3	5	4.99	0.1	235.62	194.94	21.11
Groundnut oil	6	10	1.7	28	38	6.2	4.88	4.96	0.2	19.63	228.60	67.22
Olive oil	3	22	5.3	38	38	1	4.24	4.96	0.2	325.38	211.74	34.33
FLSD (0.05)	2	1.5		5.2	0.4		0.3	0.1		76.22	25.11	

Table 1b. Physiochemical composition of fresh and frying oil during deep fat frying of akara chips

Samples	Peroxide value			Viscometric value (s)			Value			Fat content		
	Fresh	Frying	t-test	Fresh	Frying	t-test	Fresh	Frying	t-test	Fresh	Frying	t-test
Palm oil	17.6	31.4	7.2	467.7	362.3	31.2	25.30	13.20	5.3	1.80	2.00	0.1
Groundnut oil	5.0	5.0	1	126.1	173.8	20.4	28.40	25.30	2.1	4.40	4.50	0.1
Olive oil	4.7	5.8	0.1	32.20	166.1	45.6	29.80	25.00	2.4	4.80	4.70	0.1
FLSD (0.05)	0.1	0.1		19.3	28.2		0.3	0.2		0.4	0.2	

Furthermore, frying palm oil (39^o Celsius), groundnut oil (38^o Celsius) and olive oil (38^o Celsius) significantly ($p < 0.05$) melted faster than the fresh palm oil (38^o Celsius), groundnut oil (28^o Celsius) and olive oil (38^o Celsius) under deep fat frying condition at a temperature of 120^o Celsius. Similarly, frying palm oil, (39^o Celsius) groundnut oil (38^o Celsius) and olive oil (38^o Celsius) had significantly ($p < 0.05$) higher melting point value than the fresh palm oil (38^o Celsius), ground nut oil (28^o Celsius) and olive oil (38^o Celsius) respectively under deep fat frying (Table 1a). Moreso, in Table 1a, there was a significant ($p < 0.05$) differences among the three fresh and frying (palm oil, groundnut oil and olive oil) oil media respectively on the melting point value. Fresh palm oil (5%) had significantly ($p < 0.05$) the lowest melting point followed by fresh groundnut oil (4.88%) and the least was observed in fresh olive oil (4.24%). The melting point of the frying palm oil (4.99%) was significantly ($p < 0.05$) the highest than the frying groundnut (4.96%) and frying olive oil (4.96%) respectively. The saponification (Table 1a) value of fresh palm oil (255.62) and olive oil (325.38) were significantly ($p < 0.05$) higher than the frying palm oil (194.94) and olive oil (211.74) respectively. While in the frying groundnut oil (226.60) saponification value was higher than the fresh groundnut oil (19.63) under deep fat frying at a temperature of 120^o Celsius. There was a significant difference ($p < 0.05$) in the peroxide value among the fresh and frying palm oil, groundnut oil and olive oil under deep fat frying at a temperature of 120^o Celsius (Table 1b). Frying oil [olive oil (5.8) < palm oil (31.4)] had significantly more peroxide value than the non-frying oil [olive oil (4.7) < palm oil (17.6)] respectively. Both non-frying and frying groundnut had the same peroxide value [5]. Table 1b also showed that there was a significant ($p < 0.05$) variations on the visometric value of the three oil media. Fresh palm oil (467.7) had the highest ability to resist flow due to friction than fresh groundnut (126.1) and olive oil (32.20) respectively. The visometric value of the frying oil increased in the order; palm oil (362.3) > groundnut oil (173.8) > olive oil (166.1) respectively. The iodine level of the fresh olive oil (29.80) was significantly ($p < 0.05$) the highest among the frying and non-frying palm oil, groundnut oil and frying olive oil respectively. Fresh palm oil (25.30), groundnut oil (28.40), and olive oil (29.80) contained more iodine than the frying palm oil (13.20), groundnut oil (25.30), and olive oil (25) respectively under deep fat frying condition. The fat content (Table 1 b) of fresh olive oil (4.80) was significantly ($p < 0.05$) the

highest than the fresh groundnut oil (4.40). The least was recorded in the fresh palm oil which had a fat concentration of 1.80. Similar trend was observed in the frying oil media, under deep fat frying condition, frying olive oil (4.70) contained the highest fat levels followed by groundnut oil (4.50) and palm oil (2.00) respectively. [8] reported that "during frying, oils are oxidized to form hydroperoxides that can decompose further to yield the secondary oxidation products, such as alcohols, ketones, aldehydes and acids". "In deep fat frying, however, acids are also produced by hydrolysis of fats to form free fatty acids" [9]. According to [10], "low moisture in oil decreases the fatty acids content while high moisture leads to its increase".

4. CONCLUSION

The results of the physiochemical composition of the oil are useful in food processing industry. The results obtained from this work provide vital information on the physiochemical properties of fresh and frying oil. It can be concluded that frying reduced the moisture content and iodine value of palm, groundnut and olive oil than the non-frying, Fresh palm oil, groundnut oil and olive oil were high in viscometric and saponification values respectively.

ACKNOWLEDGMENTS

The authors are grateful to Department of Agricultural and Bioresource Engineering and Department of Industrial Chemistry, ESUT, for providing research enabling environment and equipment.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Codex Alimentarius Commission/ FAO/ WHO Food Standards, Codex standards for named vegetable oils: CODEX STAN 210. CXSTAN 210-1999; 2000.
2. Emelike NJT, Ujong AE, Achinewu SC. Physiochemical and antioxidant properties of oils used by local vendors in D/ line-portharcourt, River State. *Agricultural and Food Science Research*. 2020;7(1): 89-96.

3. Good J. Healthiest cooking oil chart with smoke points, baseline of health foundation; 2012.
Available:<https://www.jonbarron.org/article/healthiest-cooking-oil-chart-smokepoints>
4. Ikhuoria EU, Maliki M. Characterization of avocado pear (*Persea americana*) and African pear (*Dacryodes edulis*) extracts. Afr. J. Biotechnol. 2007;6(7):950-952.
5. Ejim IF, Kamen FL. Physicochemical characterization of algae oil from microalgae of Nike Lake Enugu. Journal of Engineering and Applied Science. 2013;5(1):181-187.
6. Gomez KA, Gomez AA. Statistical procedures for agricultural research. 2nd edition, John Wiley and Sons, New York. 1984;680.
7. Genstat for Windows, Discovery. 3rd Edition, Lawes Agricultural Trust, Rothamsted Experimental Station UK; 2007.
8. Diop A, Sarr SO, Ndao S, Cissé M, Baldé M, Ndiaye B, Diop YM. Effect of deep-fat frying on chemical properties of edible vegetable oils used by Senegalese households. African journal of agriculture, nutrition and development. 2014;14 (16).
9. Steveson, SG, Vaisey-Genser M, Eskin NAM. Quality control in the use of deep frying oils. J. Am Oil chem Soc. 1984; 61:1102-8.
10. Emelike .N, Akusu .M and Ujong A. Antioxidant and physicochemical properties of oil extracted from cashew (*Anacardium occidentale* L) Kernals. International journal of food science and nutrition. 2017;2:122-128.

© 2024 Nneji and Oluka; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/96500>*