



Effect of Different Potting Media on Seeds Germination & Growth of Dahiman (*Cordia macleodii* Hook.) in Nursery

Neha Minj^{a++*}, Pratap Toppo^{a#}, S.S. Tuteja^{b†}, Lalji Singh^{a‡},
Manish Kumar Mankur^{a^}, Alok Singh Bargah^{c^}
and Sakshi Verma^{a++}

^a Department of Forestry, College of Agriculture, IGKV, Raipur (Chhattisgarh), India.

^b Centre of Excellence on NTFPs & MAPs, College of Agriculture, IGKV, Raipur (Chhattisgarh), India.

^c Department of Forest Products and Utilization, College of Forestry and Research Station, MGUVV, Sankra Patan Durg (Chhattisgarh), India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i94952>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/121735>

Original Research Article

Received: 14/06/2024

Accepted: 16/08/2024

Published: 24/08/2024

⁺⁺M.Sc. Research Scholar

[#]Assistant Professor

[†]Principal Scientist & Technical Advisor

[‡]Professor

[^]Ph.D. Research Scholar

*Corresponding author: E-mail: neha05minj03@gmail.com;

Cite as: Minj, Neha, Pratap Toppo, S.S. Tuteja, Lalji Singh, Manish Kumar Mankur, Alok Singh Bargah, and Sakshi Verma. 2024. "Effect of Different Potting Media on Seeds Germination & Growth of Dahiman (*Cordia macleodii* Hook.) in Nursery". *International Journal of Plant & Soil Science* 36 (9):68-77. <https://doi.org/10.9734/ijpss/2024/v36i94952>.

ABSTRACT

The study was conducted at the Herbal Garden, IGKV, Raipur (C.G.) from July to January during the 2023-24 session. The study aimed to evaluate the impact of various potting mixtures on the growth of Dahiman (*Cordia macleodii*) in a nursery setting. The experiment used a Complete Randomized Design (CRD) with three replications and nine different treatments. The treatments involved different combinations of potting mixtures: T₁ (sand + soil + FYM, 1:1:1), T₂ (soil + sand + vermicompost, 1:1:1), T₃ (soil + vermicompost + FYM, 1:1:1), T₄ (sand + soil + lemongrass compost, 1:1:1), T₅ (sand + soil + FYM, 2:1:1), T₆ (sand + soil + vermicompost, 1:1:1), T₇ (sand + soil + vermicompost, 1:1:2), T₈ (sand + soil + FYM, 1:1:2), and T₉ (control). The T₇ treatment (sand + soil + vermicompost, 1:1:2) stood out as the most effective, significantly improving all growth parameters for both Dahiman. These findings indicate that the T₇ treatment (sand + soil + vermicompost, 1:1:2) is the best choice for promoting healthy growth in Dahiman seedlings.

Keywords: Seedlings; complete randomized design; FYM & vermicompost.

1. INTRODUCTION

The rare species *Cordia macleodii* is also referred to as Dahiman or Dahiphallas. The family Boraginaceae is where it belongs. The original home of *Cordia macleodii* is India. In India, it is commonly found in drug-addled, wet deciduous forests like Chhattisgarh. The tree is between 9 and 12 meters tall, with light green bark that is 12 to 15 mm thick. Broadly oval, glossy, dark green leaves are present. The hue of flowers is yellowish-white. The fruits appear in March through June, and the blooms appear in March through April. According to a number of pharmacological investigations, *Cordia macleodii* plants exhibit wound healing, hepatoprotective, analgesic, anti-inflammatory, antibacterial, and antioxidant properties.

Seedlings are small plants with enough potential to grow into mature trees. These seedlings grow naturally, but they can also be propagated by seeding and germinating, which helps them to sprout quicker and boost their chances of survival. The factors must be optimal to produce a health and vigorous seedlings, and among them, the genes and the nursery environment determine to a large degree how a tree can survive [1]. Development of seedlings begins with the acquisition of viable germplasm or rootstocks that can be germinated or developed through vegetative means [2]

All too often, nurseries operate with minimal inputs and outdated techniques, and therefore produce poor seedlings [1]. Therefore, to enhance the quality of seedlings produced, nurseries should demonstrate high quality standards that incorporates structured planning and quality control, appropriate substrates and

containers, nursery hygiene, and good equipment [1].

Nurseries provide the means to control moisture, light physical and chemical soil constituents in such a way as to produce healthy and uniform seedlings for planting [3]. Several factors can influence seedling growth in the nursery. But potting substrate is the most important factor from the nursery environment. This is because, first it is basically a plant's first food; secondly, it physically supports a growing seedling and thirdly, it stores and supplies nutrients, water and air to the root system [1].

2. MATERIALS AND METHODS

The experiment has been carried out during season of 2023 – 2024 in Herbal Garden of Indira Gandhi Krishi Vishwavidyalya, Raipur. Completely Randomized Design (CRD) was employed with nine (9) treatments and three (3) replication each. The treatment details are . : T₁ (sand + soil + FYM, 1:1:1), T₂ (soil + sand + vermicompost, 1:1:1), T₃ (soil + vermicompost + FYM, 1:1:1), T₄ (sand + soil + lemongrass compost, 1:1:1), T₅ (sand + soil + FYM, 2:1:1), T₆ (sand + soil + vermicompost, 1:1:1), T₇ (sand + soil + vermicompost, 1:1:2), T₈ (sand + soil + FYM, 1:1:2), and T₉ (control soil only) .The seeds of (*Cordia macleodii* Hook.)were sown in 2 cm depth in polybags prepared with different potting mixture on July,2023.

The polybags were set up in accordance with the experimental design, with three replications of each type of polybag per treatment. The seeds germination and growth parameters of the seedlings were observed, and the mean of these data was computed.

The height of the plant was measured every month for all the treatments using a meter scale in centimeters. The height from the ground level to the terminal apex of the plant was measured and their mean was recorded. The diameter of the plant was measured using a Vernier caliper at 1 cm above the ground level. The collar diameter was measured every month for all the treatments and their mean was recorded. The leaves number for each plant was counted every month for all the treatments and their mean was recorded.

2.1 Seeds Germination Percentage

Germination percentage [4] was calculated by number of germinating seedlings divided by the total number of seeds sown in poly bags and multiplied by 100.

2.2 Survival Percentage

Total number of survived seedlings was counted from each treatment. Success percentage was calculated by using following formula:

$$\text{SURVIVAL (\%)} = \frac{\text{total number of seedlings survived}}{\text{total number of germinated seedlings}} \times 100$$

3. RESULTS AND DISCUSSION

The current investigation was conducted to know the effect of different potting media on seeds germination and growth of Dahiman (*Cordia macleodii* Hook.) in nursery. The findings from the current study are presented under suitable headings, along with tables and figures.

Germination percentage: The study revealed that T₇ Sand + Soil + Vermicompost 1:1:2 had the highest germination percentage (60.16) % of *Cordia macleodii*, whereas T₉ Control (soil only) had the lowest germination percentage (13.46) % (Table 1 & Fig. 1). The results show highest germination percentage due to the more portion of vermicompost as it contains a lot of mineral nutrients, active soil enzymes and microbes [5]. Vermicomposts improve seed germination, seedling vigor, and plant productivity more than what would have been possible from inorganic mineral nutrient sources, while using as little as 10–40 % of the total plant rooting volume [6,7,8].

The findings of this study are consistent with the findings of Gawankar *et al.*, [9] who found that the effect of different potting media on seed germination and seedling growth in jackfruit

(*Artocarpus heterophyllus* Lam.) containing soil, vermicompost, cocopeat, and rice husk in a 1:1:1:1 proportion was best for obtaining the highest germination percentage (93.0%), 11.88 days for initiation of germination, 22.88 days for 50% germination, and 24.25 days for total germination. Panchal *et al.*, [10], Meena *et al.*, [11] and Purwantoro, [12] reported that use of vermicompost and rice husk was useful as they had components which favour the nutritional status, pH levels and organic carbon content. The findings are consistent with previous studies on papaya by Arvind [13], and Ramteke *et al.*, [14].

Survival percentage: The study revealed that T₇ Sand + Soil + Vermicompost 1:1:2 had the highest survival percentage (37.4) % of *Cordia macleodii*, whereas T₉ Control (soil only) had the least impact on seedling survival percentage (15.06) % (Table 1 & Fig. 1). The results shows very less survival rate of seedlings while more germination percentage rate is due to climatic factors as the seedlings were in open field condition therefore there is less survival percentage of seedlings..

According to Mulugeta's [15] study, which examined the effects of various potting mixtures on the growth and survival of *Albizia Gummifera* and *Cordia Africana* seedlings, the survival rates of *C. africana* (98.333%) and *A. gummifera* (95.000%) were considerably higher in T₄ than in T₁₀. These results are consistent with the findings of this study.

Plant height (cm): The study revealed that the height of *Cordia macleodii* seedlings increased significantly with the number of days in the plant height, revealing a significant impact of the treatment. The treatments with the statistically highest effects on plant height in 30 days are T₇ sand + soil+ vermicompost @ 1:1:2 (6.7 cm) and T₈ sand + soil+ FYM @ 1:1:2 (6.16 cm). The highest plant height measured in 60 days was 15.53 cm in T₇ sand+ soil+ vermicompost at a ratio of 1:1:2. The highest plant height measured in 90 days in T₇ sand + soil+ vermicompost @ 1:1:2 (22.4 cm) The maximum plant height in T₇ sand+ soil+ vermicompost at 1:1:2 (30.9 cm) was measured after 120 days. The highest plant height was measured in T₇ sand+ soil+ vermicompost @ 1:1:2 in 150 days (35.56 cm), and in T₇ sand + soil + vermicompost @ 1:1:2 in 180 days (36.7 cm) (Table 1 & Fig. 2). The highest plant results as vermicompost stimulates to influence the microbial activity of soil,

increases the availability of oxygen, maintains normal soil temperature, increases soil porosity and infiltration of water, improves nutrient content and increases growth, yield and quality of the plant [16].

The results of this study are consistent with those of Ghising *et al.*, [17], who found that a 2:1:1 potting media combination of soil, sand, and vermicompost produced the highest heights for *Tectona grandis* seedlings. The chemical and physical characteristics of the rooting media were enhanced by the addition of cocopeat, farmyard manure, and vermicompost to various media, which promoted the growth and development of seedlings [10]. Vermicompost is also beneficial to the general growth and development of the plant. Abirami *et al.*, [18] The same results on *Pinus gerardiana* seedling growth were also noted by Kumar *et al.*, [19], who found that growing media containing soil, sand, and vermicompost produced the highest heights. The growth of *Grevillea robusta* seedlings raised in a mixture of soil, sand, and vermicompost was found to be outstanding by Khaple *et al.*, [20] and Sood and Ram [21] in the case of *Oroxylum indicum*.

Collar diameter (mm). The study revealed that the collar diameter of the *Cordia macleodii* seedlings increased significantly with the number of days in the collar, revealing a significant treatment effect. The treatments with the statistically highest effect on collar diameter in 30 days are T₇ sand+ soil+ vermicompost @ 1:1:2 (1.69 mm) and T₈ sand + soil+ FYM @ 1:1:2 (1.59 mm). T₇ sand + soil+ vermicompost @ 1:1:2 recorded the maximum collar diameter in 60 days (2.79 mm), while T₇ sand+ soil+ vermicompost @ 1:1:2 recorded the maximum collar diameter in 90 days (3.89 mm). The greatest collar diameter in T₇ sand+ soil+ vermicompost @ 1:1:2 was measured after 120 days (5.92 mm). The greatest collar diameter was measured in T₇ sand+ soil+ vermicompost @ 1:1:2 (6.92 mm) in 150 days, and in T₇ sand+ soil+ vermicompost @ 1:1:2 (7.66 mm) in 180

Days (Table 1 & Fig. 3). The highest collar diameter results as vermicompost stimulates to influence the microbial activity of soil, increases the availability of oxygen, maintains normal soil temperature, increases soil porosity and infiltration of water, improves nutrient content and increases growth, yield and quality of the plant [16].

The results also have resemblance to Ghising *et al.*, [17] observations of the maximum collar

diameter for *Tectona grandis* seedlings on a mixture of soil, sand, and vermicompost. Additionally, the results closely match those of *Santalum album* by Annapurna *et al.* [22] and *Grevillea robusta* by Khaple *et al.*, [20], as well as *Oroxylum indicum* by Sood and Ram [21]. Additionally, according to Kumar *et al.*, [19], *Pinus gerardiana* collar diameters are noticeably larger in soil that has been combined with vermicompost.

Number of leaves. The study revealed that there was a rise in the number of leaves in the *Cordia macleodii* seedlings with each passing day, revealing the significant effect of the treatment. The treatments with the statistically highest effect on the number of leaves after 30 days are T₇ sand + soil + vermicompost @ 1:1:2 (4.66) and T₈ sand + soil + FYM @ 1:1:2 (4.26). In T₇ sand + soil + vermicompost @ 1:1:2 (9.56), the most leaves in 60 days were recorded. T₇ sand + soil + vermicompost @ 1:1:2 achieved the highest number of leaves in 90 days (14.5). T₇ sand + soil + vermicompost @ 1:1:2 recorded the highest number of leaves in 120 days (19.63). In T₇ sand + soil + vermicompost @ 1:1:2 during 150 days, the highest number of leaves was reported (24.83) The highest number of leaves in 180 days was seen in T₇ sand+ soil+ vermicompost @ 1:1:2 (29.93) (Table 1 & Fig. 4).

There were more leaves in T₇. The majority of plants food is produced in leaves, which are impacted by a variety of elements, the most important of which is the soil. The high number of leaves in the T₇ treatment could be attributed to increased production of photosynthesizing functional leaves [23], which is aided by a well-balanced media containing sufficient organic matter and clay. This finding is consistent with previous research, which revealed that employing soil mixtures containing organic substrates such as leaf manure [24] and farmyard manure [25,26,27] increased the number of leaves. Organic matter in potting mixtures affects water and nutrient availability [28] and increases seedling production [29], and the compost in our combination may have released nutrients for seedling growth while also boosting water holding capacity. The other findings are consistent with those reported by R.L Bhardwaj [30] in papaya, Abirami *et al.*, [18] in *Myristica fragrans*, and Panchal *et al.*, [10] in Juvekar *et al.*, [31] in Kokum, Sood and Ram [21] in *Oroxylum indicum* and Kaur (2017) in *Mangifera indica*, Khaple *et al.*, [20] in *Grevillea robusta*.

Table 1. Effect of different potting media on germination percentage, survival percentage, plant height, collar diameter and No. of leaves of *Cordia macleodii* in nursery

Treatments	Germination %	Survival %	Plant height (cm)						Collar diameter (mm)						No.of leaves					
			30 D	60 D	90 D	120 D	150 D	180 D	30 D	60 D	90 D	120 D	150 D	180 D	30 D	60 D	90 D	120 D	150 D	180 D
T1	17.1	15.3	3.9	12.7	14.6	23.5	27.8	28.5	1.2	2.2	3.3	4.4	5.4	6.6	1.4	6.6	11.4	16.4	20.4	25.5
T2	31.6	20.6	4.6	12.9	18.3	26.4	31.8	32.8	1.3	2.4	3.5	5.1	5.6	6.8	2.7	8.1	12.8	17.8	22.8	27.6
T3	46.3	23.7	5.5	14.2	20.4	28.4	33.6	37.6	1.4	2.6	3.7	5.7	6.7	7.0	3.7	8.8	13.8	18.8	23.9	28.8
T4	22.7	20.1	4.6	13.4	17.5	25.1	30.9	31.6	1.3	2.4	3.5	4.6	5.6	6.7	2.3	7.6	12.3	17.5	21.8	27.1
T5	20.5	16.7	3.8	12.6	16.4	24.4	29.8	30.3	1.2	2.3	3.4	4.5	5.5	6.6	2.1	7.4	11.8	16.8	21.0	26.7
T6	40.4	22.3	5.0	13.8	19.5	27.7	32.8	33.6	1.4	2.5	3.6	5.7	6.5	6.9	3.2	8.4	13.6	18.4	23.6	28.4
T7	60.1	37.4	6.7	15.5	22.4	30.9	35.5	36.7	1.6	2.7	3.8	5.9	6.9	7.6	4.6	9.5	14.5	19.6	24.8	29.9
T8	52.4	34.6	6.1	14.3	21.6	29.5	34.5	35.4	1.5	2.7	3.8	5.8	6.8	7.1	4.2	9.4	14.3	19.4	24.4	29.4
T9	13.4	15.0	3.5	12.2	13.6	22.3	27.4	28.5	1.1	2.1	3.2	4.4	5.3	6.4	1.5	5.8	10.8	15.5	20.3	25.0
Sem±	1.74	1.78	0.20	0.28	0.20	0.27	0.21	1.05	0.0	0.02	0.02	0.05	0.04	0.09	0.13	0.17	0.14	0.17	0.17	0.17
CD@ 0.05%	5.23	5.35	0.62	0.84	0.62	0.81	0.63	3.16	0.0	0.07	0.07	0.17	0.13	0.29	0.41	0.51	0.43	0.51	0.50	0.51
C.V.(%)	8.94	13.52	7.37	3.59	1.98	1.78	1.16	5.57	2.78	1.68	1.16	1.94	1.319	2.46	8.27	3.72	1.96	1.65	1.30	1.06




Fig. 1. effect of different potting mixture treatments on germination and survival percentage of *Cordia macleodii* in nursery




Fig. 2. effect of different potting mixture treatments on plant height of *Cordia macleodii* in nursery




Fig. 3. Effect of different potting mixture treatments on collar diameter of *Cordia macleodii* in nursery




Fig. 4. Effect of different potting mixture treatments on number of leaves of *Cordia macleodii* in nursery

4. CONCLUSION

The study revealed that the seeds germination and growth of *Cordia macclodii* seedlings in nurseries were considerably influenced by various potting mixture treatments. In terms of the germination and survival percentages of *Cordia macclodii* and seedlings in nurseries, potting mixture treatments T₇ (sand + soil+ vermicompost 1:1:2) showed superior performance. Treatments employing potting mixtures in *Cordia macclodii* seedling growth characteristics have been greatly influenced by T₇ (sand + soil + vermicompost 1:1:2), which has produced maximum plant height, collar diameter, and number of leaves in the nursery. The results of the investigation regarding the growing performance of seedlings lead to the conclusion that a 1:1:2 mixture of potting medium containing sand, soil, and vermicompost encourages the seedlings' quick growth and illustrates that Treatments 7 and 8 have a good impact.

5. FUTURE SCOPE

The findings of this study provide valuable insights into the effect of different potting media on germination and growth of Dahiman (*Cordia macleodii* Hook.) in nursery. However, further research avenues could be explored to enhance our understanding and contribute to sustainable agricultural practices. Some potential future research directions include:

- 1.To achieve faster germination of tree seedlings, other potting mixture concentrations and other tree species propagation materials can be utilized in future experiments.
2. Additionally, this research must to be carried out under various agroclimatic circumstances. In order to derive a sound conclusion and provide a solid foundation for suggesting an appropriate potting media to promote the growth of seedlings, the experiment should be conducted in an open field situation for an extended duration.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

I hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during writing or editing of manuscripts.

ACKNOWLEDGEMENT

This research project was successfully completed thanks to the combined efforts and contributions of numerous people and organizations. We would like to thank Herbal Garden, IGKV, Raipur, Chhattisgarh, for providing the resources and infrastructure needed to carry out this research. We would especially want to thank the Department of Forestry staff and other researchers at the College of Agriculture, IGKV, Raipur, for their collaboration and support during the entire research project.

We would like to express our sincere gratitude to all of the participants and coworkers who gave their valuable cooperation and suggestions during this study. Their participation was essential to gathering relevant data and ensuring the practical significance of our findings.

We would also like to thank our respected mentors, Dr. Pratap Toppo, Dr. S. S. Tuteja, and Prof. Lalji Singh, for their guidance and support. Their experience and outstanding suggestions significantly improved the quality of our research.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Jaenicke H. Potting substrate. What do seedling need to grow well. *Agroforestry today*. 1999;11:31-33.
2. Wolny E, Betekhtin A, Rojek M, Braszewska- Zalewska A, Lusinska J, Hasterok R. Germination and the Early Stages of Seedling Development in *Brachypodium distachyon*. *International Journal of Molecular Sciences*. 2018;19 (10):2916.
3. Doran JC. Australian trees and shrubs: Land rehabilitation and farm tree planting in the tropics. Australian Center of International Agricultural Research, Australia. 1997;ed.2:59-65.
4. Maguire JD. Speed of germination Aid in - selection and evaluation for seedling emergence and vigor. *Crop Sci*. 1962;2: 176-177.
5. Domínguez J. State of the art and new perspectives on vermicomposting research. In: *Earthworm ecology*.

- (Ed.):C.A. Edwards, 2nd edition, CRC Press, Boca Raton. 2004:401-424.
6. Subler S, Edwards CA, Metzger JD. Comparing vermicomposts and composts. *Biocycle*. 1998;39:63–66.
 7. Gopalakrishnan S, Humayun P, Vadlamudi S, Vijayabharathi R, Bhimineni R.K, Rupela, O . Plant growth-promoting traits of streptomycetes with biocontrol potential isolated from herbalvermicompost. *Biocontrol Sci Tech* . 2012;22(10):1199–1210.
 8. Alsina I, Dubova L, Steinberga V, Gmizo G. The effect of vermicompost on the growth of radish. *Acta Horticulturae*. 2013; 1013:359–365.
 9. Gawankar M, Haldankar P, Jamadagni B. Studies on seed germination and seedling growth in Jackfruit (*Artocarpus heterophyllus* Lam.) as influenced by media. *International Journal of Chemical Studies*. 2019;7(5);1699-1705.
 10. Panchal GP, Parasana JS, Patel SR, Patel MV. Effect of different growing Medias and level of IBA on growth and development of Kharni (*Manilkara hexandra* Roxb) seedlings cv. Local. *Global Journal of Bio-Science and Biotechnology*. 2014;3(4):379-383.
 11. Meena AK. Effect of different growing media on seed germination and seedling growth of papaya (*Carica papaya* L.) Cv. Pusa Delicious. A thesis submitted to the Sri Karan Narendra Agriculture University, Jobner, Rajasthan, 2015.
 12. Purwantoro RS. Effect of growing media on seed germination and seedling growth of *Aganope heptaphyllua*. *Nusantara Bioscience*. 2016; 8(2):150-154
 13. Arvind RK. Effect of different growing media and containers on seed germination and seedling growth of papaya (*Carica papaya* L.) cv. Madhubindu. *Trends in Bioscience*. 2014;8(1):231-235.
 14. Ramteke V, Paithankar DH, Ningot EP, Kurrey VK. Effect of GA3 and propagation media on germination, growth and vigour of papaya (*Carica papaya* L.) cv. Coorg Honey Dew. *The Bioscan*. 2015;10(3): 1011-1016.
 15. Mulugeta G. Effect of different potting mixture on the seedling growth and survival of *Albizia gummifera* and *Cordia africana*. *Journal of Natural Sciences Research* . 2014;4(3):25-33.
 16. Arora VK, Singh CB, Sidhu AS, Thind SS .Irrigation, tillage and mulching effects on Soybean yield and water productivity in relation to soil texture. *Agric Water Manag* 2011;98(4):563568
 17. Ghising K, Dey AN, Chakravarty S, Panda MR, Bhowmick N, Mukhopadhyay D. Growth performance of Teak (*Tectona grandis* Linn.) stump under different growing media in nursery. *Biological Forum*. 2022;14(3):322–330.
 18. Abirami K, Rema J, Mathew PA, Srinivasan V, Hamza S. Effect of different propagation media on seed germination, seedling growth and vigour of Nutmeg (*Myristica fragrans* Houtt.). *Journal of Medicinal Plants Research*. 2010;4(19): 2054-2058.
 19. Kumar R , Shamet G, Alam NM, Jana C. Influence of Growing Medium and Seed Size on Germination and Seedling Growth of *Pinus gerardiana* Wall. *Compost Science & Utilization*. 2016;24:98-104.
 20. Khaple AS, Devakumar AS, Maruti Niranjana SP. Influence of Potting Media on *Grevillea robusta* A. Cunn: Seedlings at Nursery Stage. *International Journal of Basic and Applied Sciences*. 2012;1 (4):328-332.
 21. Sood KK, Ram J. Growth and development of seedlings in relation to container size and potting media under nursery conditions in *Oroxylum indicum*- a multipurpose medicinal plant. *Indian Journal of Ecology*. 2019;46:143-148.
 22. Annapurna D, Rathore TS, Joshi G. Effect of container type and size on the growth and quality of Indian Sandalwood (*Santalum album* L.). *Australian Forestry*. 2004;67(2):82-87.
 23. Borah AS, Nath A, Ray AK, Bhat R, Maheswarappa HP, Subramanian P, Dileep MK, Sudhakara VA, Santhoshkumar K, Nazeema K , Ashokan PK. Effect of seed size, rooting medium and fertilizers on the growth of seedlings of silk cotton (*Ceiba pentandra* Linn.). *Indian J. For.* 1994;17 (4):293–300.
 24. Riaz A, Arshad M, Younis A, Raza A, Hameed M. Effect of different growing media on the growth and flowering of *Zinnia elegans* cv. Blue Point. *Pak. J. Bot*. 2008 ;40(4):1579–1585.
 25. Sudhakara K, Mammen W, Santosh AnV, Ashokan PK. Effect of seed size, rooting medium and fertilizers on containerized seedling of silk cotton (*Ceiba pentandra* Linn.). *Indian Forester*. 1995;21(12): 1135–1142.

26. Malewar GU, Adsul PB, Ismail S. Effect of different combinations of fly ash and soil on growth attributes of forest and dryland fruit crops. Indian J. For. 1998;21(2): 124–127.
27. Parasanna JS, Leua HN, Ray NR. Effect of different growing medias mixture on the germination and seedling growth of mango (*Mangifera indica*) cultivars under net house conditions. The Bioscan . 2013;8(3): 897-900.
28. Peter-Onoh CA, Obiefuna JC, Ngwuta A A, Onoh PA, Ibeawuchi II, Ekwugha EU, Emma-Okafor LC, Nze EO, Orji JO, Onyeji EC. Efficacy of five different growth media on seedling emergence and juvenile phenology of *Monodora myristica* (African nutmeg, Ehuru) in the nursery. IOSR J. Agric. Vet. Sci. 2014;7(5):60–63.
29. Baiyeri KP. Evaluation of nursery media for seedling emergence and early seedling growth of two tropical tree species. Moor J. Agric. Resource. 2003;4(1):60–65.
30. Bhardwaj RL. Effect of growing media on seed germination and seedling growth of papaya cv. Red lady. African journal of plant science . 2014;8(4):178-184.
31. Juvekar S, Gawankar M, Malshe K. Effect of growing media on seedling growth of Kokum (*Garcinia indica* Choisy). International Journal of Chemical Studies. 2019;7:2370-2372.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/121735>