



Influence of Different Levels of NAA and 2, 4, 5-T on Fruit Drop, Fruiting, Fruit Retention, Growth and Yield of Indian Ber (*Zizyphus mauritiana* Lamk.)

Manoj Kumar ^{a++*}, A. K. Dwivedi ^{a#}, V. K. Tripathi ^{bt}
and Akash Shukla ^{c#}

^a Department of Fruit Science, C.S. Azad University of Agriculture and Technology, Kanpur, India.

^b Department of Horticulture and Fruit Science, C.S. Azad University of Agriculture and Technology, Kanpur, India.

^c PSS Central Institute of Vocational Education, Bhopal, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i92899

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/98443>

Original Research Article

Received: 04/02/2023

Accepted: 05/04/2023

Published: 11/04/2023

ABSTRACT

The experiment was carried out at Horticulture Garden, Department of Fruit Science, Chandra Shekhar Azad University of Agriculture & Technology, Kanpur (U.P.) during the November 2020-March 2021 and November 2021- March 2022. Sixteen treatments viz. four levels of NAA (0, 20, 30 and 40 ppm) and 2,4,5-T (0, 10, 20 and 30ppm) were studied in a Factorial Completely Randomized Design with three replications. Spraying was done on eleventh November, 2020 in

⁺⁺Research Scholar;

[#]Assistant Professor;

[†]Professor & Head;

*Corresponding author: E-mail: manojgangwarbuat@gmail.com;

first year and fifteenth November, 2021 in second year at fruit setting stage (Pea Stage) with fine nozzle sprayer in each treatment to give uniform spray on all over the treatment of ber plant. Application of (NAA@40 ppm and 2,4,5-T@30 ppm) significantly maximized initial fruit set (162.00 and 163.66), maximum fruit retention (20.39 and 20.43 percent) and maximum initial fruit set percentage (79.60 and 79.56 per cent). The length and width of fruit was significantly (5.35 and 4.99 cm) and (4.24 and 4.27 cm) respectively increased by application same concentration mention above. The maximum (36.62 and 36.82 g) fruit weight and physical properties of fruits like volume (36.57 and 36.42 cc) recorded under treatment (NAA@40ppm and 2,4,5-T@30ppm). The minimum stone length (0.88 and 0.87 cm), minimum stone diameter (0.88 and 0.87 cm), minimum stone weight (0.65 and 0.66 gm) and specific gravity (0.94 and 0.94 g cc⁻¹) significantly found under (NAA@40 ppm and 2,4,5-T@30 ppm). The yield of ber was significantly increased (39.53 and 40.49) kg per plant, yield per hectare (121.30 and 121.33quintal) with treatment combination (NAA@40ppm and 2,4,5-T@30ppm) both the years of experiment.

Keywords: 2,4,5-T, fruit; stone; NAA; length; width; fruiting.

1. INTRODUCTION

Ber has changed scenario of horticulture both in arid and semi-arid regions significantly. At present rainfed ber orchards are seen at all the places across the country. Ber (*Zizyphus mauritiana* Lamk.), a member of family Rhamnaceae, is one of the ancient and common fruits of Indo-China region and has been grown in Indian subcontinent since times immemorial for fresh fruits. In fact it was one of the prominent fruits on which sages in ancient India lived during Vedic ages. Ber is known to be indigenous to the area stretching from India to South western China and Malaya [1]. The genus *Zizyphus* consists of over 100 species, out of which 18-40 species are found to be grown in India. Ber is evergreen shrub or small tree up to 15 m high, with trunk 40 cm or more in diameter, spreading crown, stipular spines and many drooping branches. The tree is associated with Lord Shiva, whose worship is considered incomplete without offering of jujube fruit, especially during Mahashivaratri.

India ranks first among the ber growing countries of the world with an area of 50,000 ha and annual production of 5.13 lakh MT (NHB Database, 2018-19).

Ber tree bears its inflorescence in the axil of leaves on current season's growth. The flowering period lasts for about two and a half months from Sept to Nov. The fruit setting starts in second week of Oct and continues up to first fortnight of Nov. The fruits reach maturity in about 180 days after fruit setting. The fruit growth in terms of length and diameter follows a 'double sigmoid' curve and traces of malic acid, oxalic acid and quercetin. That's why ber is referred to as 'the apple of arid zone'.

The ber fruit is richer than apple in protein, phosphorus, calcium and Vitamin 'C' [2] and one hundred gram of edible ber fruit contains moisture (85.9%), protein (0.8g), fat (0.1g), carbohydrate (12.88%), calcium (0.03g), phosphorus (0.03g), iron (0.8g), carotene (70 IU) and vitamin 'C' (50-100mg). Although the research work on these aspects is being carried out by various research workers throughout India yet the detailed information on the effects of growth regulator sprays on fruit drop, fruiting, fruit retention and growth of ber is lacking and there is still no recommendation keeping the above facts in view, the present study has been planned.

2. MATERIALS AND METHODS

2.1 Experimental Site

The present experiment was conducted at the Horticulture Garden, Department of Fruit Science, College of Horticulture, of Chandra Shekhar Azad University of Agriculture and Technology, Kanpur (U.P.) during November 2020- March 2021 and November 2021- March 2022. Geographically Kanpur is situated in the Gangetic alluvial belt of central U. P. its lies in altitude and longitude range between 25.26° to 26.58° north and 79.31° to 84.34° east and mean elevation of 125.90 m above the sea level.

2.2 Soil and Climate Condition

Kanpur is characterized by semi and sub-tropical climate with hot dry summer and cold winters. The annual rainfall is about 80-85 cm. the major portion of rain received between July to September, with scattered shower in winter, from the North-East monsoon. The pH of experimental field was determined by electric pH meter as described by Piper (1966) while organic carbon

Table 1. Detail of the treatment combinations

Symbols Used	Treatment combination
T ₁	N ₀ A ₀ (NAA – 0 ppm+2,4,5-T – 0 ppm)
T ₂	N ₀ A ₁ (NAA – 0 ppm+2,4,5-T – 10 ppm)
T ₃	N ₀ A ₂ (NAA – 0 ppm+2,4,5-T – 20 ppm)
T ₄	N ₀ A ₃ (NAA – 0 ppm+2,4,5-T – 30 ppm)
T ₅	N ₁ A ₀ (NAA – 20 ppm+2,4,5-T – 0 ppm)
T ₆	N ₁ A ₁ (NAA – 20 ppm+2,4,5-T – 10 ppm)
T ₇	N ₁ A ₂ (NAA – 20 ppm+2,4,5-T – 20 ppm)
T ₈	N ₁ A ₃ (NAA – 20 ppm+2,4,5-T – 30 ppm)
T ₉	N ₂ A ₀ (NAA – 30 ppm+2,4,5-T – 0 ppm)
T ₁₀	N ₂ A ₁ (NAA – 30 ppm+2,4,5-T – 10 ppm)
T ₁₁	N ₂ A ₂ (NAA – 30 ppm+2,4,5-T – 20 ppm)
T ₁₂	N ₂ A ₃ (NAA – 30 ppm+2,4,5-T – 30 ppm)
T ₁₃	N ₃ A ₀ (NAA – 40 ppm+2,4,5-T – 0 ppm)
T ₁₄	N ₃ A ₁ (NAA – 40 ppm+2,4,5-T – 10 ppm)
T ₁₅	N ₃ A ₂ (NAA – 40 ppm+2,4,5-T – 20 ppm)
T ₁₆	N ₃ A ₃ (NAA – 40 ppm+2,4,5-T – 30 ppm)

was determined by Walkley and Black (1967) rapid titration method. The available Nitrogen was determined by alkaline permagnate method as reported by Piper (1966), and available phosphorus and potash by Olsen's method Olsen et al. [3] and flame photometer Meston [4], respectively. The E.C. was determined by Conductivity Bridge as described Jackson [5]. Sulphur determination was done by method described by Richard [6] and available zinc was determined as per the method suggested by Lindsey and Norvell [7].

2.3 Detail of Treatments and Design

The experiment comprised 16 treatments consisting of foliar spray of Naphthalene Acetic Acid (NAA) and 2,4,5-T (A). The following treatments were compared. Experiment was laid out in Factorial Randomized Block Design with three replications.

2.3.1 Preparation of solution

For preparation of NAA stock solution one gram NAA was dissolved in appropriate alcohol and adding water that was converted in one liter. Thus, stock solution was prepared. For obtaining 20 ppm solution of NAA 20 ml solution of NAA was taken out from stock solution and with adding water, 1000 ml solution was prepared. Thus, it was obtained 20 ppm NAA solution. Further 40 ppm NAA solution was prepared as similar method. Similarly, stock solution of (2,4,5-T) was made and from this stock solution 10 ppm, 20ppm and 30 ppm (2,4,5-T) solution were prepared as per above method.

2.3.2 Methods of application

Above solutions with different concentration were sprayed by foot sprayer in the morning hours and selected branches were fully drenched. For control there was only water spray is allowed.

2.3.3 Time of application

Spraying was done on 11th November, 2020 in first year and 15th November, 2021 in second year at fruit setting stage (Pea Stage) with fine nozzle sprayer in each treatment to give uniform spray on all over the treatment of ber plant.

2.4 Statistical Analysis

Data were analyzed according to the method described by Panse and Sukhantme [8] and S.R.S. Chandel [9].

3. RESULTS AND DISCUSSION

3.1 Initial Fruit Set

Regarding different NAA and 2,4,5-T concentrations on initial fruit set of fruit are an expression of fruiting parameters of the plants which was influenced by NAA and 2,4,5-T growth regulators over control. It is obviously appeared with vision of the data that all the concentration of NAA significantly influenced to fruit sets in ber trees. Interactive effect of NAA and 2,4,5-T also influenced significantly on initial fruit set in ber but interactive treatment gave further improvement in fruit set was over main effect. The maximum of (162.00 and 163.66) initial fruit

set was noted under N_3A_3 treatment (NAA@40 ppm and 2,4,5-T@30 ppm) treatment over interactive control i.e. N_0A_0 recorded (151.00 and 152.00) initial fruit set during both the years of experiments. The effect of NAA on plant growth is greatly dependent on the time of admission and concentration. NAA has been shown greatly increased in plant by exogenous application. Due to these causes fruit setting was enhanced in present investigation. These findings are in accordance with the reports of Sandhu and Thind [10], Chaurasiya et al., [11], Chaudhary et al., [12] Das et al., [13] in ber, Badal and Tripathi [14] in guava, Saraswat et al., [15] in litchi. The 2,4,5-T probably might be due to providing of right concentration of 2,4,5-T during investigation causing enhancement of vegetative growth of the plants hastening the production of more photosynthesis towards the fruit bearing area which contributed to increase fruit set in plant. These findings are collaborated with the reports of Pandey [16] in ber, Bhat et al., [17] in litchi, Maurya et al., [18] in mango, Kumar et al., [19], Singh and Sharma [20] in ber.

3.2 Fruit Retention (%)

The fruit retention was influenced by various treatments of NAA and 2,4,5-T in present investigation. The combination effect of NAA and 2,4,5-T induced significant variation on fruit retention and its interactive treatment of N_3A_3 significantly maximized (20.39 and 20.43 percent) fruit retention closely followed by N_2A_3 (18.90 and 19.67 percent) over control i.e. N_0A_0 of (9.61 and 9.67 percent) during both the years of experiments. The exogenous application of NAA might have acted to prevention of abscission layer and thus, retention of fruit is increased. These findings are in line with the reports of Bankar and Prasad [21], Ghosh et al., [22], Singh and Ball [23] in ber, Chauhan et al., [24] in litchi, Deepa Lal et al., [25] in kinnow, Tiwari et al., [26] in aonla, Badal and Tripathi [27] in guava, Tripathi and Vivekanand [28] in aonla, Saraswat et al., [15] in litchi. The increase in fruit retention might be due to effective of different chemicals as well as 2, 4, 5-T on metabolic activity of the plant and improved source sink relationship with favorably influenced the metabolic status resulting in better check of fruit drop and enhancing retention of the more number of fruits on the plants. The findings are in agreement with the reports of Pandey [16] in ber, Bhat et al. [17] in litchi, Maurya et al. [18] in mango, Kumar et al. [19], Singh and Sharma [20] in ber.

3.3 Fruit Drop (%)

The fruit drop in ber fruit was significantly influenced with the sprays of NAA and 2,4,5-T treatments in ber fruits. The NAA and 2,4,5-T brought about significant treatment variation on fruit drop and of N_3A_3 expressed significantly minimum of (79.60 and 79.56 percent) fruit drop closely followed by N_2A_3 (81.09 and 80.32 percent). Significantly maximum of (82.16 and 81.43 percent) fruit drop was exhibited under control (N_0A_0) during both the years of experiments. The application of NAA might have increased the concentration of auxin in plants which possibly induced to reduction of fruit drop. These findings are in line with the reports of Pandey et al., [29], Naseem et al., [30], Chaudhary et al., [12] in ber, Haidry et al., [31] in mango, Saraswat et al., [15] in litchi. The induction in fruit drop was 2,4,5-T sprays possibly increased auxin synthesis which may cause to prevent fruit drop. These findings are collaborated with the reports of Deepa Lal et al., [25] in kinnow, Tiwari et al., [26] in aonla, Badal and Tripathi [27] in guava, Tripathi and Vivekanand [28] in aonla, Pandey et al., [29] in ber, Bhat et al., [17] in litchi, Maurya et al., [18] in mango, Kumar et al., [19], Singh and Sharma [20] in ber, Deepa Lal et al., [32] in kinnow.

3.4 Length of Fruit (cm)

The interactive effect of NAA and 2,4,5-T did differ significantly but further improvement was observed over mean values and combined treatment of N_3A_3 (NAA @40 ppm and 2,4,5-T@30 ppm) recorded maximum of (5.35 and 4.99 cm) length against the minimum of (2.62 and 2.69 cm) fruit length was expressed under control (N_0A_0) during both the years of experiment. Enhancement range on length of fruit was fruits indicated caused by NAA treatment might be due to its involvement in cell division, cell elongation and decreased volume of intracellular space in the monocarpic cells which could have boosted plant health there by producing healthy and larger fruit NAA increase the growth rate of fruit which results a bigger fruit size ultimately. These findings are in line with reports of Tiwari et al., [26] in aonla, Badal and Tripathi [27] in guava, Tripathi and Vivekanand [28] in aonla, Meena et al., [33], Arora et al., [34], Pandey et al., [16] in ber, Rathod et al., [35] in aonla, Patil et al., [36], Kumar et al., [37] in mango. The increase in size of fruits with application of 2,4,5-T might be due to significantly increase in cell division and cell

elongation also associated with active performance of Photosynthesis in the plant and photosynthetes were translocated to the fruits which caused possibly to increase in fruit size. These findings are collaborated with the reports of Randhawa et al., [38] in sweet lime, Tripathi et al., [39], Brahmachari et al. [40] in guava, Suman et al., [41] in sweet lime.

3.5 Width of Fruit (cm)

The fruit width of ber was significantly induced with NAA and 2,4,5-T combination was found to be significant. Combined treatment of N_3A_3 (NAA@40 ppm and 2,4,5-T@30ppm) induced significantly maximum (4.24 and 4.27 cm) width of fruit while the minimum (2.63 and 2.66 cm) width of fruit was presented with control (N_0A_0) during both years of experiments. The diameter of fruit was enhanced due to sprays of NAA treatments might be due to its involvement in cell division, cell elongation which ultimately induced to width of fruits. These findings are in agreement with the reports of Tiwari et al., [26] in aonla, Tripathi and Vivekanand [28] in aonla, Singh and Singh [42], Pandey et al., [16], Singh et al., [43], Arora and Singh [34] in ber and Rathod et al., [35] in aonla, Kumar et al., [37] in mango. The different concentration of 2,4,5-T might be due to significantly increased cell division and cell elongation. This result may have associated with active performance of photosynthesis in the plant and they were translocated to the fruits which caused to increase in fruit size. These finding are in line with reports of Randhawa et al., [38] in sweet lime, Tripathi et al., [39], Brahmachari et al., [40] in guava, Suman et al. [41] in sweet lime.

3.6 Fruit Weight (gm)

The effect of foliar sprays of NAA and 2,4,5-T positively influenced on fruit weight of ber. The significant variation was observed in interactive treatments of NAA and 2,4,5-T its N_3A_3 (NAA@40ppm and 2,4,5-T@30ppm) treatment induced to the maximum of (36.62 and 36.82 g) fruit weight closely followed by N_2A_3 (NAA@30ppm and 2,4,5-T@30ppm) expressed (30.56 and 31.01 g) fruit weight. Significantly poorest (25.55 and 25.55 g) fruit weight was recorded under control (N_0A_0) during both the years of experiment. The growth regulator NAA might have improved the synthesis of more photosynthetes and their translocation to the fruits which may have increased the weight of fruits in present investigation. These result in

conformity with those of Tiwari et al., [26] in aonla, Badal and Tripathi [27] in guava, Tripathi and Vivekanand [28] in aonla, Bal et al., [44], Bal et al., [45], Singh et al., [43] in ber, Haidry et al., (2001), Singh et al., [46] in mango, Kumar et al., [37] in mango. The improvement in fruit weight probably 2,4,5-T enhanced deposition of solids which increased in cell size by increasing the accumulation of water in intracellular space which might be enhanced to fruit weight. These findings collaborated with the reports of Randhawa et al., [38] in sweet lime, Tripathi et al., [39], Brahmachari et al. [40] in guava, Suman et al., [41] in sweet lime.

3.7 Fruit Volume (cc)

The volume of ber fruit was consistently influenced with sprays of NAA and 2,4,5-T treatments. In this regard (36.57 and 36.42 cc) fruit volume was expressed under the interactive treatment of N_3A_3 (NAA@40ppm and 2,4,5-T@30ppm) closely followed by treatment N_2A_3 (NAA@40ppm and 2,4,5-T@30ppm) recorded (35.10 and 35.14 cc) fruit volume. Significantly the lesser (25.49 and 25.59 cc) volume was recorded under control (N_0A_0) during both the years of experiments. It might have also being due to cell division and cell expansion, increased volume of intracellular space in the mesocarpic cell and increase the water absorption with mobilization of sugar and minerals in the expended cell and intracellular space of mesocarp which improve size and volume of the fruit. These findings collaborated with the reports of Badal and Tripathi [14] in guava, Patil et al., [36] in mango. The volume of fruit was greatly influenced with the application of 2,4,5-T treatments which was possibly might be due to 2,4,5-T which regulates semi permeability of cell wall and mobilization of water extended into fruit which ultimately help to enhancement of fruit volume. These findings are in line with the reports of Bal et al., [47] in ber.

3.8 Stone Length (cm)

The length of stone was significantly enhanced was observed over mean values and combined treatment of N_3A_3 (NAA @ 40 ppm & 2,4,5-T @ 30 ppm) recorded minimum of (0.88 and 0.87 cm) cm length against the maximum of (2.70 and 2.79 cm) stone length was expressed under control (N_0A_0) during both the year of experiments. The superiority on length of fruits

indicated caused by NAA treatment might be due to its involvement in cell division, cell elongation and decreased volume of intracellular space in the monocarpic cells which could have boosted plant health there by producing healthy and larger fruit NAA increase the growth rate of fruit which results a bigger fruit size ultimately small size of stone. These findings are in line with reports of Meena et al., [33], Arora et al., [34], Pandey [16] in ber, Rathod et al., [35] in aonla, Patil et al., [36] in mango. The increase in size of fruits with application of 2,4,5-T might be due to significantly increase in cell division and cell elongation associated with active performance of Photosynthesis in the plant and photosynthetes were translocated to the fruits which caused possibly to increase in stone size. These findings are collaborated with the reports of Bal et al., [47], Tripathi et al., [39], Pandey [16], Ram et al.,[48] in ber, Kumar et al., [49] in guava.

3.9 Stone Diameter (cm)

The diameter of stone of ber was significantly induced combined treatment of N_3A_3 (NAA @ 40 ppm & 2,4,5-T @ 30 ppm) induced significantly minimum (0.73 and 0.70cm) diameter of stone closely followed by treatment N_2A_3 (0.75 and 0.73cm). These findings are in agreement with the reports of Singh and Singh [42], Pandey [16], Singh et al., [43], Arora et al., [43] in ber, Rathod et al., [35] in aonla. This result may have associated with active performance of photosynthesis in the plant and they were translocated to the stone which caused to increase in stone size Bal et al., [47], Tripathi et al., [39], Pandey [16], Ram et al., [48] in ber, Kumar et al., [49] in guava.

3.10 Stone Weight (gm)

The effect of foliar sprays of NAA and 2,4,5-T positively influenced on stone weight of ber with interactive treatments of NAA and 2,4,5-T its (NAA @ 40 ppm and 2,4,5-T @ 30 ppm) N_3A_3 treatment induced to the minimum of (0.65 and 0.66 gm) stone weight closely followed by N_2A_3 (NAA@20ppm and 2,4,5-T@30ppm) expressed (0.77 and 0.78gm) stone weight. Significantly maximum (1.80 and 1.77 gm) stone weight was recorded under control (N_0A_0) during both the years of experiments. Probably NAA enhanced deposition of solids which increased in cell size by increasing the accumulation of water in intracellular space which might be reduced to stone weight. These findings are gets support to

the reports of Bal et al., [44], Haidry et al., (2001), Singh et al., [46], Banker and Prasad [21] in ber, Saraswat et al., [15] in litchi. The 2,4,5-T result may have due to associated with active performance of photosynthesis in the plant and they were translocated to the stone which caused to reduce in stone weight Bal et al., [47], Tripathi et al., [39], Pandey [16], Ram et al.,[48] in ber, Kumar et al., [49] in guava.

3.11 Specific Gravity ($g\ cc^{-1}$)

The specific gravity of ber fruit was influenced positively with the sprays of NAA and 2,4,5-T treatments. The significant variation was observed in interactive treatments of NAA and 2,4,5-T its (NAA @ 40 ppm and 2,4,5-T @ 30 ppm) N_3A_3 treatment induced to the minimum of (0.94 and 0.94 $g\ cc^{-1}$) specific gravity closely followed by N_2A_3 (NAA@20ppm and 2,4,5-T@30ppm) expressed (0.95 and 0.97 $g\ cc^{-1}$) specific gravity. Significantly maximum (1.32 and 1.32 $g\ cc^{-1}$) specific gravity was recorded under control (N_0A_0) during both the years of experiments. These findings get support of Tiwari et al., [26], Tripathi and Vivekanand [28] in aonla, Ghosh et al., (2013) in ber. The concentration of 2,4,5-T also influenced on specific gravity These findings are in line with the reports of Tripathi et al., [26], Pandey [16], Ram et al., [48] in ber.

3.12 Yield (Kg/Plant)

The influences were shown with sprays of NAA and 2,4,5-T growth regulators. The NAA and 2,4,5-T also influenced significantly on yield per plant in ber fruits. The maximum of (39.53 and 40.49 kg) was revealed under the interactive treatment of N_3A_3 followed by N_2A_3 (38.95 and 38.97kg) over control (N_0A_0) during both the years of experiments. The improvement brought about NAA may be attributed to its physiological activities in the plants, which could have checked fruit drop and minimized number of blemished fruits considerably thereby increasing yield. These findings are collaborated with the reports of Tiwari et al., [26] in aonla, Badal and Tripathi (2021b) in guava, Tripathi and Vivekanand [28] in aonla, Saraswat et al. [15], in litchi, Kumar et al., [37] in mango. The increase in yield by growth regulator 2,4,5-T associated with high rate of enzymatic activities as well as involvement of biosynthesis of auxin, increase in number of size of fruit which ultimately enhanced the yield. These findings are in line with the reports of Tripathi et al., [39], Kumar et al., [19] in ber, Kumar and Tripathi [50] in strawberry.

Table 2. Effect of foliar application of NAA, 2, 4, 5-T and their interaction on initial fruit set and fruit retention (%) in ber

Parameter	PGRs Doses 2,4,5-T ppm (A)	NAA ppm (N)									
		2021					2022				
		N ₀ Control	N ₁ 20	N ₂ 30	N ₃ 40	Mean B	N ₀ Control	N ₁ 20	N ₂ 30	N ₃ 40	Mean B
Initial Fruit Set	A ₀ Control	151.00	153.33	155.00	156.66	154.00	152.00	154.33	156.00	157.66	155.00
	A ₁ 10	157.33	159.00	159.66	159.33	158.83	158.66	160.33	161.00	159.66	159.91
	A ₂ 20	156.66	155.66	156.66	155.66	156.16	157.66	157.00	158.33	157.66	157.66
	A ₃ 30	156.66	159.00	160.00	162.00	159.41	158.66	161.00	162.00	163.66	161.33
	Mean A	155.41	156.75	157.83	158.41		156.75	158.16	159.33	159.66	
	Factors	A	B	AXB			A	B	AXB		
	SE(m)±	0.44	0.44	0.88			0.65	0.65	1.30		
	C.D.	1.27	1.27	2.55			1.88	1.88	3.76		
	SE(d)	0.62	0.62	1.24			0.92	0.92	1.84		
Fruit Retention (%)	A ₀ Control	9.61	10.31	10.86	11.22	10.50	9.67	10.39	10.90	11.26	10.55
	A ₁ 10	11.59	12.12	12.59	13.17	12.36	11.63	12.16	12.67	13.27	12.43
	A ₂ 20	13.66	14.10	14.77	15.42	14.48	13.74	14.12	14.83	15.49	14.54
	A ₃ 30	16.32	17.83	18.90	20.39	18.36	16.39	18.57	19.67	20.43	18.76
	Mean A	12.79	13.59	14.28	15.05		12.85	13.81	14.52	15.11	
	Factors	A	B	AXB			A	B	AXB		
	SE(m)±	0.10	0.10	0.20			0.12	0.12	0.24		
	C.D.	0.29	0.29	0.59			0.36	0.36	0.72		
	SE(d)	0.14	0.14	0.29			0.17	0.17	0.35		

Table 3. Effect of foliar application of NAA, 2, 4, 5-T and their interaction on fruit drop (%) and length (cm) of ber fruit

Parameter	PGRs doses 2,4,5-T ppm (A)	NAA ppm (N)									
		2021					2022				
		N ₀ Control	N ₁ 20	N ₂ 30	N ₃ 40	Mean B	N ₀ Control	N ₁ 20	N ₂ 30	N ₃ 40	Mean B
Fruit drop (%)	A ₀ Control	90.39	89.68	89.13	88.77	89.49	90.33	89.60	89.09	88.73	89.44
	A ₁ 10	88.41	87.88	87.40	86.82	87.63	88.36	87.83	87.33	86.73	87.56
	A ₂ 20	86.33	85.90	85.23	84.57	85.51	86.26	85.87	85.17	84.50	85.45
	A ₃ 30	83.67	82.16	81.09	79.60	81.63	83.61	81.43	80.32	79.56	81.23
	Mean A	87.20	86.40	85.71	84.94		87.14	86.18	85.48	84.88	
	Factors	A	B	AXB			A	B	AXB		
	SE(m)±										

Parameter	PGRs doses 2,4,5-T ppm (A)	NAA ppm (N)									
		2021					2022				
		N ₀ Control	N ₁ 20	N ₂ 30	N ₃ 40	Mean B	N ₀ Control	N ₁ 20	N ₂ 30	N ₃ 40	Mean B
	SE(m)±	0.10	0.10	0.20			0.12	0.12	0.24		
	C.D.	0.29	0.29	0.59			0.36	0.36	0.72		
	SE(d)	0.14	0.14	0.29			0.17	0.17	0.35		
Length of ber fruit (cm)	A ₀ Control	2.62	2.71	2.78	2.92	2.75	2.69	2.81	2.92	3.01	2.86
	A ₁ 10	3.00	3.19	3.33	3.43	3.24	3.10	3.26	3.37	3.45	3.29
	A ₂ 20	3.53	3.71	3.81	3.94	3.75	3.64	3.84	3.92	4.02	3.85
	A ₃ 30	4.03	4.41	5.01	5.35	4.70	4.19	4.29	4.71	4.99	4.55
	Mean A	3.29	3.50	3.73	3.91		3.41	3.55	3.73	3.87	
	Factors	A	B	AXB			A	B	AXB		
	SE(m)±	0.05	0.05	0.10			0.03	0.03	0.07		
	C.D.	0.14	0.14	0.29			0.11	0.11	0.22		
	SE(d)	0.07	0.07	0.14			0.05	0.05	0.10		

Table 4. Effect of foliar application of NAA, 2, 4, 5-T and their interaction on width of fruit (cm) and fruit weight in ber (g)

Parameter	PGRs Doses 2,4,5-T ppm (A)	NAA ppm (N)									
		2021					2022				
		N ₀ Control	N ₁ 20	N ₂ 30	N ₃ 40	Mean B	N ₀ Control	N ₁ 20	N ₂ 30	N ₃ 40	Mean B
Width of fruit (cm)	A ₀ Control	2.63	2.73	2.78	2.90	2.76	2.66	2.78	2.81	2.88	2.78
	A ₁ 10	3.00	3.13	3.22	3.30	3.16	3.07	3.15	3.25	3.32	3.20
	A ₂ 20	3.43	3.52	3.62	3.72	3.57	3.45	3.57	3.66	3.77	3.61
	A ₃ 30	3.83	4.00	4.12	4.24	4.05	3.89	4.05	4.15	4.27	4.09
	Mean A	3.22	3.34	3.43	3.54		3.27	3.39	3.47	3.56	
	Factors	A	B	AXB			A	B	AXB		
	SE(m)±	0.00	0.00	0.00			0.00	0.00	0.01		
	C.D.	0.00	0.00	0.01			0.02	0.02	0.05		
SE(d)	0.00	0.00	0.00			0.01	0.01	0.02			
Fruit weight (g)	A ₀ Control	25.55	25.91	26.42	26.99	26.22	25.55	26.46	26.70	27.66	26.59
	A ₁ 10	27.55	27.93	28.54	29.14	28.29	27.97	28.25	29.47	29.78	28.86
	A ₂ 20	30.16	31.27	32.13	32.96	31.63	30.53	31.46	31.99	33.22	31.80
	A ₃ 30	33.65	34.12	35.16	36.62	34.89	34.86	35.14	35.90	36.82	35.68
	Mean A	29.23	29.81	30.56	31.43		29.73	30.33	31.01	31.87	

Parameter	PGRs Doses		NAA ppm (N)							
	2,4,5-T ppm		2021				2022			
	(A)	N ₀ Control	N ₁ 20	N ₂ 30	N ₃ 40	Mean B	N ₀ Control	N ₁ 20	N ₂ 30	N ₃ 40
Factors	A	B	AXB			A	B	AXB		
SE(m)±	0.01	0.01	0.02			0.02	0.02	0.04		
C.D.	0.03	0.03	0.06			0.12	0.12	0.04		
SE(d)	0.01	0.01	0.03			0.06	0.06	0.08		

Table 5. Effect of foliar application of NAA, 2, 4, 5-T and their interaction on fruit volume (cc) and stone length in ber (cm)

Parameter	PGRs Doses		NAA ppm (N)								
	2,4,5-T ppm		2021				2022				
	(A)	N ₀ Control	N ₁ 20	N ₂ 30	N ₃ 40	Mean B	N ₀ Control	N ₁ 20	N ₂ 30	N ₃ 40	Mean B
Volume (cc)	A ₀ Control	25.49	25.87	26.33	26.86	26.13	25.59	25.90	26.38	26.90	26.19
	A ₁ 10	27.37	27.89	28.49	29.10	28.21	27.47	27.93	28.53	29.08	28.25
	A ₂ 20	30.10	31.20	32.10	32.84	31.56	30.11	31.22	32.06	32.89	31.57
	A ₃ 30	33.58	34.10	35.10	36.57	34.83	33.62	34.08	35.14	36.42	34.81
	Mean A	29.14	29.77	30.51	31.34		29.19	29.78	30.52	31.32	
	Factors	A	B	AXB			A	B	AXB		
	SE(m)±	0.02	0.02	0.04			0.02	0.02	0.05		
	C.D.	0.06	0.06	0.12			0.08	0.08	0.16		
SE(d)	0.03	0.03	0.06			0.03	0.03	0.07			
Stone length (cm)	A ₀ Control	2.70	2.62	2.52	2.32	2.54	2.79	2.66	2.55	2.38	2.59
	A ₁ 10	2.20	2.09	1.99	1.86	2.03	2.20	2.06	1.99	1.90	2.04
	A ₂ 20	1.73	1.59	1.48	1.34	1.53	1.77	1.59	1.42	1.34	1.53
	A ₃ 30	1.23	1.13	0.97	0.88	1.05	1.21	1.12	1.02	0.87	1.05
	Mean A	1.96	1.86	1.74	1.60		1.99	1.86	1.75	1.62	
	Factors	A	B	AXB			A	B	AXB		
	SE(m)±	0.00	0.00	0.01			0.00	0.00	0.01		
	C.D.	0.01	0.01	0.03			0.02	0.02	0.04		
SE(d)	0.00	0.00	0.01			0.01	0.01	0.02			

Table 6. Effect of foliar application of NAA, 2, 4, 5-T and their interaction on stone diameter (cm) and stone weight in ber (g)

Parameter	PGRs doses 2,4,5-T ppm (A)	NAA ppm (N)									
		2021					2022				
		N ₀ Control	N ₁ 20	N ₂ 30	N ₃ 40	Mean B	N ₀ Control	N ₁ 20	N ₂ 30	N ₃ 40	Mean B
Stone diameter (cm)	A ₀ Control	1.31	1.26	1.20	1.15	1.23	1.29	1.25	1.19	1.14	1.22
	A ₁ 10	1.09	1.05	1.02	0.97	1.03	1.08	1.04	1.02	0.98	1.03
	A ₂ 20	0.96	0.95	0.93	0.88	0.93	0.95	0.93	0.91	0.87	0.91
	A ₃ 30	0.85	0.79	0.75	0.73	0.78	0.82	0.79	0.73	0.70	0.76
	Mean A	1.05	1.01	0.97	0.93		1.03	1.00	0.96	0.92	
	Factors	A	B	AXB			A	B	AXB		
	SE(m)±	0.00	0.00	0.01			0.00	0.00	0.00		
	C.D.	0.01	0.01	0.02			0.01	0.01	0.02		
SE(d)	0.00	0.00	0.01			0.00	0.00	0.01			
Stone weight (gm)	A ₀ Control	1.80	1.72	1.65	1.56	1.68	1.77	1.71	1.63	1.55	1.67
	A ₁ 10	1.50	1.42	1.34	1.26	1.38	1.48	1.41	1.33	1.25	1.37
	A ₂ 20	1.20	1.09	1.04	0.98	1.08	1.18	1.08	1.03	0.94	1.06
	A ₃ 30	0.88	0.82	0.77	0.65	0.78	0.91	0.83	0.77	0.66	0.79
	Mean A	1.34	1.26	1.20	1.11		1.33	1.26	1.19	1.10	
	Factors	A	B	AXB			A	B	AXB		
	SE(m)±	0.00	0.00	0.00			0.00	0.00	0.00		
	C.D.	0.01	0.01	0.02			0.00	0.00	0.01		
SE(d)	0.00	0.00	0.01			0.00	0.00	0.00			

Table 7. Effect of foliar application of NAA, 2, 4, 5-T and their interaction on specific gravity (g /cc) and yield (kg/plant)in ber

Parameter	PGRs Doses 2,4,5-T ppm (A)	NAA ppm (N)									
		2021					2022				
		N ₀ Control	N ₁ 20	N ₂ 30	N ₃ 40	Mean B	N ₀ Control	N ₁ 20	N ₂ 30	N ₃ 40	Mean B
Specific gravity (g /cc)	A ₀ Control	1.32	1.16	1.10	1.08	1.16	1.32	1.17	1.11	1.09	1.17
	A ₁ 10	1.05	1.04	1.05	1.03	1.04	1.06	1.04	1.06	1.05	1.05
	A ₂ 20	1.02	1.00	0.99	0.98	1.00	1.03	1.00	0.99	0.98	1.00
	A ₃ 30	0.97	0.98	0.95	0.94	0.96	0.97	0.95	0.97	0.94	0.96
	Mean A	1.09	1.04	1.02	1.01		1.09	1.04	1.03	1.01	
	Factors	A	B	AXB			A	B	AXB		

Parameter	PGRs Doses 2,4,5-T ppm (A)	NAA ppm (N)										
		2021					2022					
		N ₀ Control	N ₁ 20	N ₂ 30	N ₃ 40	Mean B	N ₀ Control	N ₁ 20	N ₂ 30	N ₃ 40	Mean B	
	SE(m)±	0.01	0.01	0.02			0.01	0.01	0.02			
	C.D.	0.03	0.03	0.07			0.03	0.03	0.07			
	SE(d)	0.01	0.01	0.03			0.01	0.01	0.03			
Yield (kg/plant)	A ₀ Control	19.92	19.71	19.80	20.42	19.96	19.95	19.73	19.82	20.45	19.99	
	A ₁ 10	20.90	21.14	22.35	23.64	22.01	20.89	21.15	22.36	23.66	22.01	
	A ₂ 20	24.84	25.47	28.15	31.14	27.40	24.87	26.16	28.17	31.17	27.59	
	A ₃ 30	34.23	36.14	38.95	39.53	37.21	34.28	36.17	38.97	40.49	37.47	
	Mean A	24.97	25.61	27.31	28.68		25.00	25.80	27.33	28.94		
	Factors	A	B	AXB			A	B	AXB			
		SE(m)±	0.14	0.14	0.29			0.11	0.11	0.23		
		C.D.	0.42	0.42	0.85			0.33	0.33	0.67		
		SE(d)	0.20	0.20	0.41			0.16	0.16	0.32		

Table 8. Effect of foliar application of NAA, 2, 4, 5-T and their interaction on yield in ber (q/ha)

PGRs Doses 2,4,5-T ppm (A)	NAA ppm (N)									
	2021					2022				
	N ₀ Control	N ₁ 20	N ₂ 30	N ₃ 40	Mean B	N ₀ Control	N ₁ 20	N ₂ 30	N ₃ 40	Mean B
A ₀ Control	101.40	102.47	103.72	105.68	103.31	101.48	102.55	103.98	105.78	103.45
A ₁ 10	106.76	108.34	109.38	110.96	108.86	106.83	108.52	109.58	110.57	108.87
A ₂ 20	112.13	113.36	114.78	115.91	114.04	111.54	112.43	113.79	114.90	113.17
A ₃ 30	116.72	118.10	119.61	121.30	118.93	116.08	116.80	117.87	121.33	118.02
Mean A	109.25	110.57	111.87	113.46		108.98	110.07	111.31	113.15	
Factors	A	B	AXB			A	B	AXB		
	SE(m)±	0.05	0.05	0.10			0.13	0.13	0.26	
	C.D.	0.14	0.14	0.29			0.37	0.37	0.75	
	SE(d)	0.07	0.07	0.14			0.18	0.18	0.36	

3.13 Yield (q/hac)

The interactive effect of NAA and 2,4,5-T also influenced significantly on quintal per plants in ber fruits. The maximum of (121.30 and 121.33quintal) was revealed under the interactive treatment of N_3A_3 (NAA@40ppm and 2,4,5-T@30ppm) followed by (119.61 and 117.87 quintal) treatment N_2A_3 of (NAA@30ppm and 2,4,5-T@30ppm) over control (N_0A_0) (101.40 and 101.48 quintal) during both the years of experiments. The improvement brought about NAA may be attributed to its physiological activities in the plants, which could have checked fruit drop and minimized number of blemished fruits and yield kg per plants considerably thereby increasing yield quintal per plants. These findings are collaborated with the reports of Ghosh et al., (2013) in ber, Kumar and Tripathi [50] in strawberry. Increase in yield by growth regulator 2,4,5-T associated with high rate of enzymatic activities as well as involvement of biosynthesis of auxin, increase in number of size of fruit which ultimately enhanced the yield. These findings are in line with the reports of Tripathi et al., [39], Kumar et al., [19] in ber, Kumar and Tripathi [50] in strawberry [51-53].

4. CONCLUSION

It may be concluded that the application of NAA and 2,4,5-T resulted in to flowering fruit drop, growth, yield of Indian ber with maximum fruit set and retention as well as yield attributing characters such as size of fruit (length and diameter), weight and volume of fruit which ultimately increased the yield per plant and thereby per hectare in both NAA @40ppm and 2,4,5-T@30ppm.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Vavilov NI. The origin, variation, immunity and breeding of cultivate crops. Chronic Botanica, Waltham Mass, U.S.A. (Translation from Russian selected writings); 1951.
- Bakhshi JC, Singh P. Ber: A good choice for semi-arid and marginal soils. Indian Horticulture; 1974.
- Olsen SR, Cole CV, Watanabe FS, Dean LA. Estimation of available phosphorus in soil by extraction with sodium bicarbonate US Dept. Agric. Circ. 1954;939.
- Meston EC. Contribution of soil constituents to the cation exchange capacity as determined by different extraction methods. Journal Revista Brasileira de Ciencia do Solo. 1956;33(3):507-515.
- Jackson DI. Gibberellin and the growth of peach and apricot fruits. Australian Journal of Biological Sciences. 1968;21(2):209-216.
- Richards LA. Diagnosis and improvement of saline and alkaline soil. USDA Hand Book no. 60; 1968.
- Lindsey WL, Norwell WA. Determination of zinc. Soil Sci. Soc. Am. J. 1978;42:421.
- Panse VG, Sukhatme PV. Statistical methods for agricultural workers, 4th Edn, New Delhi; 1985.
- Chandel SRS. Principles of experimental designs. A handbook of Agricultural Statistics. 1984:36-43.
- Sandhu SS, Thind SS. Effect of NAA sprayed on fruit set on general appearance and quality of umran ber. Indian Journal of Horticulture. 1988;45:274-282.
- Chaurasiya, Gangadhar R, Sharma A, Tiwari S, Goyal G, Bhaduria AS, Singh AP, Yadav A. Influence of foliar application of GA_3 with and without NAA on fruit drop, growth, yield and quality of ber (*Zizyphus mauritiana* Lamk.). International Journal of Current Microbiology and Applied Sciences. 2019;8(3):45-56.
- Chaudhary, Raj Bala, Bairwa LN, Garhwal OP, Negi P. Effect of plant growth regulators and nutrients on yield attributing characters and yield of ber (*Zizyphus mauritiana* Lamk.) Journal of Pharmacognosy and Phytochemistry. 2020;9(4):1968-1972.
- Das KK, Yadav PK, Bhunia SR, Singh RS. Effect of plant growth regulators on flowering parameters of ber (*Zizyphus mauritiana* Lamk.) International Journal of Current Microbiology and Applied Sciences. 2020;9(3):2684-2690.
- Badal DS, Tripathi VK. Effect of foliar application of NAA and Boron on physico-chemical parameters of winter season guava (*Psidium guajava* L.) cv. Lucknow-49. The Pharma Innovation Journal. 2021a;10(9):928-932.
- Saraswat NK, Panday UN, Tripathi VK. Influence of NAA and zinc sulphate on fruit set, fruit drop, cracking, fruit size, yield and

- quality of litchi cv. Culcuttia. J Asian Horticulture. 2006;2(4):255-259.
16. Pandey V. Effect of NAA and GA₃ spray on fruit retention, growth, yield and quality of ber (*Zizyphus mauritiana* Lamk.) cv. Banarasi Karaka. Orissa J. Hort. 1999;27(10):69-73.
 17. Bhat SK, Raina BL, Chogtu SK, Muthoo AK. Effect of exogenous auxin application on fruit drop and cracking in litchi (*Litchi chinensis* Sonn.) cv. Dehradun. Advances-in-Plant Sciences. 1997;10(1):83-86.
 18. Maurya AN, Singh SN, AR. Effect of plants growth regulator on the fruit retention and quality of dashehari mango. Punjab Hort. J. 1973;13(2/3):117-21.
 19. Kumar S, Kumar U, Naresh P. Effect of plant growth regulators on yield of Indian ber (*Zizyphus mauritiana* L.) fruit. Int. J. Pure App. Biosci. 2017;5(1):966-969.
 20. Singh, Sharma. Advancement and efficacy of plant growth regulators in ber (*Zizyphus mauritiana* Lamk). A review Journal of Applied and Natural Science. 2020;12(3):372-379.
 21. Bankar GS, Prasad RN. Effect of gibberellic acid and NAA on fruit set and quality of fruit in ber cv. gola. Progressive Horticulture. 1990;22(1-4):60-62.
 22. Ghosh SN, Bera B, Kundu A, Roy S. Effect of plant growth regulators on fruit retention, yield and physico-chemical characteristics of fruits in Ber cv. Banarasi Karaka grown in close spacing. Proceedings of 1st International Jujube Symposium, Agricultural University of Hebei, Baoding, China. 2008:18.
 23. Singh C, Bal JS. Effect of nutrients and growth regulators on physico-chemical characteristics of Indian jujube (*Zizyphus mauritiana* Lamk.). Proceedings of 1st International Jujube Symposium, Agricultural University of Hebei, Baoding, China; 2008:49.
 24. Chauhan AS, Kumar K, Saini PK, Singh V, Singh JP. Effect of NAA and zinc sulphate on fruiting, yield of litchi (*Litchi chinensis* Sonn.) cv. Calcuttia. International Journal of Current Microbiology and Applied Sciences. 2019;8(3):836-843.
 25. Deepa L, Tripathi VK., Nayyer Md. Abu, Kumar Sanjeev, Ahmed M, Siddiqui MW. Pre-harvest spray of gibberellic acid, NAA, and calcium nitrate on fruit retention, yield and quality of kinnow mandarin. Environment and Ecology. 2016;34(4C): 2288-2292.
 26. Tiwari P, Tripathi VK, Singh A. Effect of foliar application of plant bio-regulators and micronutrients on fruit retention, yield and quality attributes of aonla. Progressive Research - An International Journal. 2017;12(Special-IV):2565-2568.
 27. Badal DS, Tripathi VK. Influence of foliar feeding of NAA and boron on growth, flowering, fruiting, and yield of winter season guava (*Psidium guajava* L.) cv. L-49. Biological Forum-An International Journal. 2021a;13(3):87-391.
 28. Tripathi VK, Viveka Nand. Effect of foliar application of boron, zinc and NAA on fruit retention, yield and quality attributes of aonla. Progressive Horticulture. 2022;54(1):76-81.
 29. Pandey A, Tripathi VK, Pandey M, Mishra AN, Kumar D. Influence of NAA, GA₃ and zinc sulphate on fruit drop, growth, yield and quality of ber cv. Banarasi Karaka. In Proceedings of the International Symposium on Minor Fruits and Medicinal Plants for Health and Ecological Security (ISMF & MP), West Bengal, India, 19-22 December. 2011:184-187.
 30. Naseem S, Malik Mohsin A, Noor-ul-Nisa M, Muhammad Afzal J. Comparative evaluation of naphthalene acetic acid and urea for preventing premature fruit drop and improving fruit yield and quality in ber cv. Suffon. J. of Agri. Research. 2016;54(1):55-62.
 31. Haidry GA, Jalauddin B, Ghaffoor A, Munir M. Effect of naphthalene acetic acid (NAA) on the fruit drop, yield and quality of mango (*Mangifera indica* L.) cultivar langra. Scientific-Khyber Pakistan. 1997;10(1):13-20.
 32. Deepa L, Tripathi VK, Kumar Sanjeev, Nayyer Md. Abu. Effect of pre-harvest application of gibberellic acid, NAA, and calcium nitrate on fruit drop. Maturity and Storage Quality of Kinnow Mandarin. Res. Environ Life Sci. 2015;8(4):561-564.
 33. Meena V, Eyarkai N, Kashyap P, Meena K K. Naphthalene acetic acid and ferrous sulphate induced changes in physico-chemical composition and shelf-life of ber. Indian Journal of Horticulture. 2013;70(1):37-42.
 34. Arora R, Singh S. Effect of growth regulators on quality of ber (*Zizyphus mauritiana* L.) cv. umran. Agricultural Science Digest-A Research Journal. 2014;34(2):102-106.

35. Rathod RK, Ramdevputra MV, Jadeja SR, Parmar LS, Jivani LL. Effect of foliar application of micronutrients and growth regulator on fruit yield of aonla (*Emblica officinalis*). Journal of Pharmacognosy and Phytochemistry. 2019;8(5):133-137.
36. Patil AS, Tidke SN, Tike MA, Shinde BN, Gore AK. Effect of chemicals and growth regulators on physical and chemical characters of parbhani bhushan mango. Journal of Soils and Crops. 2005;15(1):76-79.
37. Kumar R, Tripathi VK, Tomar S, Chaudhary M. Effect of best plant bio-regulators and micronutrient for achieving higher yield and quality of mango (*Mangifera indica* L.) fruits cv. amrapali. Journal of Plant Development Sciences. 2018;10(11):599-604.
38. Randhawa GS, Singh JP, Dhuria HS. Effect of gibberellic Acid, 2,4-dichlorophenoxyacetic acid and 2,4,5-trichlorophenoxyacetic acid on fruit set, drop, size and total yield in sweet lime (*Citrus Limettioides* Tanaka.). Indian Journal of Horticulture. 1959;16(4):206-209.
39. Tripathi D, Pandey AK, Pal AK, Yadav MP. Studies on effect of plant growth regulators on fruit drop, development, quality and yield of ber (*Zizyphus mauritiana* Lamk.) cv. Banarasi Karaka. Progressive Horticulture. 2009;41(2):184-186.
40. Brahmachari VS, Mandal AK, Kumar R, Rani R. Effect of growth substances on flowering and fruiting characters of sardar guava (*Psidium guajava* L.). Horticulture Journal. 1996;(9):1-7.
41. Suman M, Sangma PD, Meghawal DR, Sahu OP. Effect of plant growth regulators on fruit crops. Journal of Pharmacognosy and Phytochemistry. 2017;6(2): 331-337.
42. Singh UR, Singh N. Effect of plant regulators on fruit drop, size and quality of ber (*Zizyphus mauritiana* Lamk.) var. Banarasi. Haryana Journal of Horticultural Sciences. 1976;(5):1-8.
43. Singh K, Randhawa JS, Singh K. Effect of growth regulators and fungicides on fruit drop, yield and quality of fruit in ber cv. umran. J. Res. PAU. 2001;38(3-4):181-185.
44. Bal JS, Singh SN, Randhawa JS, Sharma SC. Effect of naphthalene acetic acid and tri-chlorophenoxy acetic acid on fruit drop, size and quality of ber. Progressive Horticulture. 1982;14(2-3):148-151.
45. Bal JS, Singh SN, Randhawa JS. Response of naphthalene acetic acid spray at fruit set and slow growth phase in ber fruits (*Zizyphus mauritiana* Lamk.). Punjab Agriculture University Journal of Research. 1986;(23):569-572.
46. Singh NP, Malhi CS, Sharma RC. Effect of Plant Bio Regulators (PBRs) on flowering, fruit yield and quality in mango cv. dashehari. Horticultural Journal. 2005;18:10-12.
47. Bal JS, Singh SN, Randhawa JS, Jawanda JS. Effect of growth regulators on fruit drop, size and quality of ber (*Zizyphus mauritiana* Lamk.). Indian Journal of Horticulture. 1984;41(3/4):182-185.
48. Ram RB, Pandey S, Kumar A. Effect of plant growth regulators on fruit retention, physico-chemical parameters and yield of ber cv. Banarasi karaka. Biochemical and cellular Archive. 2005;5(2):229-232.
49. Kumar R, Ram D, Kumar A, Kumar R, Ojha P, Dayal V. Effect of micro nutrients and plant growth regulator on fruit setting of *Psidium guajava* L. cv. Lucknow-49. The Pharma Innovation Journal. 2022;11(11):967-969.
50. Kumar R, Tripathi VK. Influence of NAA, GA₃ and boric acid on growth, yield and quality of strawberry cv. chandler. Progressive Horticulture. 2009;41(1):113-115.
51. Kumar B, Mistry NC, Singh B, Gandhi CP. Indian horticulture database. National Horticulture Board. Ministry of Agriculture, Government of India; 2011.
52. Piper CS. Soil and plant analysis. LWW; 1945.
53. Doran JW, Jones A. Methods for assessing soil quality. Soil Science Society of America; 1996.

© 2023 Kumar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/98443>